arXiv:1304.0562v1 [math.PR] 2 Apr 2013

Speed and fluctuations of N-particle branching Brownian motion with spatial selection

Pascal Maillard*

April 3, 2013

Abstract

We consider branching Brownian motion on the real line with the following selection mechanism: Every time the number of particles exceeds a (large) given number N, only the N right-most particles are kept and the others killed. After rescaling time by $\log^3 N$, we show that the properly recentred position of the $\lfloor \alpha N \rfloor$-th particle from the left, $\alpha \in (0,1)$, converges in law to an explicitly given spectrally positive Lévy process. This behaviour has been predicted to hold for a large class of models falling into the universality class of the FKPP equation with weak multiplicative noise [Brunet et al., Phys. Rev. E **73**(5), 056126 (2006)] and is proven here for the first time for such a model.

Keywords. Branching Brownian motion; selection; FKPP equation

MSC2010. 60J80; 60K35; 60J70

Contents

1 Introduction **2**
1.1 Definition of the model and statement of the main result 2
1.2 Motivation and related work 3
1.3 Overview of the proof 4
1.4 Notation guide 8

2 Brownian motion in an interval **9**
2.1 Brownian motion killed upon exiting an interval 10
2.2 The Brownian taboo process 11

3 Preliminaries on branching Markov processes **13**
3.1 Definition and notation 13
3.2 Stopping lines 14
3.3 Many-to-few lemmas and spines 15

4 BBM with absorption at a critical line **17**
4.1 Proof of Proposition 4.1 18

---

[*] Department of Mathematics and Computer Science, Weizmann Institute of Science, POB 26, Rehovot 76100, Israel. e-mail: pascal DOT maillard AT weizmann DOT ac DOT il

5 BBM in an interval **20**
5.1 Notation 20
5.2 The processes Z_t and Y_t 21
5.3 The number of particles 23
5.4 The particles hitting the right border 25
5.5 Penalizing the particles hitting the right border 26

6 BBM with absorption before a breakout **28**
6.1 Definitions 29
6.2 The time of the first breakout 32
6.3 The particles that do not participate in the breakout 35
6.4 The fugitive and its family 39

7 The B-BBM **42**
7.1 Definition of the model 42
7.2 Proof of Proposition 7.3 44
7.3 Proof of Theorems 7.1 and 7.2 48

8 The $\mathrm{B}^\flat$-BBM **50**
8.1 Definition of the model 51
8.2 Preparatory lemmas 52
8.3 The probability of $G_1^\flat$ 56
8.4 Proofs of the main results 61

9 The $\mathrm{B}^\sharp$-BBM **63**
9.1 Definition of the model 63
9.2 Preparatory lemmas 64
9.3 The probability of $G_1^\sharp$ 66
9.4 Proofs of the main results 72

10 Coupling the N-BBM with other braching Brownian motions **74**
10.1 A monotone coupling between N-BBM and more general particle systems 74
10.2 Proof of Theorem 1.1 76

Acknowledgements **77**

1 Introduction

1.1 Definition of the model and statement of the main result

In this article, we consider an instance of branching Brownian motion (BBM) with *selection*, dubbed the N-BBM and defined as follows: Given a reproduction law $(q(k))_{k\geqslant 0}$ with $m = \sum(k-1)q(k) > 0$ and finite second moment, particles diffuse according to standard Brownian motion and branch at rate $q(k)/(2m)$ into k particles, for every $k \geqslant 0$. We fix a (large) parameter N and select particles according to the following simple mechanism: Each time the number of particles exceeds N, we keep only the N right-most and instantaneously kill the others. In Section 1.2, we explain the motivation behind this system. We first state our result.

For a finite counting measure ν on $\mathbf{R}$, $\alpha \in (0,1)$ and $N \in \mathbf{N}$, we define

$$\mathrm{qu}_\alpha^N(\nu) = \inf\{x \in \mathbf{R} : \nu([x,\infty)) < \alpha N\}.$$

For $N \in \mathbf{N}$ large enough, we then define $a_N = \log N + 3\log\log N$ and

$$\mu_N = \sqrt{1 - \frac{\pi^2}{a_N^2}} = 1 - \frac{\pi^2}{2\log^2 N} + \frac{3\pi^2 \log\log N}{\log^3 N} + o\Big(\frac{1}{\log^3 N}\Big).$$

We then set $M_\alpha^N(t) = \mathrm{qu}_\alpha^N(\nu_t^N) - \mu_N t$, where ν_t^N is the counting measure formed by the positions of the particles of N-BBM at time t. Our main theorem is the following:

Theorem 1.1. *Suppose that at time* 0 *there are N particles distributed independently according to the density proportional to* $\sin(\pi x/a_N)e^{-x}\mathbf{1}_{(0,a_N)}(x)$. *Let $\alpha \in (0,1)$ and define x_α by* $\int_{x_\alpha}^\infty ye^{-y}\,\mathrm{d}y = \alpha$. *Then the finite-dimensional distributions of the process* $\big(M_\alpha^N(t\log^3 N)\big)_{t\geqslant 0}$ *converge weakly as $N \to \infty$ to those of the Lévy process $(L_t + x_\alpha)_{t\geqslant 0}$ with $L_0 = 0$ and*

$$\log E[e^{i\lambda L_1}] = i\lambda c + \pi^2 \int_0^\infty e^{i\lambda x} - 1 - i\lambda x \mathbf{1}_{(x\leqslant 1)}\,\Lambda(\mathrm{d}x). \tag{1.1}$$

Here, Λ is the image of the measure $(x^{-2}\mathbf{1}_{(x>0)})\mathrm{d}x$ *by the map $x \mapsto \log(1+x)$ and $c \in \mathbf{R}$ is a constant depending on the reproduction law $q(k)$.*

1.2 Motivation and related work

The N-BBM and the results of this article are related to several other mathematical models, a few of which we wish to outline in this section. We are not going to elaborate on the applications to other sciences such as physics or biology, for which we refer to [17, 19] and the review articles [56, 52].

Most importantly, the N-BBM is a prototype for *noisy traveling waves of FKPP type*, i.e. travelling waves of the stochastic partial differential equation $\partial_t u = \partial_x^2 u + u(1-u) + \sqrt{u/N}\,\dot{W}$, where $\dot{W}$ is space-time white noise. This equation is known as the FKPP equation with weak multiplicative noise and is believed to share the same phenomenology as the N-BBM. The connection between BBM *without* selection and the FKPP equation *without* noise is indeed well known, at least since McKean's work [48], and has had many fruitful applications, one of the most important being Bramson's study of the law of the right-most particle of BBM [14]. The relation between the N-BBM and the noisy FKPP equation is less explicit, but there is an exact duality relation between this equation and a system of branching and coalescing Brownian motions [55]. More generally, particle systems governed by the FKPP equation in the large population limit can often be modeled by the noisy FKPP equation, the parameter N having the meaning of an "effective population size" ([16], [56, Chapter 7], [52]).

Brunet, Derrida, Mueller and Munier [17] give a semi-deterministic description of such systems and basically obtain our Theorem 1.1. Their arguments, which we outline in Section 1.3, work for very general models but rely on some fundamental assumptions which are intuitively convincing but far from easy to prove in particular settings. The author is indeed aware of only three mathematically rigorous articles in this area: Bérard and Gouéré [6] prove the $\log^{-2} N$ correction to the linear speed of the N-particle branching random walk (N-BRW). Durrett and Remenik [27] study the empirical distribution of N-BRW and show that it converges to a system of integro-differential equations with moving boundary. Finally, Mueller, Mytnik and Quastel [49] obtain an estimate for the speed of the traveling waves of the noisy FKPP equation, which partly confirms the physicists' predictions. To our knowledge, the present article is therefore the first example of a rigorous proof of the full statistics for a particular system.

In order to prove Theorem 1.1, we approximate the N-BBM by BBM with absorption at a linear space-time barrier. This process is well-studied [26, 30, 32, 39, 51] and is much more tractable than N-BBM due to the greater independence between the particles and its connection with some differential equations [32, 46, 51]. Of paramount importance to us is the article by Berestycki, Berestycki and Schweinsberg, who study BBM with absorption at a barrier with slope μ_N, starting from N particles. If N_t denotes the number of particles at time t in their system, they show that the process $(N_t)_{t\geqslant 0}$ converges in law to Neveu's continuous-state branching process and the genealogy of the particles to the Bolthausen–Sznitman coalescent, which has been observed before [18]. Although we do not apply their results directly, our proof of Theorem 1.1 uses many of their ideas. We remark that the genealogy of the N-BBM should converge to the Bolthausen–Sznitman coalescent as well, but our methods do not seem to be applicable (see the end of Section 1.3 for details).

We finally mention the relation between the N-BBM or N-BRW and an instance of a Fleming–Viot-type process: Let N particles perform independent continuous-time nearest-neighbour random walks on the integers with drift towards the origin. Furthermore, as soon as a particle reaches the origin, let it jump at random onto one of the other particles. This type of model was introduced in order to yield a particle picture of quasi-stationary distributions [21], and it is indeed conjectured that the empirical distribution of this system under the stationary distribution converges, as N goes to infinity, to the quasi-stationary distribution of the random walk (see [2, 3] for recent progress on this subject). The relation with the N-BBM is obvious and we conjecture that the empirical distribution of a general N-BRW is an ergodic process whose stationary distribution converges to a deterministic measure, which is the quasi-stationary distribution of the underlying random walk with a certain drift towards the origin.

1.3 Overview of the proof

Since this article is fairly long, we give a detailed overview of our main results. We recall the heuristic semi-deterministic description of N-BBM established in [17]:

1. Most of the time, the particles are in a meta-stable state. In this state, the diameter of the cloud of particles (also called the *front*) is approximately $\log N$, the empirical density of the particles proportional to $e^{-x}\sin(\pi x/\log N)$, and the system moves at a linear speed $v_{\mathrm{cutoff}} = 1-\pi^2/(2\log^2 N)$. This is the description provided by the *cutoff* approximation from [15] mentioned above.

2. This meta-stable state is perturbed from time to time by particles moving far to the right and thus spawning a large number of descendants, causing a shift of the front to the right after a relaxation time which is of the order of $\log^2 N$. To make this precise, we fix a point in the bulk, for example the barycentre of the cloud of particles, and shift our coordinate system such that this point becomes its origin. Playing with the initial conditions of the FKPP equation with cutoff, the authors of [17] find that a particle moving up to the point $\log N + x$ causes a shift of the front by $\Delta = \log\left(1+\frac{Ce^x}{\log^3 N}\right)$, for some constant $C > 0$. In particular, in order to have an effect on the position of the front, a particle has to reach a point near $\log N + 3\log\log N$.

3. Assuming that such an event where a particle "escapes" to the point $\log N + x$ happens with rate Ce^{-x}, one sees that the time it takes for a particle to come close to $\log N + 3\log\log N$ (and thus causing shifts of the front) is of the order of $\log^3 N \gg \log^2 N$.

4. With this information, the speed of the front is found to satisfy $v_N - v_{\mathrm{cutoff}} \approx \pi^2 \frac{3\log\log N}{\log^3 N}$ and the fluctuations are found to be given by the Lévy process from Theorem 1.1.

Berestycki, Berestycki and Schweinsberg [7] made a first attempt to put this description onto a rigorous foundation. They consider BBM with absorption at the origin and with drift $-\mu_N$. Their starting point is to introduce a second barrier at the point $a_{N,A} = a_N - A$ for some large positive constant A and divide the particles at time t into two parts; on the one hand those that have stayed inside the interval $(0, a_{N,A})$, on the other hand those that have hit the point $a_{N,A}$ before hitting 0.

This corresponds roughly to the division of the process into a deterministic and a stochastic part. Indeed, killing the particles at $a_{N,A}$ prevents the number of particles from growing fast and thus permits to calculate expectations and variances of various quantities. For example, if at time 0 we have N particles distributed according to the meta-stable density, then the variance of the number of particles at the time $\log^3 N$ is of order of $e^{-A}N^2$. For large A, the particles inside the interval $(0, a_{N,A})$ therefore behave almost deterministically at the timescale $\log^3 N$. Moreover, the leading term in the Fourier expansion of the transition density of Brownian motion (with drift $-\mu_N$ and killed at the border of the interval $(0, a_{N,A})$) is proportional to $e^{-\mu_N x}\sin(\pi x/a_{N,A})$, which explains the meta-stable density predicted by the physicists.

As for the particles that hit $a_{N,A}$, the authors of [7] find that 1) the number of descendants at a later time of such a particle is of the order of $e^{-A}NW$, where W is a random variable with tail $P(W > x) \sim 1/x$, as $x \to \infty$ and 2) the rate at which particles hit the right barrier is of the order of $e^A/\log^3 N$. Putting the pieces together, they then show that the process which counts the number of particles of the system converges in the $\log^3 N$ timescale to Neveu's continuous-state branching process[1] and its genealogy to the Bolthausen–Sznitman coalescent.

Our proof of Theorem 1.1 builds upon the ideas of [7] presented above. The basic idea is to approximate the N-BBM by a BBM with absorption at a *random* barrier, which is chosen in such a way that *it keeps the number of particles almost constant.* We call the resulting system the "B-BBM" (B stands for "barrier"), see Figure 1 for a graphical description. The B-BBM takes two (large) parameters a and A, which have similar purposes than $a_{N,A}$ and A above. We then set $\mu = \sqrt{1 - \pi^2/a^2}$ and start with BBM with drift $-\mu$ and an absorbing barrier at the origin. Now, at the beginning, this barrier stays at the origin and does not move. When and only when a particle hits a *and* spawns a lot of descendants do we increase the drift to the left. This increase is in order to kill particles and thus make the population size stay almost constant. Note that moving the barrier to the right is an equivalent operation, but increasing the drift is technically more convenient. After the system has relaxed (which takes a time of order a^2) the drift is set to $-\mu$ again and the process is repeated.

As in [7], an important quantity is $Z_t = \sum w_Z(X_u(t))$, where $w_Z(x) = ae^{\mu(x-a)}\sin\frac{\pi x}{a}$. Here, we sum over all the particles u alive at time t and $X_u(t)$ denotes the position of the particle u at time t. The process $(Z_t)_{t\geqslant 0}$ is important because it is a martingale for BBM with absorption at 0 and a and drift $-\mu$. Furthermore, it gives the approximate number of particles at a time $t \gg a^2$. More precisely, set $N = \lceil 2\pi e^A a^{-3} e^{\mu a}\rceil$ and suppose we kill particles at 0 and a. The expected number of particles at a time t, with $a^2 \ll t \ll a^3$, is then approximately $Ne^{-A}Z_0$. Moreover, the variance is of the order of $N^2 e^{-2A} Z_0$. Therefore, if $Z_0 \approx e^A$ then the number of particles is concentrated around its expectation for large A.

When a particle hits the right barrier at the time τ, say, we absorb its descendants at the space-time line $\mathscr{L}_\tau : x = a - y + (1-\mu)(t-\tau)$, where y is a large constant depending on A only (this idea comes from [7]). In doing so, the number of particles absorbed at the barrier has the same law as in BBM with drift -1 and absorption at $-y$, starting from a single particle at the origin. Moreover, the time it takes for all the particles to be absorbed depends only on y, not on a. Now, if $x_1, \dots, x_n$ are the positions of the absorbed particles and $Z' = \sum_i w_Z(x_i)$, then the number of descendants of this particle at a later time is of the order of $e^{-A}NZ'$, provided that the drift stays constant. Consequently, we say that a *breakout* occurs, whenever $Z' > \varepsilon e^A$, where $\varepsilon = A^{-p}$ for some large p to be chosen later.

---

[1] A continuous-state branching process (CSBP) $(Z_t)_{t\geqslant 0}$ is a time-changed Lévy process without negative jumps: at time t, time is sped up by the factor Z_{t-}. CSBPs are scaling limits of Galton–Watson processes and thus have an inherent notion of genealogy. Neveu's CSBP is the CSBP with Lévy measure $x^{-2}\mathbf{1}_{(x>0)}\,\mathrm{d}x$, whose genealogy is given by the Bolthausen–Sznitman coalescent [8]. As a wise reader, you have read the introductory chapter, such that you know what this coalescent is.

In order to define a breakout properly, we classify the particles into *tiers*. Particles that have never hit the point a form the particles of tier 0. As soon as a particle hits a (at the time τ, say) it advances to tier 1. Its descendants then belong to tier 1 as well, but whenever a descendant hits a and has an ancestor which has hit the line $\mathscr{L}_\tau$ after τ, it advances to tier 2 and so on. Whenever a particle advances to the next tier, it has a chance to break out. We can then define *the time T of the first breakout* and will indeed show that T is approximately exponentially distributed with rate proportional to $\varepsilon^{-1}a^{-3}$. Interestingly, we will see that with high probability breakouts only occur from particles which are of tier 0 or 1. In fact, the number of breakouts occuring from particles of tier 1 between the times 0 and a^3 is approximately proportional to A (and the remaining $\approx \varepsilon^{-1}$ breakouts occur from particles of tier 0).



Figure 1: A caricatural graphical description of the B-BBM. The fugitive and its descendants are drawn with thick lines, the other particles with thin lines. A breakout happens at time T and the barrier is moved from the time T^+ on. The process starts afresh at the time Θ_1. Note that technically we increase the drift to the left instead of moving the barrier. The three important timescales (1, a^2 and a^3) are shown as well. Recall that $a \approx \log N$.

After the breakout, we will then increase the drift to the left slightly, in order to kill more particles than usual (remember that increasing the drift to the left corresponds to moving the barrier to the right). For this, we first choose a family of increasing smooth functions $(f_x)_{x\geqslant 0}$ with $f_x(0) = 0$ and $f_x(+\infty) = x$ for all $x \geqslant 0$. Such a function will be called a *barrier function*. The drift after a breakout is then set to $\mu_t = \mu + (\mathrm{d}/\mathrm{d}t)f_\Delta(t/a^2)$, where Δ is the total amount by which we have to move the barrier. Thus, the only randomness in the choice of the barrier is in its total shift, not in its shape. Looking at the definition of Z_t and the fact that $Z_0 \approx e^A$, one easily guesses that we have to choose $\Delta = \log(1 + e^{-A}Z')$ in order to get Z_t ultimately back to its initial value. This already explains the convergence of the barrier to the Lévy process given by (1.1): On the one hand, we have $Z' \approx \pi W$, where W is the random variable mentioned above. This implies that the law of $e^{-A}Z'$ conditioned on $Z' > \varepsilon e^A$ is approximately $\varepsilon x^{-2}\mathbf{1}_{(x\geqslant\varepsilon)}\,\mathrm{d}x$ for large A and a.[2] On the other hand, we will show that breakouts occur at a rate proportional to $\varepsilon^{-1}a^{-3}$. Together with the definition of Δ, this explains the Lévy measure $\Lambda(\mathrm{d}x)$ in (1.1).

---

[2] The statement "for large A and a" means that we let first a, then A go to infinity, see Section 6.1.

As for the shape of the barrier, it is determined by the fact that we want the number of particles at each time to be approximately N. This is accomplished by choosing

$$f_\Delta(t) = \log\Big(1 + (e^\Delta - 1)\pi^{-2}e^{\pi^2 t/2}\frac{\mathrm{d}}{\mathrm{d}t}\theta(1,t)\Big),$$

where the theta function $\theta(x,t)$ is defined in (2.1).

We remark that in order to study the B-BBM up to the time T of the first breakout, we need to study it *conditioned to break out at time* t for every $t \geqslant 0$. We will see in Section 6.4 that this will lead to a decomposition of the particles into 1) a *fugitive*, which is conditioned to break out at t and which will effectively be a *spine*, and 2) the other particles conditioned *not to break out* before t. This is essentially a Doob transform of the process. Furthermore, we will see that the tier 1 particles will have an essential role: At the timescale a^3 they will lead to an additional shift of the barrier by an amount of the order of A. This term will play the role of the linear compensation that is necessary in order to obtain in the limit the Lévy process of infinite variation stated in Theorem 1.1.

We now describe how we use the results on the B-BBM in order to prove Theorem 1.1. Initially, our plan was to *couple* the N-BBM and the B-BBM, i.e. construct them on the same probability space. We would then assign a colour to each particle: *blue* to the particles which appear in the N-BBM but not in the B-BBM, *red* to those that appear in the B-BBM but not in the N-BBM, and *white* to the particles that appear in both processes. Our aim was then to show that the number of blue and red particles was negligible after a time of order a^3. This, unfortunately, did not work out, because we were not able to handle the intricate dependence between the red and blue particles.

Instead, we couple the N-BBM with two different processes, the $\mathrm{B}^\flat$- and the $\mathrm{B}^\sharp$-BBM, which are variants of the B-BBM and which bound the position of the N-BBM in a certain sense from below and above, respectively. The $\mathrm{B}^\flat$-BBM is defined as follows: Initially, all particles are coloured white and evolve as in the B-BBM. A white particle is coloured red as soon as it has N or more white particles to its right. Children inherit the colour of their parent. After a breakout and the subsequent relaxation, all the red particles are killed immediately and the process restarts with the remaining particles. It is intuitive that the collection of white particles then bounds the N-BBM from below (in some sense) because we kill "more" particles than in the N-BBM. Indeed, in Section 10.1, we show by a coupling method that the empirical measure of the white particles in $\mathrm{B}^\flat$-BBM is stochastically dominated by the one of the N-BBM with respect to the usual stochastic ordering of measures. It then remains to show that the number of red particles in $\mathrm{B}^\flat$-BBM is negligible when A and a are large. We do this through precise estimates on the number of particles in the interval $[r,\infty)$ for every $r \geqslant 0$. These allow us to estimate the expected number of particles which turn red at the point r. It turns out that this expectation is small enough, which permits to conclude.

The definition of the $\mathrm{B}^\sharp$-BBM, which is used to bound the N-BBM from above, is more intricate than the one of $\mathrm{B}^\flat$-BBM. Again, we colour all initial particles white and particles evolve as in B-BBM with the following change: Whenever a white particle hits 0 *and* has less than N particles to its right, instead of killing it immediately, we colour it blue and let it survive for a time of order a^2. More precisely, we cut time into intervals $I_n = [t_n, t_{n+1})$, with $t_n = Kna^2$ for some large constant K. A particle which gets coloured blue during I_n then survives until the time t_{n+2}. At this time, all of its descendants to the left of the origin are killed and the others survive. It will turn out that this system bounds the N-BBM from above with high probability and that the number of blue particles will remain negligible during a time of order a^3, as long as A and a are large.

We note that although our technique of bounding the N-BBM from below and from above works well for the *position* of the particles, it does not give us information about the genealogy; the reason being that the coupling deforms the genealogical tree of the process. Thus, although it should not be difficult to show that the genealogy of the B-BBM (and of the $\mathrm{B}^\flat$- and $\mathrm{B}^\sharp$-BBM) converges to the

Bolthausen–Sznitman coalescent we do not know at present how one could transfer this information to the N-BBM.

1.4 Notation guide

This article is quite long and uses a wealth of different notation. Below is a list of recurrent symbols, roughly in the order of their first appearance. Following this list are some further remarks about notational conventions.

Symbol	**Meaning**	**Sect.**
$q(k)$	Reproduction law	1, 4
m	$m = \sum_k (k-1)q(k)$	1, 4
β_0	Branching rate, $\beta_0 = 1/(2m)$	1, 4
$\theta, \overline{\theta}$	Theta functions	2.1
W^x	Law of Brownian motion started at x	2.1
$p_t^a(x,y)$	Transition density of Brownian motion killed outside $[0,a]$	2.1
E_t	Error term	2.1
$I^a(x,S), J^a(x,S)$	Integrals related to Brownian motion killed outside $[0,a]$	2.1
$W^x_{\text{taboo}}, W^{x,t,y}_{\text{taboo}}$	Law of Brownian taboo process and its bridge	2.2
$\mathbf{N}$, $\mathbf{N}^*$	The set of natural numbers including and excluding 0, resp.	3.1
U	The space of individuals	3.1
Ξ	A realisation of a branching Markov process	3.1
ζ_u, b_u, d_u	Lifetime, birth and death times of an individual u	3.1
$\mathscr{N}(t)$	Set of individuals alive at time t	3.1
$X_u(t)$	Position of the individual u at the time t	3.1
$\mathbf{P}^x, \mathbf{E}^x, \mathbf{P}^\nu, \mathbf{E}^\nu$	Law of and expectation w.r.t. a branching Markov process (later: BBM) started at x or with particles distributed according to a finite counting measure ν	3.1
$\mathscr{F}_t$	σ-algebra with information up to time t	3.2
$\mathscr{F}_{\mathscr{L}}$	σ-algebra with information up to the stopping line $\mathscr{L}$	3.2
$\mathscr{L}_T$	Stopping line generated by a stopping time T	3.2
L	A random variable with law $q(k)$	4
m_2	$m_2 = E[L(L-1)]$	4
W	Seneta-Heyde martingale limit of BBM with absorption	4
μ	$\mu = \sqrt{1-\pi^2/a^2}$	5.1
$\mathbf{p}_t(x,y)$	Density of BBM killed outside an interval	5.1
$\mathbf{P}_f, \mathbf{E}_f$	Law of and expectation w.r.t. BBM with varying drift	5.1
$w_Z(x)$	$w_Z(x) = a\sin(\pi x/a)e^{\mu(x-a)}\mathbf{1}_{(x\in[0,a])}$	5.2
$w_Y(x)$	$w_Y(x) = e^{\mu(x-a)}\mathbf{1}_{(x\geqslant 0)}$	5.2
Z_t, Y_t	Sums of $w_Z(x), w_Y(x)$ over the positions of time t particles	5.2
N_t	Number of particles at time t	5.3
R_t	Number of particles hitting a up to time t	5.4
$\widetilde{\mathbf{P}}, \widetilde{\mathbf{E}}$	Law/expectation of BBM weakly conditioned not to hit a	5.5
$A, a, \varepsilon, \eta, y, \zeta$	Parameters of BBM before a breakout and B-BBM	6.1
$\mathscr{N}_t^{(l)}$	Stopping line of tier l particles at time t	6.1
$\mathscr{S}_t^{(l)}$	Stopping line of tier l particles hitting the critical line before t	6.1
$\mathscr{R}_t^{(l)}$	Stopping line of tier l particles hitting a before t	6.1
$\mathscr{S}^{(u,t)}$	Stopping line of descendants of (u,t) hitting the critical line	6.1
$Z_t^{(l)}$	Sum of $w_Z(x)$ over particles from $\mathscr{N}_t^{(l)}$	6.1

Z_t, Y_t, R_t, N_t	$Z_t = Z_t^{(0+)}$ (from Section 6 on). Same for Y_t, R_t and N_t	6.1
p_B	Probability of a breakout	6.1
$\mathbf{Q}^a$	$\mathbf{P}^a$ conditioned not to break out	6.1
$T^{(l)}$	Time of first breakout from tier l	6.1
$\widehat{\mathbf{P}}, \widehat{\mathbf{E}}$ (also $\widehat{\mathbf{P}}_\lambda$,$\widehat{\mathbf{E}}_\lambda$)	Law/expectation of BBM conditioned not to break out before some fixed time	6.3
$Z_{\varnothing}^{(l)}$	Sum of $w_Z(x)$ over particles from $\mathscr{S}_t^{(l)}$	6.3
$\mathscr{U}$	The fugitive (the particle which breaks out)	6.4
$\overline{\mathscr{N}_t}, \breve{\mathscr{N}_t}$	The descendants of the fugitive at time t (split into two groups)	6.4
$\bar{Z}_t$, $\breve{Z}_t$	Value of Z_t restricted to particles from $\overline{\mathscr{N}_t}$ and $\breve{\mathscr{N}_t}$, resp.	6.4
$\widehat{Z}_t, \widehat{Y}_t, \widehat{N}_t, \dots$	Value of $Z_t, Y_t, N_t \dots$ restricted to particles not related to the fugitive	7
$Z_t^{\mathrm{fug}}, Y_t^{\mathrm{fug}}, N_t^{\mathrm{fug}}, \dots$	Value of $Z_t, Y_t, N_t \dots$ restricted to particles descending from the fugitive	7
T, T_n	Time of the first/n-th breakout	7.1
T^-, T_n^-, T^+, T_n^+	Random times related to the times T, T_n	7.1
Δ	The total shift of the barrier after a breakout	7.1
$X_t^{[n]}$	The position of the first n pieces of the barrier at time t (barrier process)	7.1
Θ_n	Beginning of the $n+1$-th piece of B-BBM (i.e. time of the n-th breakout plus relaxation time)	7.1
G_n	"Good event" related to the first n pieces of B-BBM	7.1
γ_0	$\gamma_0 = (\pi p_B e^A)^{-1}$.	7.1
$G_{\mathscr{U}}, G_{\mathrm{fug}}, \widehat{G}$, $\breve{G}, G_\Delta, G_{\mathrm{nbab}}$	Several good sets related to a piece of B-BBM	7.2

The complement of an event G is denoted by G^c.

From Section 5 on, the symbol C stands for a *positive* constant, which may only depend on the reproduction law q and the value of which may change from line to line. Furthermore, if X is any mathematical expression, then the symbol $O(X)$ stands for a possibly random term whose absolute value is bounded by $C|X|$.

In Section 6, we introduce two parameters A and a and will first let a then A go to infinity. This will be expressed by the statements "for large A and a we have…" or "as A and a go to infinity…" (see Section 6.1 for a precise definition). These phrases will become so common that in Sections 7 to 9 they will often be used implicitly, although they will always be explicitly stated in the theorems, propositions, lemmas etc. Section 6 furthermore introduces the notation $o(1)$, which stands for a (non-random) term that only depends on the reproduction law q and the parameters A, a, ε, η, y and ζ and which goes to 0 as A and a go to infinity.

Sections 8 and 9 each use special notation which only appears in those sections. This notation is defined at the beginning of both sections. Moreover, in both sections, we sometimes denote quantities which refer to descendants of the fugitive *after* a breakout by the superscript "fug".

2 Brownian motion in an interval

In this section, we recall some explicit formulae concerning real-valued Brownian motion killed upon exiting or conditioned not to exit an interval.

2.1 Brownian motion killed upon exiting an interval

We define for $x \in \mathbf{R}$ and $t > 0$ the following function of Jacobi theta-type:

$$\begin{aligned} \theta(x,t) &= \tfrac{1}{2} + \textstyle\sum_{n=1}^{\infty} e^{-\frac{\pi^2}{2}n^2 t}\cos(\pi n x) &(2.1)\\ &= \textstyle\sum_{n\in\mathbf{Z}} \frac{1}{\sqrt{2\pi t}}\exp\Big(-\frac{(x-2n)^2}{2t}\Big), &(2.2)\end{aligned}$$

the two representations being related by the Poisson summation formula (see [5], §9). We also define

$$\overline{\theta}(t) = \frac{2}{\pi^2} e^{\frac{\pi^2}{2}t}\frac{\mathrm{d}}{\mathrm{d}t}\theta(1,t),$$

which is a smooth function on $\mathbf{R}_+$. By (2.1) and (2.2), one can show that $\overline{\theta}$ is stricly increasing[3] with $\overline{\theta}(0) = 0$ and $\overline{\theta}(+\infty) = 1$.

Various quantities of Brownian motion killed upon exiting an interval can be expressed by $\theta(x,t)$. For $x \in \mathbf{R}$, let W^x be the law of Brownian motion started at x, let $(X_t)_{t\geqslant 0}$ be the canonical process and let $H_y = \inf\{t \geqslant 0 : X_t = y\}$. For $a > 0$ and $x \in [0,a]$, denote by $W^x_{\text{killed},a}$ the law of Brownian motion started at x and killed upon leaving the interval $(0,a)$. Let $p^a_t(x,y)$ be its transition density, i.e.

$$p^a_t(x,y)\mathrm{d}y = W^x_{\text{killed},a}(X_t \in \mathrm{d}y) = W^x(X_t \in \mathrm{d}y,\ H_0 \wedge H_a > t), \quad x,y \in [0,a]. \tag{2.3}$$

Then (see [36], Problem 1.7.8 or [13], formula 1.1.15.8),

$$p^a_t(x,y) = a^{-1}\left(\theta\Big(\frac{x-y}{a},\frac{t}{a^2}\Big) - \theta\Big(\frac{x+y}{a},\frac{t}{a^2}\Big)\right). \tag{2.4}$$

Equation (2.1) then yields

$$p^a_t(x,y) = \frac{2}{a}\sum_{n=1}^{\infty} e^{-\frac{\pi^2}{2a^2}n^2 t}\sin(\pi n\tfrac{x}{a})\sin(\pi n\tfrac{y}{a}). \tag{2.5}$$

This representation is particularly useful for large t: Define

$$E_t = \pi^2\sum_{n=2}^{\infty} n^2 e^{-\pi^2(n^2-1)t/2}. \tag{2.6}$$

By (2.5) and the inequality $|\sin nx| \leqslant n\sin x$, $x \in [0,\pi]$, one sees that

$$\left| e^{\frac{\pi^2}{2a^2}t}p^a_t(x,y) - \frac{2}{a}\sin(\pi x/a)\sin(\pi y/a)\right| \leqslant E_{t/a^2}\frac{2}{a}\sin(\pi x/a)\sin(\pi y/a). \tag{2.7}$$

The Green function (see e.g. [38], Lemma 20.10, p379) is given by

$$\int_0^\infty p^a_t(x,y)\,\mathrm{d}t = 2a^{-1}(x\wedge y)(a - x\vee y). \tag{2.8}$$

Set $H = H_0 \wedge H_a$ and define $r^a_t(x) = W^x(H \in \mathrm{d}t,\ X_H = a)/\mathrm{d}t$. Then (see [13], formula 1.3.0.6),

$$r^a_t(x) = \frac{1}{a^2}\theta'\left(\frac{x}{a} - 1, \frac{t}{a^2}\right), \tag{2.9}$$

where θ' denotes the derivative of θ with respect to x.

---

[3] More precisely, by elementary computations, (2.1) gives $t_0 \in \mathbf{R}_+$, such that $\overline{\theta}$ is strictly increasing on (t_0,∞) and (2.2) gives $t_1 > t_0$, such that $\overline{\theta}$ is strictly increasing on $[0,t_1)$.

The following two integrals are going to appear several times throughout the article, which is why we give some useful estimates here. For a measurable subset $S \subset \mathbf{R}$, define

$$I^a(x,S) = W^x\Big(e^{\frac{\pi^2}{2a^2}H_a}\mathbf{1}_{(H_0>H_a\in S)}\Big) = \int_{S\cap(0,\infty)} e^{\frac{\pi^2}{2a^2}s} r_s^a(x)\mathrm{d}s, \quad \text{and} \tag{2.10}$$

$$J^a(x,y,S) = \int_{S\cap(0,\infty)} e^{\frac{\pi^2}{2a^2}s} p_s^a(x,y)\mathrm{d}s, \tag{2.11}$$

which satisfy the scaling relations

$$I^a(x,S) = I\Big(\frac{x}{a},\frac{S}{a^2}\Big), \quad J^a(x,y,S) = aJ\Big(\frac{x}{a},\frac{y}{a},\frac{S}{a^2}\Big), \tag{2.12}$$

with $I = I^1$ and $J = J^1$. The following lemma provides estimates on $I(x,S)$ and $J(x,y,S)$. It is easily proven from the above equations (see [47, Lemma 2.2.1]).

Lemma 2.1. *There exists a universal constant C, such that for every $x \in [0,1]$ and every measurable $S \subset \mathbf{R}_+$, we have*

$$|I(x,S) - \pi\lambda(S)\sin(\pi x)| \leqslant C\Big(x \wedge E_{\inf S}(1\wedge\lambda(S))\sin(\pi x)\Big), \quad \textit{and}$$

$$|J(x,y,S) - 2\lambda(S)\sin(\pi x)\sin(\pi y)| \leqslant C\Big([(x\wedge y)(1-(x\vee y))] \wedge E_{\inf S}\sin(\pi x)\sin(\pi y)\Big),$$

where $\lambda(S)$ denotes the Lebesgue measure of S and $E_{\inf S}$ is defined in (2.6).

2.2 The Brownian taboo process

The *Brownian taboo process* on the interval $(0,a)$ is the diffusion with infinitesimal generator $\frac{1}{2}\left(\frac{\mathrm{d}}{\mathrm{d}x}\right)^2 + \frac{\pi}{a}\cot\frac{\pi x}{a}\frac{\mathrm{d}}{\mathrm{d}x}$. The name of this process was coined by F. Knight [40] who showed that it can be interpreted as Brownian motion conditioned to stay inside the interval $(0,a)$. It is the Doob transform of Brownian motion killed at 0 and a, with respect to the space-time harmonic function $h(x,t) = \sin(\pi x/a)\exp(\pi^2 t/(2a^2))$. As a consequence, its transition density is given for $x,y \in [0,a]$ by

$$p_t^{\mathrm{taboo}(0,a)}(x,y) = \frac{\sin(\pi y/a)}{\sin(\pi x/a)} e^{\frac{\pi^2}{2a^2}t} p_t^a(x,y) = \frac{2}{a}\sin^2(\pi y/a)(1+O(1)E_{t/a^2}), \tag{2.13}$$

by (2.7). For $x \in [0,a]$ we denote the law of the Brownian taboo process on $(0,a)$ started from x by $W^x_{\mathrm{taboo},a}$. Often we will drop the a if its value is clear from the context. We also denote by $W^{x,t,y}_{\mathrm{taboo}}$ the law of taboo bridge from x to y of length t. Note that the taboo process is self-dual in the sense that for a measurable functional F and $t > 0$, we have $W^{x,t,y}_{\mathrm{taboo}}[F((X_s;0\leqslant s\leqslant t))] = W^{y,t,x}_{\mathrm{taboo}}[F((X_{t-s};0\leqslant s\leqslant t))]$.

The following lemma will be needed in Sections 6 and 7.

Lemma 2.2. *Let $c > 0$. There exists a constant C, depending only on c, such that we have for every $x,y \in [0,a]$,*

$$W^x_{\mathrm{taboo}}\Big[\int_0^t e^{-cX_s}\,\mathrm{d}s\Big] \leqslant C\Big(t/a^3 + (1\wedge x^{-1})\Big), \tag{2.14}$$

and for $t \geqslant a^2$,

$$W^{x,t,y}_{\mathrm{taboo}}\Big[\int_0^t e^{-cX_s}\,\mathrm{d}s\Big] \leqslant C\Big(t/a^3 + (1\wedge x^{-1}) + (1\wedge y^{-1})\Big). \tag{2.15}$$

If $t \leqslant a^2$, we still have,

$$W^{x,t,y}_{\mathrm{taboo}}\Big[\int_0^t e^{-cX_s}\,\mathrm{d}s\Big] \leqslant C. \tag{2.16}$$

Proof. We write $k(x) = e^{-cx}$. We first show that (2.14) implies (2.15). By the self-duality of the taboo process, we have

$$W^{x,t,y}_{\text{taboo}}\Big[\int_0^t k(X_s)\,\mathrm{d}s\Big] = W^{x,t,y}_{\text{taboo}}\Big[\int_0^{t/2} k(X_s)\,\mathrm{d}s\Big] + W^{y,t,x}_{\text{taboo}}\Big[\int_0^{t/2} k(X_s)\,\mathrm{d}s\Big].$$

It therefore remains to prove that

$$E(x,y) = W^{x,t,y}_{\text{taboo}}\Big[\int_0^{t/2} k(X_s)\,\mathrm{d}s\Big] \leqslant C(t/a^3 + (1 \wedge x^{-1})).$$

Conditioning on $\sigma(X_s; 0 \leqslant t \leqslant t/2)$, this integral equals

$$E(x,y) = W^{x}_{\text{taboo}}\Big[\frac{p^{\text{taboo}}_{t/2}(X_{t/2}, y)}{p^{\text{taboo}}_{t}(x,y)}\int_0^{t/2} k(X_s)\,\mathrm{d}s\Big].$$

By (2.13), there exists a universal constant C, such that for $t \geqslant a^2$,

$$E(x,y) \leqslant C\, W^{x}_{\text{taboo}}\Big[\int_0^{t/2} k(X_s)\,\mathrm{d}s\Big].$$

Equation (2.14) therefore implies (2.15).

Heuristically, one can estimate the left side of (2.14) in the following way: Since $k(x)$ is decreasing very fast, only the times at which X_s is of order 1 contribute to the integral. When started from the stationary distribution, the process takes a time of order a^3 to reach a point at distance $O(1)$ from the origin [43] and it stays there for a time of order 1, hence the integral is of order t/a^3. When started from the point x, an additional error is added, which is of order 1, when x is at distance of order 1 away from 0. Adding both terms gives the bound appearing in the statement of the lemma.

The actual calculations can be performed in the following way: Let Y be a random variable with values in $(0,a)$ distributed according to $\widetilde{m}(\mathrm{d}x) := 2/a \sin^2(\pi x/a)\,\mathrm{d}x$, which is the stationary probability measure of the taboo process. Let $H_Y = \inf\{t > 0 : X_s = Y\}$. We then have

$$\begin{aligned} W^{x}_{\text{taboo}}\Big[\int_0^t k(X_s)\,\mathrm{d}s\Big] &= W^{x}_{\text{taboo}}\Big[\int_0^{H_Y} k(X_s)\,\mathrm{d}s + \int_{H_Y}^t k(X_s)\,\mathrm{d}s\Big] \\ &\leqslant W^{x}_{\text{taboo}}\Big[\int_0^{H_Y} k(X_s)\,\mathrm{d}s\Big] + W^{\widetilde{m}}_{\text{taboo}}\Big[\int_0^t k(X_s)\,\mathrm{d}s\Big]. \end{aligned}$$

Both of these quantities can now be expressed through the scale function and speed measure of the taboo process and after some calculations (see [47, Lemma 2.2.2] for details), one obtains (2.14).

In order to prove (2.16), a different method is needed. We may assume that $x, y \leqslant a/2$, otherwise we decompose the path at the first and/or last time it hits $a/2$ and bound the parts above $a/2$ trivially by $a^2 e^{-ca/2} = O(1)$. The transition density of the taboo bridge can be written

$$W^{x,t,y}_{\text{taboo}}\Big[X_s \in \mathrm{d}z\Big] = \frac{p^a_s(x,z)p^a_{t-s}(z,y)}{p^a_t(x,y)}\,\mathrm{d}z.$$

If we denote by $p^0_t(x,y) = (2\pi t)^{-1/2}\exp(-(z^2+x^2)/2t)2\sinh(zx/t)$ the transition density of Brownian motion killed at 0, then we have the trivial inequality $p^a_t(x,y) \leqslant p^0_t(x,y)$ and furthermore $p^a_t(x,y) \geqslant Cp^0_t(x,y)$ for $x,y \leqslant a/2$ and $t \leqslant a^2$ by Brownian scaling. It follows that

$$W^{x,t,y}_{\text{taboo}}\Big[\int_0^t k(X_s)\,\mathrm{d}s\Big] \leqslant CR^{x,t,y}\Big[\int_0^t k(X_s)\,\mathrm{d}s\Big],$$

where $R^{x,t,y}$ denotes the law of the Bessel bridge of dimension 3. This Bessel bridge is the Doob transform of the Bessel process started at x with respect to the space-time harmonic function $h_y(z,s) = p^0_{t-s}(z,y)/p^0_t(x,y)$. By the standard theory of Doob transforms, this is the Bessel process with additional drift

$$\frac{\mathrm{d}}{\mathrm{d}z}(\log h_y(z,s)) = -\frac{z^2}{t-s} + \frac{\mathrm{d}}{\mathrm{d}z}\log\sinh\frac{zy}{t-s} = -\frac{z^2}{t-s} + \frac{y}{t-s}\coth\frac{zy}{t-s}.$$

This in an increasing function in y and standard comparison theorems for diffusions (see e.g. [53], Theorem IX.3.7) now yield that for $y_1 \leqslant y_2$, we have $R^{x,t,y_2}[k(X_s)] \leqslant R^{x,t,y_1}[k(X_s)]$, since k is a decreasing function. This is true in particular for $y_1 = 0$. Using the self-duality of the Bessel bridge, we can repeat the same reasoning with x. We thus have altogether

$$W^{x,t,y}_{\mathrm{taboo}}\Big[\int_0^t k(X_s)\,\mathrm{d}s\Big] \leqslant C R^{0,t,0}\Big[\int_0^t k(X_s)\,\mathrm{d}s\Big],$$

for any $x,y \leqslant a/2$. This calculation can be done explicitly and yields (2.16). $\square$

3 Preliminaries on branching Markov processes

In this section we recall the definition and basic properties of branching Brownian motion and branching Markov processes in general.

3.1 Definition and notation

A branching Markov process in a Polish space $\mathscr{E}$ can formally be defined in many ways, e.g. as a Markov process in the space $\bigcup_n \mathscr{E}^n$ [35] or as a measure-valued Markov process [25]. A more powerful definition which maintains the genealogy, uses Neveu's *marked trees* [50, 23, 22]. We will briefly describe this construction and refer to these references or to [47, Section 3] for details. The basic idea is to define a *tree* $\mathfrak{t}$ as a suitable subset of $U = \{\varnothing\} \cup \bigcup_{n\geqslant 1} \mathbf{N}^{*n}$, the space of *individuals* (we use the notation $\mathbf{N}^* = \{1,2,3,\dots\}$ and $\mathbf{N} = \{0\} \cup \mathbf{N}^*$). The space U is endowed with the ordering relations $\preceq$ and $\prec$ defined by

$$u \preceq v \iff \exists w \in U : v = uw \quad \text{and} \quad u \prec v \iff u \preceq v \text{ and } u \neq v.$$

To each individual u in a tree $\mathfrak{t}$, we then associate a *trajectory* $\Xi_u \in D(\mathscr{E})$. Here, $D(\mathscr{E})$ is the Skorokhod space of functions $\Xi : [0,\infty) \to \mathscr{E} \cup \{\Delta\}$ ($\Delta \notin \mathscr{E}$ being a cemetary symbol) which are right-continuous with left limits, with $\Xi(0) \neq \Delta$ and for which $\Xi(t) = \Delta$ implies $\Xi(s) = \Delta$ for all $s \geqslant t$. For a trajectory $\Xi \in D(\mathscr{E})$, we define $\zeta(\Xi) = \inf\{t \geqslant 0 : \Xi(t) = \Delta\}$ and set $\zeta_u = \zeta(\Xi_u)$ for an individual $u \in U$. The *birth* and *death* times of the individual u are the defined by $b_u = \sum_{v\prec u} \zeta_v$ and $d_u = b_u + \zeta_u$. The set of individuals *alive at time* t (i.e. for which $b_u \leqslant t < d_u$), is denoted by $N(t)$. The *position* of an individual u at time t is defined by $X_u(t) = \Xi_v(t - b_v)$, where $v \in U$ is such that $v \in \mathscr{N}(t)$ and $v \preceq u$. If $d_u \leqslant t$, then we set $X_u(t) = \Delta$.

Now suppose we are given a defective strong Markov process $\overline{X} = (\overline{X}_t)_{t\geqslant 0}$ on $\mathscr{E}$, with paths in $D(\mathscr{E})$. The law of $\overline{X}$ started in $x \in \mathscr{E}$ will be denoted by $\overline{P}^x$. For simplicity, we will assume that for every $x \in \mathscr{E}$, we have $\zeta(\overline{X}) < \infty$, $\overline{P}^x$-almost surely. Furthermore, let $((q(x,k))_{k\in\mathbf{N}})_{x\in\mathscr{E}}$ be a family of probability measures on $\mathbf{N}$, measurable with respect to x. Then we define the branching Markov process with particle motion $\overline{X}$ and reproduction law q as the (unique) family of probability measures $(\mathbf{P}^x)_{x\in\mathscr{E}}$ on Ω which satisfies

$$\mathbf{P}^x(\mathrm{d}\omega) = \overline{P}^x(\mathrm{d}X_\varnothing) q(X_\varnothing(\zeta_\varnothing -), k_\varnothing) \prod_{i=1}^{k_\varnothing} \mathbf{P}^{X_\varnothing(\zeta_\varnothing -)}(\mathrm{d}\omega^{(i)}). \tag{3.1}$$

Note that by looking at the space-time process $(X_t, t)_{t\geqslant 0}$, we can (and will) extend this definition to the time-inhomogeneous case.

3.2 Stopping lines

The analogue to stopping times for branching Markov processes are *(optional) stopping lines*, for which several definitions exist. For branching Brownian motion, they have first been defined by Chauvin [22], the definition there is however too restricted for our purposes. Jagers [37] has given a definition of more general stopping lines for discrete-time branching processes; our definition will in fact mix both approaches. Note also that Biggins and Kyprianou [9] build up on Jagers' definition of stopping lines and define the subclasses of *simple* and *very simple* stopping lines (again for discrete-time processes). Chauvin's definition then corresponds to the class of very simple stopping lines.

We first define a *(random) line* (called "stopping line" in [37]) to be a set $\ell = \ell(\omega) \subset U \times [0,\infty)$, such that

1. $u \in \mathscr{N}(t)$ for all $(u,t) \in \ell$ and
2. $(u,t) \in \ell$ implies $(v,s) \notin \ell$ for all $v \preceq u$ and $s < t$.

Note that a line is at most a countable set. For a pair $(u,t) \in U \times [0,\infty)$ and a line ℓ, we write $\ell \preceq (u,t)$ if there exists $(v,s) \in \ell$, such that $v \preceq u$ and $s \leqslant t$. For a subset $A \subset U \times [0,\infty)$, we write $\ell \preceq A$ if $\ell \preceq (u,t)$ for all $(u,t) \in A$. If ℓ_1 and ℓ_2 are two lines, we define the line $\ell_1 \wedge \ell_2$ to be the maximal line (with respect to $\preceq$), which is smaller than both lines.

We now define for each $u \in U$ two filtrations on Ω_u by

$$\mathscr{F}_u(t) = (\Omega_u \cap \sigma(\Xi_u(s); 0 \leqslant s \leqslant t - b_u)) \vee \bigvee_{v \prec u} (\Omega_v \cap \sigma(\Xi_v))$$

$$\mathscr{F}_u^{\mathrm{pre}}(t) = (\Omega_u \cap \sigma(\Xi_u(s); 0 \leqslant s \leqslant t - b_u)) \vee \bigvee_{v \not\succeq u} (\Omega_v \cap \sigma(\Xi_v)).$$

Informally, $\mathscr{F}_u(t)$ contains the information on the path from u to the root between the times 0 and t, and $\mathscr{F}_u^{\mathrm{pre}}(t)$ contains this information and of all the other particles *excluding* the descendants of u. In particular, we have $\mathscr{F}_u(t) \subset \mathscr{F}_u^{\mathrm{pre}}(t)$ The filtration $\mathscr{F}_u(t)$ is denoted by $\mathscr{A}_u(t)$ in Chauvin's paper [22] and $\mathscr{F}_u^{\mathrm{pre}}(t)$ corresponds to the *pre-(u,t)-sigma-algebra* as defined by Jagers [37].

We can now define a *stopping line* ("optional line" in [37]) $\mathscr{L}$ to be a random line with the additional property

3a. $\forall (u,t) \in U \times [0,\infty) : \{\omega \in \Omega_u : \mathscr{L} \preceq (u,t)\} \in \mathscr{F}_u^{\mathrm{pre}}(t)$.

The sigma-algebra $\mathscr{F}_{\mathscr{L}}$ of the past of $\mathscr{L}$ is defined to be the set of events $E \in \mathscr{F}$, such that for all $(u,t) \in U \times [0,\infty)$,

$$E \cap \{\omega \in \Omega_u : \mathscr{L} \preceq (u,t)\} \in \mathscr{F}_u^{\mathrm{pre}}(t).$$

For example, for any $t \geqslant 0$, the set $\mathscr{N}(t) \times \{t\}$ is a stopping line. This permits us to define the filtration $(\mathscr{F}_t)_{t \geqslant 0}$ by

$$\mathscr{F}_t = \mathscr{F}_{\mathscr{N}(t) \times \{t\}}.$$

Following Biggins and Kyprianou [9], we now say that $\mathscr{L}$ is a *simple stopping line* or a *very simple stopping line*, if it satisfies the property 3b or 3c below, respectively:

3b. $\forall (u,t) \in U \times [0,\infty) : \{\omega \in \Omega_u : \mathscr{L} \preceq (u,t)\} \in \Omega_u \cap \mathscr{F}_t$.

3c. $\forall (u,t) \in U \times [0,\infty) : \{\omega \in \Omega_u : \mathscr{L} \preceq (u,t)\} \in \mathscr{F}_u(t)$.

Then $\mathscr{N}(t) \times \{t\}$ is obviously a very simple stopping line. Furthermore, if $T : D(\mathscr{E}) \to \mathbf{R}_+$ is a stopping time, then

$$\mathscr{L}_T = \{(u,t) \in U \times [0,\infty) : u \in \mathscr{N}(t) \text{ and } t = T(X_u)\}\}$$

is a very simple stopping line as well. We recall that the definition of stopping lines in [22] is equivalent to the definition of very simple stopping lines given here.

The first important property of stopping lines is the strong branching property. In order to state it, we define the subtree rooted at (u,t) for $t \geqslant 0$ and $u \in \mathscr{N}(t)$ by

$$\omega^{(u,t)} = (\mathfrak{t}^{(u)}, (\Xi'_{uv}; v \in \mathfrak{t}^{(u)})),$$

with $\Xi'_u(\cdot) = \Xi_u(\cdot + t - b_u)$ and $\Xi'_{uv} = \Xi_{uv}$ for $v \in \mathfrak{t}^{(u)} \backslash \{\varnothing\}$. The *strong branching property* ([22, Proposition 2.1], [37, Theorem 4.14]) then states that for every stopping line $\mathscr{L}$, conditioned on $\mathscr{F}_{\mathscr{L}}$, the subtrees $\omega^{(u,t)}$, for $(u,t) \in \mathscr{L}$, are independent with respective distributions $\mathbf{P}^{X_u(t)}$.

3.3 Many-to-few lemmas and spines

Another important tool in the theory of branching processes is the so-called Many-to-one lemma and its recently published extension, the Many-to-few lemma [33] along with the *spine decomposition* technique which comes along with it and has its origins in [45], although it appeared implicitly in the literature before that, see e.g. the references in the same paper. Here we state stopping line versions of these lemmas, which to the knowledge of the author have not yet been stated in this generality in the literature, although they belong to the common folklore. We will therefore only sketch how they can be derived from the existing literature.

We assume for simplicity that the strong Markov process $\overline{X}$ admits a representation as a conservative strong Markov process X with paths in $D(\mathscr{E})$, which is killed at a rate $R(x)$, where $R : \mathscr{E} \to [0,\infty)$ is measurable. The law of X started at x is denoted by P^x and the time of killing by ζ. Given a stopping time T for X, we can then define a stopping time $\overline{T}$ for $\overline{X}$ by setting $\overline{T} = T$, if $T < \zeta$ and $\overline{T} = \infty$ otherwise. For simplicity, we write $\mathscr{L}_T$ for $\mathscr{L}_{\overline{T}}$. Finally, for every $x \in \mathscr{E}$, define $m(x) = \sum_{k\geqslant 0}(k-1)q(x,k)$, $m_1(x) = \sum_{k\geqslant 0} kq(x,k)$ and $m_2(x) = \sum_{k\geqslant 0} k(k-1)q(x,k)$.

We are now going to present the spine decomposition technique, following [31]. They assume that $q(x,0) \equiv 0$, but this restriction is actually not necessary, as noted in [33]. Given a tree $\mathfrak{t}$, a *spine* of $\mathfrak{t}$ is an element of the *boundary* of $\mathfrak{t}$, i.e. it is a line of descent $\xi = (\xi_0 = \varnothing, \xi_1, \xi_2, \dots)$ from the tree, which is finite if and only if the last element is a leaf of the tree. We augment our space Ω to the space Ω^* by

$$\Omega^* = \{(\omega,\xi) : \omega \in \Omega,\ \xi \text{ is a spine of the tree underlying } \omega\}$$

We are going to denote by ξ_t the individual $u \in U$ that satisfies $u \in \mathscr{N}(t)$ and $u \in \xi$ if it exists, and $\xi_t = \varnothing$ otherwise. Instead of the redundant $X_{\xi_t}(t)$, we write $X_\xi(t)$. We also note that the definition of stopping lines can be extended to Ω^* by projection.

Now, for every $x \in \mathscr{E}$, one can define a probability measure $\mathbf{P}^{*,x}$ on Ω^* in the following way:

- Initially, $X_\xi(0) = x$.
- The individuals on the spine move according to the strong Markov process X and die at the rate $m_1(y)R(y)$, when at the point $y \in \mathscr{E}$.
- When an individual on the spine dies at the point $y \in \mathscr{E}$, it leaves k offspring at the point where it has died, with probability $(m_1(x))^{-1}kq(x,\cdot)$ (this is also called the size-biased distribution of $q(x,\cdot)^4$).
- Amongst those offspring, the next individual on the spine is chosen uniformly. This individual repeats the behaviour of its parent (started at the point y).
- The other offspring initiate independent branching Markov processes according to the law $\mathbf{P}^y$, independently of the spine.

This decomposition first appeared in [23].

If we start with n initial particles $1,\dots,n$ at positions $x_1,\dots,x_n$, we can extend this definition by defining a (non-probability) measure $\mathbf{P}^*$, which is the sum of n probability measures $\mathbf{P}^*_1 + \dots + \mathbf{P}^*_n$,

---

[4] The size-biased distribution of the Dirac-mass at 0 is again the Dirac-mass at 0.

where under $\mathbf{P}^*_i$, the particle i follows the law $\mathbf{P}^{*,x_i}$ and the remaining particles $j \neq i$ follow the law $\mathbf{P}^{x_j}$.

We now have the important

Lemma 3.1 (Many-to-one)**.** *Let $\mathscr{L}$ be a simple stopping line. Define T by $(\xi_T, T) \in \mathscr{L}$ if it exists, and $T = \infty$ otherwise. Let Y be a random variable of the form $Y = \sum_{(u,t)\in\mathscr{L}} Y_u \mathbf{1}_{(u\in\xi)}$, where Y_u an $\mathscr{F}_{\mathscr{L}}$-measurable random variable for every $u \in U$. Then*

$$\mathbf{E}\Big[\sum_{(u,t)\in\mathscr{L}} Y_u\Big] = \mathbf{E}^*\Big[Y e^{\int_0^T R(X_\xi(t))m(X_\xi(t))\,\mathrm{d}t}\mathbf{1}_{(T<\infty)}\Big]. \tag{3.2}$$

Proofs of this result can be found for fixed time in [42], [31] or [33]. With simple stopping lines, it has been proven in the discrete setting [9, Lemma 14.1] and their arguments can be used to adapt the proofs in [31] and [33] to yield the result stated here.

Often, we will use a simpler version of the Many-to-one lemma, which is the following

Lemma 3.2 (Simple Many-to-one)**.** *Let $T = T(X)$ be a stopping time for the strong Markov process X which satisfies $P^x(T < \infty) = 1$ for every $x \in \mathscr{E}$. Let $f : \mathscr{E} \to [0,\infty)$ be measurable. Then we have*

$$\mathbf{E}^x\Big[\sum_{(u,t)\in\mathscr{L}_T} f(X_u(t))\Big] = E^x\Big[e^{\int_0^T R(X_t)m(X_t)\,\mathrm{d}t} f(X_T)\Big]$$

The next lemma tells us about second moments of sums of the previous type. To state it, we define for a stopping time T for X the density of the branching Markov process before $\mathscr{L}_T$, by

$$p_T(x,\mathrm{d}y,t) = \mathbf{E}^x\Big[\sum_{u\in\mathscr{N}(t)} \mathbf{1}_{(X_u(t)\in\mathrm{d}y,\ t<T(X_u))}\Big]. \tag{3.3}$$

Lemma 3.3. *Let H be the hitting time functional of a closed set $F \subset \mathscr{E}$ on $D(\mathscr{E})$ which satisfies $P^x(H < \infty) = 1$ for every $x \in \mathscr{E}$. Let $f : \mathscr{E} \to [0,\infty)$ be measurable. Then we have*

$$\begin{aligned}\mathbf{E}^x\Big[\Big(\sum_{(u,t)\in\mathscr{L}_H} f(X_u(t))\Big)^2\Big] &= \mathbf{E}^x\Big[\sum_{(u,t)\in\mathscr{L}_H} (f(X_u(t)))^2\Big] \\ &\quad + \int_0^\infty \int_{\mathscr{E}} p_H(x,\mathrm{d}y,s)R(y)m_2(y)\Big(\mathbf{E}^y\Big[\sum_{(u,t)\in\mathscr{L}_H} f(X_u(t))\Big]\Big)^2\,\mathrm{d}s \end{aligned}\tag{3.4}$$

Remark 3.4. The intuitive explanation of Lemma 3.3 (see, e.g. the proof of Proposition 18 in [7]) is that a particle at the point y spawns an expected number of $R(y)m_2(y)\,\mathrm{d}s$ of ordered pairs of particles during the time interval $[s, s+\mathrm{d}s]$, and by the strong branching property, the two particles in a pair evolve independently. This lemma can be proven using the Many-to-few lemma from [33] (which is valid for stopping lines as well by the same argument as the one above) or with Lemma 3.1, by noting that

$$\Big(\sum_{(u,t)\in\mathscr{L}_H} f(X_u(t))\Big)^2 = \sum_{(u,t)\in\mathscr{L}_H} (f(X_u(t)))^2 + \sum_{(u,t)\in\mathscr{L}_H} \Big(f(X_u(t)) \sum_{(v,s)\in\mathscr{L}_H,\ v\neq u} f(X_v(s))\Big).$$

Taking for X the space-time process $(Y_t, t)_{t\geqslant 0}$ of a possibly non-homogeneous strong Markov process $(Y_t)_{t\geqslant 0}$ with paths in $D(\mathscr{E})$ and the closed set $F = \mathscr{E} \times \{t\}$, for some $t \geqslant 0$, we obtain the following useful corollary, which appeared already in [57] and [54] in the homogeneous case.

Lemma 3.5. *Let $f : \mathscr{E} \times \mathbf{R}_+ \to [0,\infty)$ be measurable and let $t \geqslant 0$. Then we have*

$$\begin{aligned}\mathbf{E}^{(x,0)}\Big[\Big(\sum_{u\in\mathscr{N}(t)} f(Y_u(t),t)\Big)^2\Big] &= \mathbf{E}^{(x,0)}\Big[\sum_{u\in\mathscr{N}(t)} (f(Y_u(t),t))^2\Big] \\ &\quad + \int_0^t \int_{\mathscr{E}} p(x,\mathrm{d}y,s)R(y,s)m_2(y,s)\Big(\mathbf{E}^{(y,s)}\Big[\sum_{u\in\mathscr{N}(t)} f(Y_u(t),t)\Big]\Big)^2\,\mathrm{d}s \end{aligned}\tag{3.5}$$

4 BBM with absorption at a critical line

From this section on, $q(k)$ will denote a law on $\{0,1,2,\dots\}$ and L a random variable with law $q(k)$. We define $m = \mathbf{E}[L-1]$ and $m_2 = \mathbf{E}[L(L-1)]$ and suppose that $m > 0$ and $m_2 < \infty$. We study the branching Markov process where, starting with a single particle at the origin, particles move according to standard Brownian motion with drift -1 and branch at rate $\beta_0 = 1/(2m)$ into k particles according to the reproduction law $q(k)$. At the point $-y$, we add an *absorbing barrier* to the process, i.e. particles hitting this barrier are instantly killed. Formally, we are considering the process up to the stopping line $\mathscr{L}_{H_{-y}}$, where H_{-y} is the hitting time functional of the point $-y$. It is well-known since Kesten [39] that this process gets extinct almost surely. As a consequence, the number of particles absorbed at the barrier, i.e. the random variable

$$N_y = \#\mathscr{L}_{H_{-y}},$$

is almost surely finite. By the strong branching property and the translational invariance of Brownian motion, one sees that the process $(N_y)_{y\geqslant 0}$ is a continuous-time Galton–Watson process, a fact which was first noticed by Neveu [51] (see [4], Chapter III or [34], Chapter V for an introduction to continuous-time Galton–Watson processes). Let $u(s)$ be its infinitesimal generating function. Neveu stated that $u = \psi' \circ \psi^{-1}$, where ψ is a so-called travelling wave of the FKPP (Fisher–Kolmogorov–Petrovskii–Piskounov) equation: Write $f(s) = \sum_k s^k q(k)$. Then ψ is a solution of the equation

$$\frac{1}{2}\psi'' - \psi' = \beta_0(\psi - f\circ\psi), \tag{4.1}$$

with $\psi(-\infty) = 1$ and $\psi(+\infty)$ is the extinction probability of the process, i.e. the smaller root of $f(s) = s$. For a proof of these results, see [46, Section 3].

In the same paper [51], Neveu introduced his *multiplicative martingales*, which he used to derive the Seneta-Heyde norming for the martingale $e^{-y}N_y$. He proved that in the case of binary branching, one has

$$W_y := ye^{-y}N_y \to W \quad \text{almost surely as } y\to\infty, \tag{4.2}$$

where $W > 0$ almost surely. His proof relied on a known asymptotic for the travelling wave ψ, namely that

$$1-\psi(-x) \sim Kxe^{-x}, \qquad \text{as } x\to\infty, \tag{4.3}$$

for some constant $K > 0$. It was recently shown [59] that this asymptotic is true if and only if $E[L\log^2 L] < \infty$ and the proof of (4.2) works in this case as well. We also still have in this case, for every $x \in \mathbf{R}$,

$$E[e^{-e^x W}] = \psi(x), \tag{4.4}$$

a fact which was already proven by Neveu [51] for dyadic branching.

In [7], further properties of the limit W have been established under the hypothesis of dyadic branching, namely

$$\mathbf{P}(W > x) \sim \frac{1}{x}, \qquad \text{as } x\to\infty, \tag{4.5}$$

and

$$\mathbf{E}[W\mathbf{1}_{(W\leqslant x)}] - \log x \to c_{4.6}, \qquad \text{as } x\to\infty, \tag{4.6}$$

for some constant $c_{4.6} \in \mathbf{R}$. Equation (4.5) has been proven in Propositions 27 and 40 of [7], and (4.6) appears in the proof of Proposition 39 of the same paper. Their arguments were very ingenious but indirect and although they could be extended to general reproduction laws with finite variance, we will reprove them here directly under (probably) minimal assumptions, based on methods of [46]. The main result in this section is

Proposition 4.1. *If* $E[L\log^2 L] < \infty$*, then* (4.5) *holds. If* $E[L\log^3 L] < \infty$*, then* (4.6) *holds.*

See also [20] for a proof of (4.5) in the case of branching random walk. Before proving this result in the next subsection, we state a lemma which is immediate from (4.2) and the fact that N_y is almost surely finite (see also Corollary 25 in [7]):

Lemma 4.2. *Suppose $E[L\log^2 L] < \infty$. For any $\eta > 0$, there exist y and ζ, such that $y \geqslant \eta^{-1}$ and*

$$\mathbf{P}(|W_y - W| > \eta) + \mathbf{P}(N_y > \zeta) + \mathbf{P}(\mathscr{L}_{H_{-y}} \nsubseteq U \times [1,\zeta]) + \mathbf{P}\Big(\sup_{0\leqslant t\leqslant\zeta} \#\{u \in \mathscr{N}(t) : (u,t) \preceq \mathscr{L}_{H_{-y}}\} > \zeta\Big) \leqslant \eta.$$

4.1 Proof of Proposition 4.1

Define $\chi(\lambda) = E[e^{-\lambda W}]$ for $\lambda \geqslant 0$. Our first result is:

Lemma 4.3. *Suppose $\mathbf{E}[L\log^2 L] < \infty$. Then $\chi''(\lambda) \sim \lambda^{-1}$ as $\lambda \to 0+$. Furthermore, $\mathbf{E}[L\log^3 L] < \infty$ if and only if $r(\lambda) = \lambda^{-1} - \chi''(\lambda) \geqslant 0$ for $\lambda \geqslant 0$, with $\int_0^1 r(\lambda)\,\mathrm{d}\lambda < \infty$.*

Proof. Define $\phi(x) = 1 - \psi(-x)$, such that $u(s) = \phi'(\phi^{-1}(s))$. By (4.3) and the hypothesis $\mathbf{E}[L\log^2 L] < \infty$,

$$\phi(x) \sim Kxe^{-x}, \qquad \text{as } x \to \infty. \tag{4.7}$$

Furthermore, by (4.1), we have

$$\frac{1}{2}\phi''(x) + \phi'(x) = \beta_0(f(1-\phi(x)) - (1-\phi(x))). \tag{4.8}$$

Setting $g(s) = 2\beta_0[f(1-s) - 1 + f'(1)s] \geqslant 0$ and $\rho = \phi + \phi'$, we get from (4.8), and the fact that $\beta_0 = 1/(2m)$ and $f'(1) = m+1$ by definition,

$$\rho'(x) = -\rho(x) + g(\phi(x)). \tag{4.9}$$

As in the proof of Theorem 1.1 of [46], we will study the function ρ through the integral equation corresponding to (4.9), namely

$$\rho(x) = e^{-x}\Big(\rho(0) + \int_0^x e^y g(\phi(y))\,\mathrm{d}y\Big) = e^{-x}\Big(\rho(0) + \int_{\phi(x)}^{\phi(0)} \frac{e^{\phi^{-1}(s)}g(s)}{-u(s)}\,\mathrm{d}s\Big). \tag{4.10}$$

Now, by Theorem B of [11] (see also Theorem 8.1.8 in [12]) we have for every $d \geqslant 0$,

$$\int_0^1 \frac{\log^d \frac{1}{s}}{s^2} g(s)\,\mathrm{d}s < \infty \quad\iff\quad \int_0^1 \frac{\log^d \frac{1}{s}}{s} g'(s)\,\mathrm{d}s < \infty \quad\iff\quad E[L\log^{1+d} L] < \infty. \tag{4.11}$$

Furthermore, by Proposition 3.2 of [46], we have $-u(s) \sim s$ as $s \to 0$, and by (4.7), we have $e^{\phi^{-1}(s)} \sim (\log 1/s)/s$ as $s \to 0$. By the hypothesis $E[L\log^2 L] < \infty$, this gives

$$\int_0^{\phi(0)} \frac{e^{\phi^{-1}(s)}g(s)}{-u(s)}\,\mathrm{d}s < \infty,$$

whence, by (4.10),

$$\rho(x) \sim Ke^{-x}, \qquad \text{as } x \to \infty, \tag{4.12}$$

where the constant K is actually the same as the one in (4.7), see again the proof of Theorem 1.1 of [46]. Now, from (4.4), we get $\chi(\lambda) = 1 - \phi(-\log\lambda)$, whence, by (4.9) and (4.10),

$$\begin{aligned} \chi''(\lambda) = -\frac{1}{\lambda^2}\rho'(-\log\lambda) &= \frac{K}{\lambda} + \frac{1}{\lambda^2}\Big(-\lambda\int_{-\log\lambda}^{\infty} e^y g(\phi(y))\,\mathrm{d}y - g(\phi(-\log\lambda))\Big) \\ &= \frac{K}{\lambda} + \frac{1}{\lambda}\Big(\int_{-\log\lambda}^{\infty} e^y \phi'(y) g'(\phi(y))\,\mathrm{d}y\Big), \end{aligned} \tag{4.13}$$

where the last equation follows from integration by parts. This proves the first statement, with the constant K instead of 1, since the last integral vanishes as $\lambda \to 0$. Now, setting

$$r(\lambda) = -\frac{1}{\lambda}\Big(\int_{-\log\lambda}^{\infty} e^y \phi'(y) g'(\phi(y))\,\mathrm{d}y\Big),$$

we first remark that $r(\lambda) \geqslant 0$, since the integrand is negative for $y \in \mathbf{R}$. By the Fubini–Tonelli theorem, we then have

$$\begin{aligned}\int_0^1 r(\lambda)\mathrm{d}\lambda &= \int_0^1 -\frac{1}{\lambda}\int_{-\log\lambda}^{\infty} e^y\phi'(y)g'(\phi(y))\,\mathrm{d}y\,\mathrm{d}\lambda = \int_0^{\infty} -ye^y\phi'(y)g'(\phi(y))\,\mathrm{d}y\\ &= \int_0^{\phi(0)} e^{\phi^{-1}(y)}\phi^{-1}(y)g'(y)\,\mathrm{d}y,\end{aligned}$$

which is finite if and only if $E[L\log^3 L] < \infty$, by (4.11) and the fact that $e^{\phi^{-1}(y)}\phi^{-1}(y) \sim (\log^2 1/s)/s$. This proves the second statement, again with the constant K instead of 1.

The previous arguments worked for every travelling wave ψ. In order to show that that the constant K is equal to 1 in our case, we use Neveu's multiplicative martingale (this idea was also used in [44], Theorem 2.5). It was observed by Neveu [51] (see also [22] for a rigorous proof), that $((1-\phi(x+y))^{N_y})_{y\geqslant 0}$ is a martingale for every $x \in \mathbf{R}$ with values in $[0,1]$. By (4.2) and (4.7), we then get by dominated convergence, for every $x \in \mathbf{R}$,

$$\chi(Ke^x) = \lim_{y\to\infty} E[e^{-Kye^{x-y}N_y}] = \lim_{y\to\infty} E[(1-\phi(y-x))^{N_y}] = 1-\phi(-x) = \chi(e^x).$$

This yields $K = 1$. □

Remark 4.4. Choosing arbitrary initial points $x_0, x_1 \in \mathbf{R}$ instead of 0 in (4.10), one sees that

$$e^{x_0}\rho(x_0) + \int_{x_0}^{\infty} e^y g(\phi(y))\,\mathrm{d}y = e^{x_1}\rho(x_1) + \int_{x_1}^{\infty} e^y g(\phi(y))\,\mathrm{d}y.$$

In particular, since ρ is bounded, letting $x_0 \to -\infty$ and $x_1 \to +\infty$ yields $\int_{-\infty}^{\infty} e^y g(\phi(y))\,\mathrm{d}y = 1$. One could hope (see the proof of Proposition 4.1 below) that this helps in determining the constant $c_{4.6}$, but apparently this does not seem to be the case.

Proof of Proposition 4.1. For $n \in \mathbf{N}$, we define the function

$$V_n(x) = \int_0^x y^n P(W \in \mathrm{d}y) = E[W^n \mathbf{1}_{(W\leqslant x)}],$$

such that with $\chi^{(n)}$ denoting the n-th derivative of χ, we have for $\lambda > 0$,

$$\chi^{(n)}(\lambda) = (-1)^n \int_0^{\infty} e^{-\lambda x}\mathrm{d}V_n(x).$$

If $E[L\log^2 L] < \infty$, Proposition 4.1 and Karamata's Tauberian theorem ([29], Theorem XIII.5.2 or [12], Theorem 1.7.1) now yields

$$V_2(x) \sim x, \qquad \text{as } x \to \infty. \tag{4.14}$$

By an integration by parts argument (see also [29], Theorem VIII.9.2 or [12], Theorem 8.1.2), we get (4.5). Now suppose that $E[L\log^3 L] < \infty$. By Lemma 4.3, we have $\chi'(\lambda) - \log\lambda \to c \in \mathbf{R}$, as $\lambda \to 0$. By Theorem 3.9.1 from [12] (with $\ell(x) \equiv 1$), this yields

$$V_1(x) - \log x \to \gamma - c, \qquad \text{as } x \to \infty,$$

where γ is the Euler–Mascheroni constant. This is exactly (4.6).

□

5 BBM in an interval

In this section we study branching Brownian motion killed upon exiting an interval. Many ideas in this section (except for Section 5.5 and parts of Section 5.3) stem from Sections 2 and 3 of [7] and for completeness, we will reprove some of their results with streamlined proofs. However, we will also extend their results to the case of Brownian motion with variable drift.

5.1 Notation

During the rest of the paper, the symbol C stands for a *positive* constant, which may only depend on the reproduction law q. Its value may change from line to line. If a subscript is present, then this subscript is the number of the equation where this constant appears for the first time (example: $C_{5.36}$). In this case, this constant is fixed after its value has been chosen in the corresponding equation. If X is any mathematical expression, then the symbol $O(X)$ stands for a possibly random term whose absolute value is bounded by $C|X|$.

Recall the definition of $q(k)$, m, m_2 and β_0 from Section 4 and the hypotheses on m and m_2. In this section, we let $a \geqslant \pi$ and set

$$\mu = \sqrt{1 - \frac{\pi^2}{a^2}}. \tag{5.1}$$

From (5.1), one easily gets the basic estimate

$$0 \leqslant 1 - \mu \leqslant \frac{\pi^2}{2\mu a^2}. \tag{5.2}$$

We then denote by $\mathbf{P}^x$ the law of the branching Markov process where, starting with a single particle at the point $x \in \mathbf{R}$, particles move according to Brownian motion with variance 1 and drift $-\mu$ and branch at rate 1 into k particles according to the reproduction law $q(k)$. Expectation with respect to $\mathbf{P}^x$ is denoted by $\mathbf{E}^x$. On the space of continuous functions from $\mathbf{R}_+$ to $\mathbf{R}$, we define H_0 and H_a to be the hitting time functionals of 0 and a. We further set $H = H_0 \wedge H_a$. Then note that the density of the branching Brownian motion before $\mathscr{L}_H$, as defined in (3.3), has a density with respect to Lebesgue measure given for $t > 0$ and $x, y \in (0, a)$ by

$$\mathbf{p}_t(x, y) = e^{\mu(x-y) + \frac{\pi^2}{2a^2}t} p_t^a(x, y), \tag{5.3}$$

where p_t^a was defined in (2.3).

Now, let $f : \mathbf{R}_+ \to \mathbf{R}_+$ be non-decreasing, with $f(0) = 0$, continuous and such that the left-derivative f' exists everywhere and is of bounded variation. Such a function will be called a *barrier function.* We define

$$\|f\| = \max\Big\{\|f\|_\infty, \|f'\|_\infty, \|f'\|_\infty^2, \int_0^\infty |\mathrm{d}f'(s)|\Big\}, \tag{5.4}$$

where $\|\cdot\|_\infty$ is the usual supremum norm. Furthermore, we set $\mathrm{Err}(f, t) = \|f\|\Big(\frac{1}{a} + \frac{t}{a^3}\Big)$. Now define

$$\mu_t = \mu + \frac{\mathrm{d}}{\mathrm{d}t} f(t/a^2) = \mu + \frac{1}{a^2} f'(t/a^2), \tag{5.5}$$

such that $\mu_0 = \mu$ and $\mu_t \geqslant \mu$ for all $t \geqslant 0$. We denote by $\mathbf{P}_f^x$ the law of the branching Brownian motion described above, but with infinitesimal drift $-\mu_t$. Expectation with respect to $\mathbf{P}_f^x$ is denoted by $\mathbf{E}_f^x$ and the density of the process is denoted by $\mathbf{p}_t^f(x, y)$.

The above definitions can be extended to arbitrary initial configurations of particles distributed according to a counting measure ν on $(0, a)$. In this case the superscript x is replaced by ν or simply omitted if ν is known from the context.

5.2 The processes Z_t and Y_t

Recall from Section 3 that the set of particles alive at time t is denoted by $\mathscr{N}(t)$. We define

$$\mathscr{N}_{0,a}(t) = \{u \in \mathscr{N}(t) : H(X_u) > t\},$$

where H was defined in the previous subsection. Now set $w_Z(x) = ae^{\mu(x-a)}\sin(\pi x/a)\mathbf{1}_{(x\in[0,a])}$ and $w_Y(x) = e^{\mu(x-a)}\mathbf{1}_{(x\geqslant 0)}$ and define

$$Z_t = \sum_{u\in\mathscr{N}_{0,a}(t)} w_Z(X_u(t)) \quad \text{and} \quad Y_t = \sum_{u\in\mathscr{N}_{0,a}(t)} w_Y(X_u(t)).$$

Then Z_t is a martingale under $\mathbf{P}^x$, the proof of which is standard and relies on the branching property, the Many-to-one lemma (Lemma 3.2) and the fact that $e^{t/2}w_Z(B_t)$ is a martingale for a Brownian motion with drift $-\mu$ killed at 0 and a, which is easily seen by Itō's formula, for example. Furthermore, it is easy to see as well that Z_t is a supermartingale under $\mathbf{P}^x_f$.

The following lemma relates the density of BBM with variable drift to BBM with fixed drift.

Lemma 5.1. *For all $x, y \in [0,a]$ and $t \geqslant 0$, we have $\mathbf{p}^f_t(x,y) = \mathbf{p}_t(x,y)e^{-f(t/a^2)+O(\mathrm{Err}(f,t))}$.*

Proof. By the Many-to-one lemma and Girsanov's theorem, we have

$$\begin{aligned}\mathbf{p}^f_t(x,y) &= e^{\beta_0 m t} W^x_{-\mu_t}\left(B_t \in \mathrm{d}y,\, H > t\right)\\ &= \exp\left(t/2 - \int_0^t \frac{\mu_s^2-\mu^2}{2}\mathrm{d}s\right) W^x_{-\mu}\left(\exp\Big(-\int_0^t \mu_s - \mu\,\mathrm{d}B_s\Big),\, B_t\in\mathrm{d}y, H > t\right).\end{aligned} \tag{5.6}$$

By the integration by parts formula $\int_0^t \mu_s\,\mathrm{d}B_s = \mu_t B_t - \mu B_0 - \int_0^t B_s\,\mathrm{d}\mu_s$ and some easy estimates, we get with (5.3),

$$\mathbf{p}^f_t(x,y) = \mathbf{p}_t(x,y)e^{-\mu f(t/a^2)+O(\mathrm{Err}(f,t))}.$$

The lemma now follows from (5.2). □

Proposition 5.2. *Under any initial configuration of particles, for every $t \geqslant 0$, we have*

$$\mathbf{E}_f[Z_t] = Z_0 e^{-f(t/a^2)+O(\mathrm{Err}(f,t))}, \tag{5.7}$$

and if in addition $\mu \geqslant 1/2$, then

$$\mathrm{Var}_f(Z_t) \leqslant Ce^{-f(t/a^2)+O(\mathrm{Err}(f,t))}\Big(\frac{t}{a^3}Z_0 + Y_0\Big). \tag{5.8}$$

Furthermore, we have for every $t \geqslant 0$ (without hypothesis on μ),

$$\mathbf{E}_f[Y_t] \leqslant Ce^{-f(t/a^2)+O(\mathrm{Err}(f,t))}Y_0. \tag{5.9}$$

and for $t \geqslant a^2$,

$$\mathbf{E}_f[Y_t] \leqslant Ce^{-f(t/a^2)+O(\mathrm{Err}(f,t))}\frac{Z_0}{a}. \tag{5.10}$$

Moreover, for every $a^2 \leqslant t \leqslant a^3$, we have

$$\mathrm{Var}_f(Y_t) \leqslant Ce^{-f(t/a^2)+O(\mathrm{Err}(f,t))}\frac{Y_0}{a}. \tag{5.11}$$

Proof. Equation (5.7) follows from Lemma 5.1 and the fact that Z_t is a martingale under $\mathbf{P}^x$. In order to show (5.9) and (5.10), it suffices by Lemma 5.1 to consider the case without variable drift. We first suppose that $t \geqslant a^2$. By (5.3) and (2.7), we get

$$\mathbf{E}^x[Y_t] \leqslant e^{\mu(x-a)} \int_0^a e^{\frac{\pi^2}{2a^2}t} p_t(x,y)\,\mathrm{d}y \leqslant Ce^{\mu(x-a)} \sin(\pi x/a) \int_0^a \frac{2}{a} \sin(\pi y/a)\,\mathrm{d}y.$$

The last integral is independent of a. Summing over x yields (5.10) as well as (5.9) in the case $t \geqslant a^2$. Now, if $t < a^2$, by the Many-to-one lemma and Girsanov's theorem, we have

$$\mathbf{E}^x[Y_t] = e^{\beta_0 m t} W^x_{-\mu}\Big[e^{\mu(X_t-a)},\ H_0 \wedge H_a < t\Big] = e^{\pi^2 t/(2a^2)} W^x[H_0 \wedge H_a < t] e^{\mu(x-a)}.$$

Summing over x yields (5.9).

In order to prove (5.8), we have by Lemma 3.5,

$$\mathbf{E}^x_f[Z_t^2] = \mathbf{E}^x_f\Big[\sum_{u\in\mathscr{N}_{0,a}(t)} w_Z(X_u(t))^2\Big] + \beta_0 m_2 \int_0^a \int_0^t \mathbf{p}^f_s(x,y)(\mathbf{E}^{(y,s)}_f[Z_t])^2\,\mathrm{d}s\,\mathrm{d}y. \tag{5.12}$$

By Lemma 5.1 and the fact that Z_t is a martingale with respect to the law $\mathbf{P}^x$, this yields

$$\mathbf{E}^x_f[Z_t^2] \leqslant Ce^{-f(t/a^2)+O(\mathrm{Err}(f,t))} \left(\mathbf{E}^x\Big[\sum_{u\in\mathscr{N}_{0,a}(t)} w_Z(X_u(t))^2\Big] + \int_0^a \int_0^t \mathbf{p}_s(x,y) w_Z(y)^2\,\mathrm{d}s\,\mathrm{d}y\right). \tag{5.13}$$

Now we have $w_Z(x)^2 = (a\sin(\pi x/a)e^{-\mu(a-x)})^2 \leqslant \pi^2(a-x)^2 e^{-2\mu(a-x)} \leqslant Cw_Y(x)$ for $x \in (0,a)$, because $\mu \geqslant 1/2$ by hypothesis. This yields

$$S_1 := \mathbf{E}^x\Big[\sum_{u\in\mathscr{N}_{0,a}(t)} w_Z(X_u(t))^2\Big] \leqslant C\mathbf{E}^x[Y_t] \leqslant Cw_Y(x), \tag{5.14}$$

by (5.9). Now, by (5.3) and (2.11), we have

$$S_2 := \int_0^a \int_0^t \mathbf{p}_s(x,y) w_Z(y)^2\,\mathrm{d}s\,\mathrm{d}y = ae^{\mu(x-a)} \int_0^a ae^{\mu(y-a)} \sin^2(\pi y/a) J^a(x,y,t)\,\mathrm{d}y.$$

Lemma 2.1 now gives

$$\begin{aligned} S_2 &\leqslant Cae^{\mu(x-a)} \int_0^a e^{-\mu y} \sin^2(\pi y/a)\Big(t\sin(\pi x/a)\sin(\pi y/a) + ay\Big)\,\mathrm{d}y \\ &\leqslant Cae^{\mu(x-a)}\Big(\sin(\pi x/a)\frac{t}{a^3} + \frac{1}{a}\Big) \int_0^\infty e^{-\mu y} y^3\,\mathrm{d}y, \end{aligned} \tag{5.15}$$

the last line following again from the change of variables $y \mapsto a-y$ and the inequality $\sin x \leqslant x$. Using again the fact that $\mu \geqslant 1/2$, equations (5.13), (5.14) and (5.15) now imply

$$\mathbf{E}^x_f[Z_t^2] \leqslant Ce^{-f(t/a^2)+O(\mathrm{Err}(f,t))}\Big(\frac{t}{a^3} w_Z(x) + w_Y(x)\Big). \tag{5.16}$$

If we write the positions of the initial particles as $x_1,\dots,x_n$, then by the independence of their contributions to Z_t,

$$\mathrm{Var}_f(Z_t) = \sum_i \mathrm{Var}^{x_i}_f(Z_t) \leqslant \sum_i \mathbf{E}^{x_i}_f[Z_t^2]. \tag{5.17}$$

Equations (5.16) and (5.17) now prove (5.8). Equation (5.11) is proven similarly. □

5.3 The number of particles

In this subsection, we establish precise first and second moment estimates for the number of particles alive at a time t. These estimates extend those of [7], which are effective only when $t \gg a^2$. For $r \in [0,a]$ and $t \geqslant 0$, we denote by $N_t(r)$ the number of particles in $[r,a]$ at time t.

Proposition 5.3. *Suppose $\mu \geqslant 1/2$. Let $t \geqslant 0$, $x, r \in [0,a)$ and suppose that $x \geqslant (r + a/20)\mathbf{1}_{(t\leqslant a^2)}$. Then*

$$\mathbf{E}^x[N_t(r)] = e^{\mu(x-r)}\left[-\frac{2(1+\mu r)e^{\pi^2 t/(2a^2)}}{a^2}\theta'\Big(\frac{x}{a},\frac{t}{a^2}\Big) + O\left((1+r^3)\frac{\sin(\pi x/a)\vee \mathbf{1}_{(t\leqslant a^2)}}{a^4}\right)\right]. \quad (5.18)$$

In particular, if $a/20 + r\mathbf{1}_{(t\leqslant a^2)} \leqslant x < a$, then

$$\mathbf{E}^x[N_t(r)] = \frac{2\pi(1+\mu r)e^{\mu(a-r)}}{a^3}\Big[w_Z(x)\bar\theta\left(\frac{t}{a^2}\right) + O\left(\frac{(a-x)^2+(1+r^2)}{a}\left(\frac{w_Z(x)}{a}\vee w_Y(x)\mathbf{1}_{(t\leqslant a^2)}\right)\right)\Big]. \quad (5.19)$$

Moreover, for every $x \in (0,a)$ and $t > 0$, we have

$$\mathbf{E}^x[N_t(r)] = \frac{2\pi(1+\mu r)e^{\mu(a-r)}}{a^3}w_Z(x)\Big(1+\mathrm{error}\Big), \quad (5.20)$$

with $|\mathrm{error}| \leqslant E_{t/a^2}(1+O(1/a^2))$.

Proof. Let $t \geqslant 0$ and $0 \leqslant x < 1$. By (5.3) and Brownian scaling, we have

$$\mathbf{E}^{ax}[N_{ta^2}(r)] = e^{\mu a x + \pi^2 t/2}\int_{r/a}^1 e^{-\mu a z}p_t^1(x,z)\,\mathrm{d}z. \quad (5.21)$$

Note that $(\partial/\partial z)p_t^1(x,z)|_{z=0} = -2\theta'(x,t)$, by (2.4). Taylor's formula then implies that there exists $\xi \in [0,z]$, such that

$$p_t^1(x,z) = -2z\theta'(x,t) + z^2\frac{\partial^2}{\partial z^2}p_t^1(x,z)\big|_{z=\xi}. \quad (5.22)$$

We first note that for all $x \in (0,1)$ and $t \geqslant 1$, or for all $x \in [\frac{1}{40},1]$ and $t \leqslant 1$, one has by (2.1) and (2.2),

$$|\theta'(x,t)| \leqslant C(\sin(\pi x)\vee \mathbf{1}_{(t\leqslant 1)}). \quad (5.23)$$

Furthermore, if $t \geqslant 1$, we have

$$\begin{aligned}\Big|\frac{\partial^2}{\partial z^2}p_t^1(x,z)\big|_{z=\xi}\Big| &= \pi^2\Big|\sum_{n=1}^\infty e^{-\pi^2 n^2 t/2}n^2\sin(\pi n x)\sin(\pi n \xi)\Big| \\ &\leqslant \pi^4\xi\sin(\pi x)\sum_{n=1}^\infty n^4 e^{-\pi^2n^2t/2} \leqslant Cz\sin(\pi x)e^{-\pi^2 t/2},\end{aligned} \quad (5.24)$$

by the inequality $|\sin nx| \leqslant n\sin x$, $x \in [0,\pi]$. Equations (5.22), (5.23) and (5.24) now give

$$\int_{r/a}^1 e^{-\mu a z}p_t^1(x,z)\,\mathrm{d}z = -\frac{2\theta'(x,t)}{\mu^2a^2}(1+\mu r)e^{-\mu r} + O\Big((1+r^3)e^{-\mu r}\frac{\sin(\pi x)}{a^4}e^{-\pi^2t/2}\Big). \quad (5.25)$$

Now suppose that $t \leqslant 1$, $x, r \in [0,a)$ and $x \geqslant r + a/20$. By (2.4) and the mean value theorem, there exists $\xi' \in [0,z]$, such that $(\partial^2/\partial z^2)p_t^1(x,z) = 2z\theta'''(x+\xi',t)$. With (2.2), one then easily sees that for $z \leqslant x - a/40$, one has

$$\Big|\frac{\partial^2}{\partial z^2}p_t^1(x,z)\Big| \leqslant Cze^{-C_{5.26}/t}. \quad (5.26)$$

Equations (5.22), (5.23) and (5.26) now give, for $x' = x - a/40$,

$$\int_{r/a}^{x'/a} e^{-\mu a z} p_t^1(x,z)\,\mathrm{d}z = -\frac{2\theta'(x,t)}{\mu^2 a^2}(1+\mu r)e^{-\mu r} + O\Big(\frac{(1+r^3)e^{-\mu r}}{a^4}\Big). \tag{5.27}$$

Furthermore, we have

$$\int_{x'/a}^{1} e^{-\mu a z} p_t^1(x,z)\,\mathrm{d}z \leqslant e^{-\mu x'} \int_{x'/a}^{1} p_t^1(x,z)\,\mathrm{d}z \leqslant e^{-\mu x'}. \tag{5.28}$$

Equations (5.21), (5.25), (5.27), (5.28) and the hypothesis on μ now imply (5.18).

In order to prove (5.19), let $x \in [1/20, 1]$. By Taylor's formula and the fact that $\theta(x,t)$ is a solution to the heat equation with diffusion coefficient $1/2$, there exists $\xi \in [x,1]$, such that

$$-\theta'(x,t) = 2(1-x)\frac{\partial}{\partial t}\theta(1,t) - (1-x)^2\theta'''(\xi,t). \tag{5.29}$$

With (2.1) and (2.2), one now easily sees that $|\theta'''(\xi,t)| \leqslant C(e^{-\pi^2 t/2}\sin(\pi x) \vee \mathbf{1}_{(t\leqslant 1)})$. Equations (5.2), (5.18) and (5.29) now readily imply (5.19). For the last equation, by (5.3) and (2.7), we have for every $x \in (0,a)$ and $t > 0$,

$$\mathbf{E}^x[N(ta^2)] = (1+\mathrm{err})w_Z(x)\frac{2e^{\mu a}}{a^2}\int_0^a e^{-\mu y}\sin(\pi y/a)\,\mathrm{d}y, \tag{5.30}$$

with $|\mathrm{err}| \leqslant E_t$. Evaluating this integral and using (5.2) and the hypothesis on μ yields (5.20). $\square$

We will often use the following handier upper bounds on $\mathbf{E}^x[N_t(r)]$:

Lemma 5.4. *Suppose $\mu \geqslant 1/2$. Let $t \geqslant 0$, $x, r \in [0,a]$ and suppose that $x \geqslant (r + a/20)\mathbf{1}_{(t\leqslant a^2)}$.*

$$\mathbf{E}^x[N_t(r)] \leqslant C(1+r)\frac{e^{\mu(a-r)}}{a^3}\Big(w_Z(x) + \frac{1+r^2}{a}w_Y(x)\mathbf{1}_{(t\leqslant a^2)}\Big). \tag{5.31}$$

Furthermore, we have for all $x, r \in [0,a]$ and $t \geqslant 0$, $\mathbf{E}^x[N_t(r)] \leqslant Ce^{\mu(x-r)}$.

Proof. One sees from (2.1) and (2.2) that $\theta''(x,t) \leqslant C$ for $(x,t) \in [0,1]\times[1,\infty) \cup [1/20,1]\times[0,1]$. Equation (5.31) now follows from (5.18) and the mean value theorem. The second statement on the other hand follows from the inequality $N_t(r) \leqslant e^{\mu(a-r)}Y_t$ together with (5.9). $\square$

Lemma 5.5. *Suppose $\mu \geqslant 1/2$ and $r \leqslant 9a/10$. For every $t \geqslant 0$ and $x \in [0,a]$, we have for large a,*

$$\mathbf{E}_f^x[N_t(r)^2] \leqslant Ce^{-f(t/a^2)+O(\mathrm{Err}(f,t))}\left(\frac{e^{\mu a}}{a^3}\right)^2(1+r^2)\Big(1+\frac{1+r^4}{a^2}\Big)e^{-2\mu r}\Big(w_Z(x)\frac{t}{a^3} + w_Y(x)\Big) \tag{5.32}$$

Proof. As in the proof of (5.8), we have by Lemmas 3.5 and 5.1,

$$\mathbf{E}^x[N_t(r)^2] \leqslant e^{-f(t/a^2)+O(\mathrm{Err}(f,t))}\Big(\mathbf{E}^x[N_t(r)] + \beta_0 m_2\int_0^a\int_0^t \mathbf{p}_s(x,z)(\mathbf{E}^z[N_{t-s}(r)])^2\,\mathrm{d}s\,\mathrm{d}z\Big). \tag{5.33}$$

By (5.3), Lemma 5.4 and the hypotheses on x and r, we have after a change of variables $z \to a - z$ in the integral,

$$\int_0^a \mathbf{p}_s(x,z)(\mathbf{E}^z[N_{t-s}(r)])^2\,\mathrm{d}z \leqslant C\left(\frac{e^{\mu a}}{a^3}\right)^2 e^{\mu(x-a-2r)+\pi^2 s/(2a^2)}$$
$$\times\Big((1+r^2)\int_0^a p_s^a(a-x,z)e^{-\mu z}\Big(z^2+\frac{1+r^4}{a^2}\Big)\,\mathrm{d}z + a^6\int_{a/20}^a p_s^a(a-x,z)e^{-\mu z}\,\mathrm{d}z\Big). \tag{5.34}$$

Integrating (5.34) over s from 0 to t and splitting the interval at $t \wedge a^2$, we have by (2.8) (for the first piece) and (2.7) (for the second),

$$\int_0^a \int_0^t \mathbf{p}_s(x,z)(\mathbf{E}^z[N_{t-s}(r)])^2 \,\mathrm{d}s\,\mathrm{d}z \leqslant C\left(\frac{e^{\mu a}}{a^3}\right)^2 \times (1+r^2)\Big(1+\frac{1+r^4}{a^2}\Big)e^{-2\mu r}(1+a^7e^{-\mu a/20})\Big(w_Z(x)\frac{t}{a^3}+w_Y(x)\Big) \tag{5.35}$$

The lemma follows from (5.33) and (5.35), together with the second statement of Lemma (5.4) and the hypotheses on μ and r. □

5.4 The particles hitting the right border

In this section we recall some formulae from [7] about the number of particles hitting the right border of the interval. We reprove these formulae here for completeness and because Lemma 2.1 makes their proofs straightforward. For most formulae we will assume that $f \equiv 0$, i.e. that we are working under the measure $\mathbf{P}$. Only Lemma 5.7 contains an upper bound on the expected number of particles for general f, which will be useful in Section 7.

For a measurable subset $S \subset \mathbf{R}$, define R_S to be the number of particles killed at the right border during the (time) interval S, i.e.

$$R_S = \#\{(u,t) : u \in \mathscr{N}(t) \text{ and } H_0(X_u) > H_a(X_u) = t \in S\}.$$

Lemma 5.6. *For any initial configuration ν and any $0 \leqslant s \leqslant t$, we have*

$$|\mathbf{E}[R_{[s,t]}] - \frac{\pi(t-s)}{a^3}Z_0| \leqslant C_{5.36}\Big(Y_0 \wedge E_{s/a^2}(1\wedge (t-s)/a^3)Z_0\Big), \tag{5.36}$$

where E_s is defined in (2.6)*. Furthermore, if $\mu \geqslant 1/2$ and $0 \leqslant t \leqslant a^3$, then for each $x \in (0,a)$,*

$$\mathbf{E}^x[R_t^2] \leqslant C_{5.37}\Big(\frac{t}{a^3}w_Z(x) + w_Y(x)\Big), \tag{5.37}$$

Proof. Recall that for a real-valued process, H_0 and H_a denote the hitting time functionals of 0 and a and $H = H_0 \wedge H_a$. Note that $W^x_{-\mu}(H < \infty) = 1$ for all $x \in [0,a]$. By Lemma 3.2, we then have

$$\mathbf{E}^x[R_S] = \mathbf{E}^x\Big[\sum_{(u,t)\in\mathscr{L}_H} \mathbf{1}_{(X_u(t)=a,\, t\in S)}\Big] = W^x_{-\mu}\Big[e^{\beta_0 m H_a}\mathbf{1}_{(H_0>H_a\in S)}\Big] = e^{\mu(x-a)}I^a(x,S), \tag{5.38}$$

where the last equality follows from Girsanov's transform and (2.10). Equation (5.36) now follows Lemma 2.1. The second-moment estimate is obtained from Lemma 3.3 and (5.38) by a calculation similar to the one in the proof of Proposition 5.2 (see [47, Lemma 5.8] for details). □

Lemma 5.7. *Let f be a function as in Section 5.1. Then for every $x \in (0,a)$, we have*

$$\mathbf{E}^x_f[R_S] \leqslant \mathbf{E}^x[R_S].$$

Proof. As in the proof of Lemma 5.6, we have

$$\mathbf{E}^x_f[R_S] = W^x_{-\mu_t}\Big[e^{\beta_0 m H_a}\mathbf{1}_{(H_0>H_a\in S)}\Big] = W^x_0\Big[e^{-\int_0^{H_a}\mu_t\,\mathrm{d}B_t - \int_0^{H_a}\mu_t^2/2\,\mathrm{d}t + H_a/2}\mathbf{1}_{(H_0>H_a\in S)}\Big],$$

by Girsanov's theorem and the definition of β_0. On the event $\{H_0 > H_a\}$, we now have by integration by parts,

$$\int_0^{H_a} \mu_t\,\mathrm{d}B_t = \mu(a-x) + a(\mu_{H_a} - \mu) - \int_0^{H_a} B_t\,\mathrm{d}\mu_t \geqslant \mu(a-x),$$

since $B_t \in [0,a]$ for $t \in [0,H_a]$. This gives

$$\mathbf{E}_f^x[R_S] \leqslant e^{\mu(x-a)} W_0^x\Big[e^{(1-\mu^2)H_a/2}\mathbf{1}_{(H_0>H_a\in S)}\Big] = \mathbf{E}^x[R_S],$$

by the proof of Lemma 5.6. □

We finish this section with a lemma which links BBM with absorption at a critical line to our BBM with selection model. The proof is simple, see [47, Lemma 2.5.10] for details.

Lemma 5.8. *Let $\zeta \geqslant 1$, $y \geqslant 1$, $\mu \geqslant 1/2$ and f be a barrier function (defined in Section 5.1). Suppose that $\sqrt{a} \geqslant y + \zeta$ and $\|f\| \leqslant \sqrt{a}$. Let $(x_i,t_i)_{i=1}^N$ be a collection of space-time points with*

$$x_i = a - y + (1-\mu)t_i - f(s/a^2), \quad i = 1,\dots,N,$$

and $t_i \leqslant \zeta$ for all i. Define $Z = \sum_i w_Z(x_i)$, $Y = \sum_i w_Y(x_i)$ and $W_y = ye^{-y}N$. Then,

$$Z = \pi W_y\Big(1 + O\Big(\frac{1}{a}\Big)\Big) \quad \text{and} \quad Y = \frac{1}{y}W_y\Big(1 + O\Big(\frac{1}{a}\Big)\Big).$$

In particular, for large a, we have

$$Y \leqslant Z/y.$$

5.5 Penalizing the particles hitting the right border

In this section, let $(U_u)_{u\in U}$ be iid random variables, uniformly distributed on $(0,1)$, independent of the branching Brownian motion. Furthermore, let $p : \mathbf{R}_+ \in (0,1]$ be measurable and such that $p(t) = 0$ for large enough t. Recall that $H = H_0 \wedge H_a$. We define the event

$$E = \{\nexists (u,t) \in \mathscr{L}_H : X_u(t) = a \text{ and } U_u \leqslant p(t)\}.$$

In order to describe the law $\widetilde{\mathbf{P}}_f^x = \mathbf{P}_f^x(\cdot|E)$, set

$$h(x,t) = \mathbf{P}_f^{(x,t)}(E), \quad Q(x,t) = \sum_{k=0}^{\infty} q(k)h(x,t)^{k-1}, \quad \tilde{q}(x,t,k) = q(k)h(x,t)^{k-1}/Q(x,t) \tag{5.39}$$

The law $\widetilde{\mathbf{P}}_f^x$ can be seen as a Doob transform of $\mathbf{P}_f^x$ by the space-time harmonic function $h(x,t)$. Under $\widetilde{\mathbf{P}}_f^x$, the BBM stopped at $\mathscr{L}_H$ is again a branching Markov process where (see [47, Section 3.4] and the errata in the arXiv version)

- particles move according to the particles move according to the law obtained from Brownian motion with drift $-\mu_t$ (stopped at 0 and a) through a change of measure by the martingale

$$h(X_{t\wedge H}, t\wedge H)\exp\left(-\int_0^{t\wedge H}\beta_0(1-Q(X_s,s))\,\mathrm{d}s\right).$$

- a particle located at the point $x \in (0,a)$ at time t branches at rate $\beta_0 Q(x,t)\mathbf{1}_{x\in(0,a)}$, throwing k offspring with probability $\tilde{q}(x,t,k)$.

Together with Lemma 3.2, this immediately gives the following useful Many-to-one lemma for the conditioned process stopped at the stopping line $\mathscr{L}_t = \mathscr{L}_{H\wedge t}$: Define the function

$$e(x,t) = \beta_0\sum_{k\geqslant 0}(k+1)(1-h(x,t)^{k-1})q(k) \leqslant \beta_0(m_2+m)(1-h(x,t)). \tag{5.40}$$

Lemma 5.9. *For any measurable function $g : [0,a] \to \mathbf{R}_+$, $x \in (0,a)$ and $t \geqslant 0$, we have*

$$\widetilde{\mathbf{E}}^x_f\Big[\sum_{u\in\mathscr{L}_t} g(X_u(t))\Big] = W^x_{-\mu_t}\Big[g(X_{H\wedge t})\frac{h(X_{H\wedge t}, H\wedge t)}{h(x,0)}e^{(H\wedge t)/2-\int_0^{H\wedge t} e(X_s,s)\,\mathrm{d}s}\Big]. \tag{5.41}$$

In particular, if we denote by $\widetilde{\mathbf{p}}^f_t(x,y)$ the density of the $\widetilde{\mathbf{P}}^x_f$-BBM, then

$$\widetilde{\mathbf{p}}_t(x,y) = \frac{h(y,t)}{h(x,0)}\mathbf{p}_t(x,y)W^{x,t,y}_{\text{taboo}}\Big[e^{-\int_0^t e(X_s,s)\,\mathrm{d}s}\Big] \leqslant (h(x,0))^{-1}\mathbf{p}_t(x,y), \tag{5.42}$$

and for general f,

$$\widetilde{\mathbf{p}}^f_t(x,y) \leqslant (h(x,0))^{-1}\mathbf{p}^f_t(x,y). \tag{5.43}$$

This lemma immediately gives an upper bound for the quantities we are interested in:

Corollary 5.10. *Let $x \in (0,a)$, $t \geqslant 0$ and $g : [0,a] \to \mathbf{R}_+$ be measurable with $g(0) = g(a) = 0$. Define $S_t = \sum_{u\in\mathscr{L}_t} g(X_u(t))$. Then,*

$$\widetilde{\mathbf{E}}^x_f[S_t] \leqslant (h(x,0))^{-1}\mathbf{E}^x_f[S_t], \tag{5.44}$$

$$\widetilde{\mathbf{E}}^x_f[S_t^2] \leqslant (h(x,0))^{-1}\mathbf{E}^x_f[S_t^2]. \tag{5.45}$$

Proof. Equation (5.44) immediately follow from (5.42). In order to prove the second-moment estimates, we note that by Lemma 3.5 and the description of the conditioned process,

$$\widetilde{\mathbf{E}}^x_f[S_t^2] = \widetilde{\mathbf{E}}^x_f\Big[\sum_{u\in\mathscr{L}_t} g(X_u(t))^2\Big] + \int_0^t\int_0^a \widetilde{\mathbf{p}}_s(x,y)\widetilde{m_2}(y,s)\beta_0 Q(y,s)\left(\widetilde{\mathbf{E}}^{(y,s)}_f[S_t]\right)^2\,\mathrm{d}y\,\mathrm{d}s,$$

where $\widetilde{m_2}(x,t) = \sum_{k\geqslant 0} k(k-1)\tilde{q}(x,t,k)$. By (5.39), we have $\widetilde{m_2}(x,t)Q(x,t) \leqslant h(x,t)m_2$. Equation (5.45) then follows from (5.12), (5.42) and (5.44). ☐

The following lemma gives a good lower bound on the first-moment estimates in the case where $f \equiv 0$.

Lemma 5.11. *Suppose $\mu \geqslant 1/2$, $t \leqslant a^3$ and $p(s) = 0$ for all $s \geqslant a^3$. Let S_t be as in Corollary 5.10. We have*

$$\widetilde{\mathbf{E}}^x[S_t] \geqslant \mathbf{E}^x[S_t](1 - C_{5.46}\|p\|_\infty). \tag{5.46}$$

This follows from the following estimate on $h(x,0)$, which will be sharpened in Lemma 6.3.

Lemma 5.12. *Suppose $p(s) = 0$ for all $s \geqslant a^3$. Then for all $x \in (0,a)$, we have*

$$1 - h(a-x,0) \leqslant C\|p\|_\infty(x+1)e^{-\mu x}.$$

Proof. By Markov's inequality, we have

$$1 - h(x,0) \leqslant \mathbf{E}^x_f(\#\{(u,s) \in \mathscr{L}_{a^3} : X_u(s) = a,\ U_u \leqslant \|p\|_\infty\}) \leqslant \|p\|_\infty\mathbf{E}^x_f(R_{a^3}).$$

The lemma now follows from Lemmas 5.6 and 5.7 and the inequality $\sin x \leqslant x$, $x \in [0,\pi]$. ☐

Proof of Lemma 5.11. By (5.42),

$$\widetilde{\mathbf{E}}^x[S_t] \geqslant \mathbf{E}^x[S_t]\inf_{y\in(0,a)}\left(h(y,t)W^{x,t,y}_{\text{taboo}}\Big[e^{-\int_0^t e(X_s,s)\,\mathrm{d}s}\Big]\right). \tag{5.47}$$

By Lemma 5.12 and the hypothesis on μ, we have for every $y \in (0,a)$,

$$h(a-y,t) \geqslant h(a-y,0) \geqslant 1 - C\|p\|_\infty e^{-y/3}. \tag{5.48}$$

This gives

$$W^{x,t,y}_{\text{taboo}}\Big[e^{-\int_0^t e(X_s,s)\,\mathrm{d}s}\Big] \geqslant 1 - W^{x,t,y}_{\text{taboo}}\Big[\int_0^t e(X_s,s)\,\mathrm{d}s\Big] \geqslant 1 - C\|p\|_\infty, \tag{5.49}$$

by (5.40), (5.48), Lemma 2.2, the inequality $e^{-x} \geqslant 1-x$ for $x \geqslant 0$ and the fact that the law of the Brownian taboo process is preserved under the map $y \mapsto a-y$. The lemma now follows from (5.47), (5.48) and (5.49). $\square$

Finally, we study the law of R_t under the new probability.

Lemma 5.13. *We have for every $x \in [0,a]$,*

$$\mathbf{E}^x_f[R_t] - \|p\|_\infty \mathbf{E}^x_f[R_t^2] \leqslant \widetilde{\mathbf{E}}^x_f[R_t] \leqslant (h(x,0))^{-1}\mathbf{E}^x_f[R_t], \tag{5.50}$$

and if there is a $p \in [0,1]$, such that $p(s) \equiv p$ for $s \leqslant t$, then we even have

$$\widetilde{\mathbf{E}}^x_f[R_t] \leqslant \mathbf{E}_f[R_t]. \tag{5.51}$$

Proof. Define the stopping line $\mathscr{R}_t = \{(u,s) \in \mathscr{L}_{H_a} : s \leqslant t\}$. By the definition of the law $\widetilde{\mathbf{P}}$,

$$\widetilde{\mathbf{E}}^x_f[R_t] = \frac{\mathbf{E}^x_f\Big[R_t \prod_{(u,s)\in\mathscr{R}_t}(1-p(X_u(s)))\Big]}{\mathbf{E}^x_f\Big[\prod_{(u,s)\in\mathscr{R}_t}(1-p(X_u(s)))\Big]}. \tag{5.52}$$

Now the denominator is $h(x,0)$ by (5.39), which yields the right-hand side of (5.50). The left-hand side follows by noticing that

$$\mathbf{E}^x_f\Big[R_t \prod_{(u,s)\in\mathscr{R}_t}(1-p(X_u(s)))\Big] \geqslant \mathbf{E}^x_f[R_t(1-\|p\|_\infty)^{R_t}] \geqslant \mathbf{E}^x_f[R_t] - \|p\|_\infty \mathbf{E}^x_f[R_t^2].$$

For (5.51), we note that if $p(s) \equiv p$ for $s \leqslant t$, then by (5.52),

$$\widetilde{\mathbf{E}}^x_f[R_t] = \frac{\mathbf{E}^x_f[R_t(1-p)^{R_t}]}{\mathbf{E}^x_f[(1-p)^{R_t}]}.$$

Since $(1-p)^k$ is decreasing in k, this yields (5.51). $\square$

6 BBM with absorption before a breakout

In this section, we study branching Brownian motion with drift $-\mu$ and absorption at 0 until a *breakout* occurs, an event which will be defined in Section 6.1 and which corresponds to a particle going far to the right and spawning a large number of descendants. In (6.20), we decompose the system into a particle conditioned to break out at a specific time T (this particle will be called the *fugitive*) and the remaining particles, which are conditioned not to break out before time T. These two parts will be studied separately, the former in Section 6.4 and the latter in Section 6.3. Before that, in Section 6.2, we study the law of the time of the first breakout, showing that it is approximately exponentially distributed. First of all, however, we start with the necessary definitions:

6.1 Definitions

We will introduce several parameters which will be used during the rest of the paper. The two most important parameters are a and A, which are both large positive constants. The meaning of a is as in the previous sections: It is the right border of an interval in which the particles are staying most of the time. The parameter A has a more subtle meaning and controls the number of particles of the system and with it the intensity at which particles hit the point a. In Section 7, we will indeed choose the initial conditions such that $Z_0 \approx e^A$.

When we study the system for large A and a, we first let a go to infinity, then A. Thus, the statement "For large A and a we have…" means: "There exist A_0 and a function $a_0(A)$, both depending on the reproduction law q only, such that for $A \geqslant A_0$ and $a \geqslant a_0(A)$ we have…". Likewise, the statement "As A and a go to infinity…" means "For all A there exists $a_0(A)$, depending on the reproduction law q only, such that as A goes to infinity and $a \geqslant a_0(A)$…". These phrases will become so common that in Sections 7 to 9 they will often be used implicitly, although they will always be explicitly stated in the theorems, propositions, lemmas etc. We further introduce the notation $o(1)$, which stands for a (non-random) term that only depends on the reproduction law q and the parameters A, a, ε, η, y and ζ and which goes to 0 as A and a go to infinity.

The remaining parameters we introduce are all going to depend on A, but not on a. First of all, there is the small parameter ε, which controls the intensity of the breakouts. Indeed, when $Z_0 \approx e^A$, the mean time one has to wait for a breakout will be approximately proportional in ε. Morally, one could choose ε such that $e^{-A/2} \ll \varepsilon \ll A^{-1}$, but for technical reasons we will require that

$$\varepsilon \leqslant C_{6.1} A^{-17}, \quad \text{and} \tag{6.1}$$

$$\varepsilon \geqslant C_{6.2} e^{-A/6}. \tag{6.2}$$

Another protagonist is η, which we will choose as small as we need and which will be used to bound the probability of very improbable events, as well as the contribution of the variable Y. It will be enough to require that

$$\eta \leqslant e^{-2A}, \tag{6.3}$$

which, by (6.2), implies

$$\eta \leqslant C\varepsilon^{12}. \tag{6.4}$$

The last parameters are y and ζ, which are defined as in Lemma 4.2, with η there being the η defined above. Note that the parameters η, y and ζ appeared already in [7] and had the same meaning there.

We can now proceed to the definition of the process. Recall the definition of μ in (5.1). As in Section 5.1, we denote by $\mathbf{P}^x$ and $\mathbf{E}^x$ the law and expectation of branching Brownian motion with drift $-\mu$ starting from a particle at the point $x \in \mathbf{R}$; we extend this definition to general initial distributions of particles according to a counting measure ν. Recall from Section 3.1 that $\mathscr{N}(t)$ denotes the set of individuals alive at time t. We want to absorb the particles at 0 and do this formally by setting

$$\mathscr{N}_0(t) = \{u \in \mathscr{N}(t) : H_0(X_u) > t\},$$

where H_0 is again the hitting time functional of 0.

Instead of absorbing particles at a, we are now going to classify them into *tiers* as described in the introduction. Let $u \in U$, $t \geqslant 0$. We define two sequences of random times $(\tau_n(u))_{n\geqslant -1}$ and $(\sigma_n(u))_{n\geqslant 0}$ by $\tau_{-1}(u) = 0$, $\sigma_0(u) = 0$ and for $n \geqslant 0$:

$$\begin{aligned} \tau_n(u) &= \inf\{s \geqslant \sigma_n(u) : X_u(s) = a\}, \\ \sigma_{n+1}(u) &= \inf\{s \geqslant \tau_n(u) : X_u(s) = a - y + (1-\mu)(s - \tau_n(u))\}, \end{aligned} \tag{6.5}$$



Figure 2: Left picture: A graphical view of the *tiers*. The trajectory of one particle u is singled out (thick line) and the times $\tau_l(u)$ and $\sigma_l(u)$ are shown. The tier 0 particles are drawn with straight lines, the tier 1 particles with dashed and the tier 2 particles with dotted lines. Right picture: The stopping line $\mathscr{N}_t$ (encircled particles).

where we set $\inf \varnothing = \infty$. See Figure 2 for a graphical description. We now define for $t \geqslant 0$ the stopping lines

$$\mathscr{R}_t^{(l)} = \{(u,s) \in U \times [0,t] : s = \tau_l(u) \text{ and } u \in \mathscr{N}_0(s)\},\ l \geqslant -1, \text{ and} \tag{6.6}$$

$$\mathscr{S}_t^{(l)} = \{(u,s) \in U \times [0,t] : s = \sigma_l(u) \text{ and } u \in \mathscr{N}_0(s)\},\ l \geqslant 0. \tag{6.7}$$

That means, $\mathscr{R}_t^{(l)}$ contains the particles of tier l at the moment at which they touch the right barrier and $\mathscr{S}_t^{(l)}$ contains the particles of tier l at the moment at which they come back to the critical line. Note that the sets $\mathscr{R}_t^{(l)}$ and $\mathscr{S}_t^{(l)}$ are increasing in t and $\mathscr{R}_t^{(l-1)} \preceq \mathscr{S}_t^{(l)} \preceq \mathscr{R}_t^{(l)}$ for every $l \geqslant 0$. We also set

$$R_t^{(l)} = \#\mathscr{R}_t^{(l)}.$$

In order to extend the definitions of the variables Z_t and Y_t to the current setup, we could simply replace $\mathscr{N}_{0,a}(t)$ by $\mathscr{N}_0(t)$ in their definition (see Section 5.1). However, it will be more useful to take special care of the individuals u for which $\tau_l(u) \leqslant t < \sigma_{l+1}(u)$ for some $l \geqslant 0$, since these are in some kind of “intermediary” state which is difficult to analyse. We therefore define the stopping lines

$$\begin{aligned}\mathscr{N}_t^{(l)} = \{(u,s) \in U \times \mathbf{R}_+ : u \in \mathscr{N}_0(s),\ \tau_{l-1}(u) \leqslant s < \tau_l(u),\\ \text{and either } t < \sigma_l(u) = s \text{ or } \sigma_l(u) \leqslant t = s\},\ l \geqslant 0,\end{aligned} \tag{6.8}$$

and

$$\mathscr{N}_t = \bigcup_{l \geqslant 0} \mathscr{N}_t^{(l)}. \tag{6.9}$$

In other words, the stopping line $\mathscr{N}_t^{(l)}$ contains the particles of tier l that have already come back to the critical line at time t, as well as the descendants of those that haven't, at the moment at which they hit the critical line. We then define for $l \geqslant 0$ (recall the definitions of w_Z and w_Y from Section 5),

$$Z_t^{(l)} = \sum_{u \in \mathscr{N}_t^{(l)}} w_Z(X_u(t)), \quad Y_t^{(l)} = \sum_{u \in \mathscr{N}_t^{(l)}} w_Y(X_u(t)),$$

Furthermore, for any symbol S and $0 \leqslant k \leqslant l \leqslant \infty$, we write,

$$S^{(k;l)} = \sum_{i=k}^{l} S^{(i)}, \quad S^{(l+)} = S^{(l;\infty)}, \quad S = S^{(0+)}.$$

For a particle $(u,s) \in \mathscr{R}_t^{(l)}$, we now define the stopping line

$$\mathscr{S}^{(u,s)} = \{(v,r) \in U \times \mathbf{R}_+ : v \in \mathscr{N}_0(r),\ (u,t) \preceq (v,r) \text{ and } r = \sigma_{l+1}(v)\}.$$

This stopping line yields a collection $(X_v(r), r-s)_{(v,r)\in\mathscr{S}^{(u,s)}}$ of space-time points and we denote by $Z^{(u,s)}$, $Y^{(u,s)}$ and $W_y^{(u,s)}$ the quantities from Lemma 5.8 corresponding to this collection of points (in particular, $W_y^{(u,s)} = ye^{-y}\#\mathscr{S}^{(u,s)}$). Of course, we have chosen the stopping line in such a way that the variable $W_y^{(u,s)}$ follows the same law as the variable W_y defined in (4.2). We also define $\tau_{\max}^{(u,s)} = \max_{(v,r)\in\mathscr{S}^{(u,s)}}(r-s)$. We then define the "good" event

$$\begin{aligned} G^{(u,s)} = \{\#\mathscr{S}^{(u,s)} \leqslant \zeta\} \cap \{\mathscr{S}^{(u,s)} \subseteq U \times [1,\zeta]\} \\ \cap \{\sup_{s\leqslant r\leqslant\zeta} \#\{v \in \mathscr{N}_0(r) : (u,s) \preceq (v,r) \preceq \mathscr{S}^{(u,s)}\} \leqslant \zeta\} \end{aligned} \tag{6.10}$$

and the event of a *breakout*,

$$B^{(u,s)} = \{Z^{(u,s)} > \varepsilon e^A\} \cup (G^{(u,s)})^c, \tag{6.11}$$

(the inclusion of the "bad" event $(G^{(u,s)})^c$ is for technical reasons). If the event $B^{(u,s)}$ occurs, the particle u is then also called the *fugitive*. We set

$$p_B = \mathbf{P}^a(B^{(\varnothing,0)}), \tag{6.12}$$

and define the law of BBM started at a with the first particle conditioned not to break out:

$$\mathbf{Q}^a(\cdot) = \mathbf{P}^a(\cdot \mid B^c) = \frac{\mathbf{P}^a(\cdot,\ B^c)}{1-p_B},$$

where we set $B = B^{(\varnothing,0)}$. We further set $Z = Z^{(\varnothing,0)}$ and $W_y = W_y^{(\varnothing,0)}$ and note that by Lemmas 5.8 and 4.2, we have for large a,

$$\mathbf{P}^a(|Z - \pi W| > 2\eta) + \mathbf{P}^a((G^{(\varnothing,0)})^c) < \eta, \tag{6.13}$$

where W is defined as in (4.2). Hence, by (6.3), (4.5) and Lemma 4.2, we get

$$p_B = \Big(\pi + o(1)\Big)\frac{1}{\varepsilon e^A}, \tag{6.14}$$

which goes to 0 as A and a go to infinity, by (6.2). Furthermore, (6.13) yields for large A and a,

$$\begin{aligned} \mathbf{Q}^a[Z] &= (\mathbf{E}[\pi W \mathbf{1}_{(\pi W \leqslant \varepsilon e^A(1+o(1))+O(\eta))}] + O(\eta\varepsilon e^A))(1+O(p_B)) \\ &= \pi(A + \log\varepsilon + c_{6.15} + o(1)), \end{aligned} \tag{6.15}$$

by (4.6), (6.1), (6.3) and (6.14). In particular, we have for $A \geqslant 1$ and large a,

$$\mathbf{Q}^a[Z] \leqslant CA. \tag{6.16}$$

Moreover, by (4.5), (6.13) and (6.3), we have for $A \geqslant 1$ and large a,

$$\mathbf{Q}^a[Z^2] \leqslant C\varepsilon e^A. \tag{6.17}$$

Finally, note that by Lemma 5.8, we have $Y \leqslant \eta Z$, $\mathbf{Q}^a$-almost surely, a fact that will often be used without further reference.

We now define for every $l \in \mathbf{N}$ the time of the first breakout of a particle of tier l,

$$T^{(l)}(\omega) = \inf\{t \geqslant 0 : \omega \in \bigcup_{(u,s)\in\mathscr{R}_t^{(l)}} B^{(u,s)}\}, \tag{6.18}$$

and set

$$T^{(0;l)} = \min_{0\leqslant j\leqslant l} T^{(l)}, \quad \text{with } T = T^{(0;\infty)} = \min_{j\geqslant 0} T^{(l)}. \tag{6.19}$$

Now fix $t > 0$ and $l \in \mathbf{N} \cup \{\infty\}$. We want to describe the system conditioned on $T^{(0;l)} = t$. For this, suppose that at time 0 the particles are distributed according to a counting measure $\nu = \sum_{i=1}^n \delta_{x_i}$. We denote by $\mathscr{U}$ the fugitive of the breakout that happened at time $T^{(0;l)}$ and define $p_i = \mathbf{P}^\nu(i \preceq \mathscr{U} \,|\, T^{(0;l)} = t)$. This yields a law $(p_i)_{i=1}^n$ on the initial particles, depending on ν and t. Since the variable $T^{(0;l)}$, the time of the first breakout, is the minimum of the variables $T_i^{(0;l)}$, $i = 1, \dots, n$, the times of the first breakout of the BBM descending from the particle i, we can decompose the process into

$$\mathbf{P}^\nu\Big(\prod_{i=1}^n \mathrm{d}\omega^{(i)} \,\Big|\, T^{(0;l)} = t\Big) = \sum_{i=1}^n p_i \times \mathbf{P}^{x_i}(\mathrm{d}\omega^{(i)} \,|\, T^{(0;l)} = t) \times \prod_{j\neq i} \mathbf{P}^{x_j}(\mathrm{d}\omega^{(j)} \,|\, T^{(0;l)} > t). \tag{6.20}$$

That is, we first choose according to the law $(p_i)_{i=1}^n$ the initial particle that is going to cause the breakout. This particle spawns a BBM conditioned to break out at time t. The remaining particles spawn independent BBM conditioned not to break out before time t.

Remark 6.1. Note that many results in this section can be done for BBM with varying drift given by a barrier function f. For example, with Lemma 5.7 one gets immediately that $\mathbf{P}_f(T^{(0)} > t) \geqslant \mathbf{P}(T^{(0)} > t)$ for all $t \geqslant 0$ and it will be clear from the next section that in fact $\mathbf{P}_f(T > t) \geqslant \mathbf{P}(T > t)$ as well. However, in order to simplify notation and because we will not need the results often in this generality, we state them here only for BBM with fixed drift.

6.2 The time of the first breakout

For large A and a, define

$$t_{6.21} = \tfrac{1}{20A} a^3. \tag{6.21}$$

We want to prove that the random variable T defined in the previous section is approximately exponentially distributed with parameter $p_B \pi Z_0/a^3$, which is the statement of the following proposition:

Proposition 6.2. *Let* $0 \leqslant t \leqslant t_{6.21}$. *For A and a large enough, we have*

$$\mathbf{P}(T > t) = \exp\Big(-\pi p_B Z_0 \frac{t}{a^3}\Big(1 + O(At/a^3 + p_B)\Big) + O(p_B Y_0)\Big). \tag{6.22}$$

The proof proceeds by a sequence of lemmas. Lemma 6.3 gives a estimate on $\mathbf{P}(T^{(0)} > t)$. This is used in Lemma 6.4, in order to obtain an estimate on $\mathbf{Q}^a(T > t)$, using a recursive argument. Finally, Proposition 6.2 is proven by combining Lemmas 6.3 and 6.4.

Lemma 6.3. *Let $0 \leqslant t \leqslant a^3$. Suppose that $p_B \leqslant 1/2$. Then,*

$$\mathbf{P}(T^{(0)} > t) = \exp\Big(- \pi p_B Z_0 \frac{t}{a^3}\big(1 + O(p_B)\big) + O(p_B Y_0)\Big). \tag{6.23}$$

Proof. Let $x_1, \ldots, x_n$ be the positions of the initial particles. Since the initial particles spawn independent branching Brownian motions, we have

$$\mathbf{P}(T^{(0)} > t) = \prod_i \mathbf{P}^{x_i}(T^{(0)} > t). \tag{6.24}$$

We have for every $x \in (0, a)$,

$$\mathbf{P}^x(T^{(0)} > t) = \mathbf{E}^x\Big[\prod_{(u,s)\in\mathscr{R}_t^{(0)}} \mathbf{1}_{B(u,s)}\Big] = \mathbf{E}^x\Big[(1 - p_B)^{R_t^{(0)}}\Big], \tag{6.25}$$

since by the strong branching property, the random variables $Z^{(u,s)}$ are independent conditioned on $\mathscr{R}_t^{(0)}$. By Lemma 5.6 and the assumption $t \leqslant a^3$, we have

$$|\mathbf{E}^x[R_t^{(0)}] - \pi w_Z(x)t/a^3| \leqslant C w_Y(x), \tag{6.26}$$

$$\mathbf{E}^x[(R_t^{(0)})^2] \leqslant C(w_Z(x)t/a^3 + w_Y(x)). \tag{6.27}$$

By Jensen's inequality, (6.26) and the inequality $|\log(1-z)| \leqslant z + z^2$ for $z \leqslant 1/2$,

$$\mathbf{E}^x\Big[(1 - p_B)^{R_t^{(0)}}\Big] \geqslant \exp\Big(- \pi p_B w_Z(x) \frac{t}{a^3}(1 + Cp_B) - Cp_B w_Y(x)\Big). \tag{6.28}$$

Furthermore, the inequality $(1-p)^n \leqslant 1 - np + n(n-1)p^2/2$ and Equations (6.26) and (6.27) give

$$\mathbf{E}^x\Big[(1 - p_B)^{R_t^{(0)}}\Big] \leqslant 1 - \pi p_B w_Z(x) \frac{t}{a^3}(1 - Cp_B) + Cp_B w_Y(x). \tag{6.29}$$

The lemma now follows from (6.25), (6.28) and (6.29) together with the inequality $1 + z \leqslant e^z$ for $z \in \mathbf{R}$. ☐

In the following lemma, note that according to the definition of the tiers, a particle starting at a starts immediately in tier 1.

Lemma 6.4. *Let $0 \leqslant t \leqslant t_{6.21}$. Then, for large A and a,*

$$\mathbf{Q}^a(T > t) \geqslant \exp\Big(- Cp_B A(\tfrac{t}{a^3} + \eta)\Big). \tag{6.30}$$

Proof. We have

$$\mathbf{Q}^a(T > t) = \mathbf{Q}^a\Big[\prod_{(u,s)\in\mathscr{S}_t^{(1)}} \mathbf{P}^{X_u(s)}(T > t - s)\Big] \geqslant \mathbf{Q}^a[\mathbf{P}^\nu(T > t)], \tag{6.31}$$

where $\nu = \sum_{(u,s)\in\mathscr{S}_t^{(1)}} \delta_{X_u(s)}$. Since $T > t$ implies $T^{(0)} > t$, we have

$$\mathbf{P}^\nu(T > t) = \mathbf{P}^\nu(T > t | T^{(0)} > t)\mathbf{P}^\nu(T^{(0)} > t). \tag{6.32}$$

Let $Z = Z^{(\varnothing,0)}$ and $Y = Y^{(\varnothing,0)}$, such that $Y \leqslant \eta Z$, $\mathbf{Q}^a$-almost surely, by Lemma 5.8 and the definition of the "bad" event $B^{(\varnothing,0)}$. By Lemma 6.3, we have for large A,

$$\mathbf{P}^\nu(T^{(0)} > t) \geqslant \exp\Big(- Cp_B Z\left(\tfrac{t}{a^3} + \eta\right)\Big). \tag{6.33}$$

Furthermore, with the notation from Section 5.5, with $p(s) \equiv p_B$,

$$\mathbf{P}^\nu(T > t | T^{(0)} > t) = \widetilde{\mathbf{P}}^\nu(T > t) = \widetilde{\mathbf{P}}^\nu\Big[\prod_{(u,s)\in\mathscr{R}_t^{(0)}} \mathbf{Q}^a(T > t - s)\Big] \geqslant \widetilde{\mathbf{P}}^\nu\Big[\mathbf{Q}^a(T > t)^{R_t^{(0)}}\Big].$$

By Jensen's inequality and Lemmas 5.13 and 5.6, this implies

$$\mathbf{P}^\nu(T > t | T^{(0)} > t) \geqslant \mathbf{Q}^a(T > t)^{\widetilde{\mathbf{E}}^\nu[R_t^{(0)}]} \geqslant \mathbf{Q}^a(T > t)^{\pi Z(t/a^3 + O(\eta))}. \tag{6.34}$$

Equations (6.31), (6.32), (6.33) and (6.34), together with Jensen's inequality and (6.15), now yield for large A and a,

$$\mathbf{Q}^a(T > t) \geqslant \mathbf{Q}^a(T > t)^{\pi^2 A(t/a^3 + O(\eta))} \times \exp\Big(- Cp_B A\left(\tfrac{t}{a^3} + \eta\right)\Big). \tag{6.35}$$

By the hypothesis on t and (6.3), the exponent of $\mathbf{Q}^a(T > t)$ in (6.35) is smaller than 1/2 for large A and a. This yields the statement. □

Proof of Proposition 6.2. The upper bound follows from Lemma 6.3 and the trivial inequality $T \leqslant T^{(0)}$. For the lower bound, we note that as in the proof of Lemma 6.4, we have by Jensen's inequality and Lemma 5.13,

$$\mathbf{P}(T > t) = \mathbf{P}(T > t \,|\, T^{(0)} > t)\mathbf{P}(T^{(0)} > t) \geqslant \mathbf{Q}^a(T > t)^{\mathbf{E}[R_t^{(0)}]}\mathbf{P}(T^{(0)} > t). \tag{6.36}$$

By Lemmas 6.4 and 5.6, we have

$$\mathbf{Q}^a(T > t)^{\mathbf{E}[R_t^{(0)}]} \geqslant \exp\Big(- Cp_B A(\tfrac{t}{a^3} + \eta)(\tfrac{t}{a^3}Z_0 + Y_0)\Big), \tag{6.37}$$

The lower bound in (6.22) now follows from (6.36), (6.37) and Lemma 6.3, together with the hypothesis on t, (6.3) and (6.14). □

Lemma 6.5. *Define* $\gamma = (\pi p_B Z_0)^{-1}$. *Suppose that* $Y_0 \leqslant C$ *and let* $\alpha \geqslant 0$ *and* $n \in \mathbf{N}$. *Then, for large* A, *for every* $l \in \mathbf{N} \cup \{\infty\}$,

$$\mathbf{E}[(T^{(0;l)}/a^3 + \alpha)^n \mathbf{1}_{(T^{(0;l)} \leqslant a^3)}] \leqslant C \sum_{k=0}^n \frac{n!\alpha^k (2\gamma)^{n-k}}{k!} \tag{6.38}$$

Furthermore, if $0 \geqslant \delta = A^{-1}o(1)$, *then for large* A *and* a,

$$\mathbf{E}[(T/a^3)\mathbf{1}_{(T \leqslant \delta a^3)}] = \gamma(1 + O(A\delta + p_B)) + O((\delta + \gamma)e^{-O(\delta/\gamma)}) \tag{6.39}$$

Proof. We first note that we have, for $n \geqslant 1$ and $\rho > 0$,

$$\int_0^\infty (t + \alpha)^{n-1} e^{-\rho t}\,\mathrm{d}t = \sum_{k=0}^{n-1} \frac{(n-1)!\alpha^k}{k!\rho^{n-k}}. \tag{6.40}$$

Now, we have

$$\begin{aligned}\mathbf{E}[(T^{(0;l)}/a^3 + \alpha)^n \mathbf{1}_{(T^{(0)} \leqslant a^3)}] &= \int_0^1 (t + \alpha)^n\,\mathbf{P}(T^{(0;l)}/a^3 \in \mathrm{d}t)\\ &\leqslant n\int_0^1 (t + \alpha)^{n-1}\,\mathbf{P}(T^{(0;l)} > ta^3)\,\mathrm{d}t + \alpha^n.\end{aligned}$$

The inequality (6.38) now follows from Lemma 6.3 and (6.40) and the hypothesis on Y_0. For the second part, we note that

$$\mathbf{E}[(T/a^3)\mathbf{1}_{(T\leqslant\delta a^3)}] = \int_0^\delta \mathbf{P}(T > ta^3)\,\mathrm{d}t - \delta\mathbf{P}(T > \delta a^3),$$

and by Proposition 6.2 and the hypothesis on Y_0, we have for $t \leqslant \delta$ and large A and a,

$$\mathbf{P}(T > ta^3) = (1 + O(p_B))\exp(-\gamma^{-1}t(1 + O(A\delta + p_B))).$$

Equation (6.39) now follows from the last two equations. □

We now show how we can couple the variable T with an exponentially distributed variable:

Lemma 6.6. *Suppose that $e^{-A}Z_0 = 1 + O(\varepsilon^{3/2})$ and that $Y_0 \leqslant \eta Z_0$. Then there exists a coupling (T, V), such that V is a random variable which is exponentially distributed with parameter $\pi p_B e^A$, T is $\sigma(V)$-measurable and $\mathbf{P}(|T/a^3 - V| > \varepsilon^{3/2}) \leqslant C\varepsilon^2$ for large A and a.*

Proof. For brevity, set $\rho := \pi p_B e^A$. Let F be the tail distribution function of T, i.e. $F(t) := \mathbf{P}(T \geqslant t)$. The number of individuals in a BBM tree being at most countable, T has no atoms except ∞. We can therefore define a random variable U which is uniformly distributed on $(0, 1)$ by setting

$$U = F(T)\mathbf{1}_{(T<\infty)} + U'F(\infty)\mathbf{1}_{(T=\infty)},$$

where U' is a uniformly distributed random variable on $(0, 1)$, independent of T. Now define $V = -\rho^{-1}\log U$. Then V is exponentially distributed with parameter ρ and $T = F^{-1}(e^{-\rho V})$, where F^{-1} denotes the generalised right-continuous inverse of F. Hence, T is $\sigma(V)$-measurable. On $\{T < \infty\}$, we have by Proposition 6.2, for a large enough,

$$\begin{aligned} V &= -\rho^{-1}(\pi p_B e^A e^{-A} Z_0 T/a^3(1 + O(AT/a^3 + p_B))) + O(p_B Y_0) \\ &= T/a^3(1 + O(\varepsilon^{3/2} + AT/a^3 + p_B)) + O(p_B e^A \eta), \end{aligned} \tag{6.41}$$

by the hypotheses on Z_0 and Y_0. Hence, by (6.2), (6.4) and (6.14), we have for a large enough,

$$|T/a^3 - V| = O(\varepsilon^{3/2}T/a^3 + A(T/a^3)^2) + O(\varepsilon^2).$$

But now we have by Lemma 6.3, for large A and a,

$$\mathbf{P}(T/a^3 > \varepsilon^{7/8}) \leqslant \mathbf{P}(T^{(0)}/a^3 > \varepsilon^{7/8}) \leqslant Ce^{-O(\varepsilon^{-1/8})} \leqslant \varepsilon^2/2. \tag{6.42}$$

The statement now follows from (6.41) and (6.42) together with (6.1) □

6.3 The particles that do not participate in the breakout

In this section, we fix $\tau \leqslant t_{6.21}$. We define the law of BBM conditioned not to break out before τ in the tiers $0, \dots, l$ by

$$\widehat{\mathbf{P}}_l(\cdot) = \mathbf{P}(\cdot \,|\, T^{(0;l)} > \tau), \quad \text{with } \widehat{\mathbf{P}} = \widehat{\mathbf{P}}_\infty.$$

Expectation w.r.t. $\widehat{\mathbf{P}}_l$ is denoted by $\widehat{\mathbf{E}}_l$. Under the law $\widehat{\mathbf{P}}_l$, the process stopped at $\mathscr{L}_H$ then follows the law $\widetilde{\mathbf{P}}$ from Section 5.5, with

$$p(t) = p_B\mathbf{1}_{(t\leqslant\tau)} + (1 - p_B)\mathbf{Q}^a(T^{(0;l)} \leqslant \tau - t), \tag{6.43}$$

such that by Lemma 6.4, (6.3), (6.16) and the trivial inequality $T^{(0;l)} \geqslant T$, we have for large A and a,

$$\|p\|_\infty \leqslant Cp_B. \tag{6.44}$$

In particular, by Lemma 5.12, (5.2) and (6.44), we have for large A and a,

$$(h(x, 0))^{-1} \leqslant 1 + Cp_B. \tag{6.45}$$

Lemmas 5.9 and 5.11, together with (6.44) and (6.45) now immediately imply the following:

Corollary 6.7. *Let* $x \in (0,a)$*,* $0 \leqslant t \leqslant \tau \leqslant t_{6.21}$ *and* $g : [0,a] \to \mathbf{R}_+$ *be measurable with* $g(0) = g(a) = 0$*. Define* $S_t^{(0)} = \sum_{u \in \mathscr{N}_t^{(0)}} g(X_u(t))$ *or* $S_t^{(0)} = R_t^{(0)}$*. Then, with* $\widehat{\mathbf{E}}_l$ *as above,*

$$\widehat{\mathbf{E}}_l^x[S_t^{(0)}] = (1 + O(p_B))\mathbf{E}^x[S_t^{(0)}] \quad \text{and} \quad \widehat{\mathbf{E}}_l^x[(S_t^{(0)})^2] \leqslant (1 + O(p_B))\mathbf{E}^x[(S_t^{(0)})^2].$$

Moreover, as in the proof of Lemma 5.13, one can show that

$$\widehat{\mathbf{Q}}_l^a[Z] = (1 + O(p_B))\mathbf{Q}^a[Z] \quad \text{and} \quad \widehat{\mathbf{Q}}_l^a[Z^2] \leqslant (1 + O(p_B))\mathbf{Q}^a[Z^2]. \tag{6.46}$$

We define two filtrations $(\mathscr{G}_l)_{l \geqslant 0}$ and $(\mathscr{H}_l)_{l \geqslant 0}$ by

$$\mathscr{G}_l = \mathscr{F}_{\mathscr{S}_\tau^{(l)}}, \quad \mathscr{H}_l = \mathscr{F}_{\mathscr{R}_\tau^{(l)}},$$

such that $\mathscr{G}_l \subset \mathscr{H}_l \subset \mathscr{G}_{l+1}$ for every l. Now define for measurable $I \subset [0,\tau]$,

$$Z_{\varnothing,I}^{(l)} = \sum_{(u,t) \in \mathscr{S}_\tau^{(l)} \cap (U \times I)} Z^{(u,t)}, \quad Y_{\varnothing,I}^{(l)} = \sum_{(u,t) \in \mathscr{S}_\tau^{(l)} \cap (U \times I)} Y^{(u,t)},$$

with $Z_\varnothing^{(l)} = Z_{\varnothing,[0,\tau]}^{(l)}$ and $Y_\varnothing^{(l)} = Y_{\varnothing,[0,\tau]}^{(l)}$. Furthermore, recall from Section 6.1 the definition of $Z_\varnothing^{(k;l)}$, $Z_\varnothing^{(l+)}$ and $Z_\varnothing$ and the corresponding quantities for Y.

Lemma 6.8. *We have for all* $0 \leqslant k \leqslant l$*,* $x \in (0,a)$ *and large* A *and* a*,*

$$\widehat{\mathbf{E}}_l[Z_\varnothing^{(k+1)} \,|\, \mathscr{G}_k] \leqslant (\pi + Cp_B)\mathbf{Q}^a[Z](\tfrac{\tau}{a^3} Z_\varnothing^{(k)} + CY_\varnothing^{(k)}). \tag{6.47}$$

In particular, for large A *and* a*,*

$$\widehat{\mathbf{E}}_l[Z_\varnothing^{(k+1)}] \leqslant CA(\tfrac{\tau}{a^3} Z_0 + CY_0)\Big(\pi^2 A(\tfrac{\tau}{a^3} + C_{6.48}\eta)\Big)^k. \tag{6.48}$$

In the case $k = 0$*, we also have for large* A *and* a*,*

$$\widehat{\mathbf{E}}_l[Z_\varnothing^{(1)}] \geqslant (\pi - Cp_B)\mathbf{Q}^a[Z](\tfrac{\tau}{a^3} Z_0 - CY_0). \tag{6.49}$$

Proof. We have for $0 \leqslant k \leqslant l$,

$$\widehat{\mathbf{E}}_l[Z_\varnothing^{(k+1)} \,|\, \mathscr{H}_k] = \widehat{\mathbf{E}}_l\Big[\sum_{(u,t) \in \mathscr{R}_\tau^{(k)}} Z^{(u,t)} \,\Big|\, \mathscr{H}_k\Big] = \widehat{\mathbf{Q}}_{l-k}^a[Z] R_\tau^{(k)}, \tag{6.50}$$

since conditioned on $\mathscr{H}_k$, the random variables $Z^{(u,t)}$ are iid under $\widehat{\mathbf{P}}_l$ of the same law as Z under $\widehat{\mathbf{Q}}_{l-k}^a$ and independent of $\mathscr{H}_k$, by the strong branching property. Now, we have

$$\widehat{\mathbf{E}}_l[R_\tau^{(k)} \,|\, \mathscr{G}_k] = \sum_{(u,t) \in \mathscr{S}_\tau^{(k)}} \mathbf{E}^{X_u(t)}[R_{\tau-t}^{(0)} \,|\, T > \tau - t],$$

such that by Lemma 5.6 and Corollary 6.7,

$$\widehat{\mathbf{E}}_l[R_\tau^{(k)} \,|\, \mathscr{G}_k] \leqslant (1 + O(p_B))\Big(\pi \frac{\tau}{a^3} Z_\varnothing^{(k)} + C_{5.36} Y_\varnothing^{(k)}\Big). \tag{6.51}$$

Equations (6.50) and (6.51) together with (6.46) give (6.47). Equation (6.48) follows easily from (6.47) by (6.16) and the fact that $Y_\varnothing^{(k)} \leqslant \eta Z_\varnothing^{(k)}$, $\widehat{\mathbf{P}}_l$-almost surely for $1 \leqslant k \leqslant l$. Now, in the case $k = 0$, we have $\mathscr{G}_0 = \mathscr{F}_0$ by definition. Denote the positions of the initial particles by $x_1, \ldots, x_n$. By Lemmas 5.6, 5.13 and (6.44),

$$\begin{aligned}
\widehat{\mathbf{E}}_l[R_\tau^{(0)}] = \sum_{i=1}^n \widehat{\mathbf{E}}_l^{x_i}[R_\tau^{(0)}] &\geqslant \sum_{i=1}^n \mathbf{E}^{x_i}[R_\tau^{(0)}] - \|p\|_\infty \mathbf{E}^{x_i}[(R_\tau^{(0)})^2] \\
&\geqslant \pi \frac{\tau}{a^3} Z_0 - C_{5.36} Y_0 - \|p\|_\infty C_{5.37} \left(\tfrac{\tau}{a^3} Z_0 + Y_0\right),
\end{aligned}$$

which yields (6.49). ☐

Lemma 6.9. *Suppose that* $t \leqslant \tau$. *Then for large* A *and* a, *we have*

$$\widehat{\mathbf{E}}[Z_t] = Z_0\left(1 + \pi \mathbf{Q}^a[Z]\tfrac{t}{a^3} + O\left(p_B + \left(A\tfrac{t}{a^3}\right)^2\right)\right) + O(AY_0), \quad \widehat{\mathbf{E}}[Y_t] \leqslant C\left(Y_0 + \eta A \tfrac{t}{a^3} Z_0\right).$$

If moreover $t \geqslant 2a^2$, *then*

$$\widehat{\mathbf{E}}[Y_t \mathbf{1}_{(R_{[t-a^2,t]}=0)}] \leqslant \frac{C}{a}(Z_0 + AY_0) \quad \textit{and} \quad \widehat{\mathbf{P}}(R_{[t-a^2,t]} \neq 0) \leqslant C\eta(Y_0 + Z_0).$$

Proof. First note that we have $\inf_{x\in[0,a]} h(x,0) \geqslant 1/2$ for large A, by (6.45). Furthermore, by the hypothesis $\tau \leqslant t_{6.21}$ and (6.3), we have $\pi^2 A(\frac{\tau}{a^3} + C\eta) \leqslant 1/2$ for large A and a. The first two inequalities now follow from Proposition 5.2, Corollary 6.7, Lemma 6.8, (6.46) and (6.3) by summing over k. In particular, if $t \geqslant 2a^2$, then for large A and a, $\widehat{\mathbf{E}}[Z_{t-a^2}] \leqslant C(Z_0 + AY_0)$ and $\widehat{\mathbf{E}}[Y_{t-a^2}] \leqslant C\eta(Z_0 + Y_0)$. Together with Proposition 5.2, Lemma 5.6 and Corollary 6.7, this proves the other two inequalities. $\square$

In order to estimate second moments, we will make use of the following extension to the Many-to-two lemma (Lemma 3.5). For $x, z \in (0,a)$ and $0 \leqslant t \leqslant \tau$, we define $\widehat{m_2}(x,t)$ to be the quantity $\widetilde{m_2}(x,t)$ from Section 5.5 corresponding to the penalisation from (6.43) and $\widehat{\mathbf{p}}_t^{(l)}(x,z)$ to be the density of tier l particles at position z and time t under the law $\widehat{\mathbf{P}}^x$, not counting the particles u with $t < \sigma_l(u)$. Then set $\widehat{\mathbf{p}}_t = \widehat{\mathbf{p}}_t^{(0+)}$.

Lemma 6.10. *Let* $w : [0,a] \to \mathbf{R}_+$ *be measurable and define* $S_t = \sum_{(u,t)\in\mathscr{N}_t} w(X_u(t))$. *Then*

$$\begin{aligned}\widehat{\mathbf{E}}^x[S_t^2] = \widehat{\mathbf{E}}^x\Big[\sum_{(u,s)\in\mathscr{N}_t} w(X_u(s))^2\Big] + \beta_0 \int_0^t \int_0^a \widehat{m_2}(z,s)\widehat{\mathbf{p}}_s(x,z)\widehat{\mathbf{E}}^{(z,s)}[S_t]^2 \,\mathrm{d}z\,\mathrm{d}s \\ + \widehat{\mathbf{E}}^x\Big[\sum_{(u,s)\in\mathscr{R}_t} \sum_{(v_1,s_1),(v_2,s_2)\in\mathscr{S}^{(u,s)},\, v_1\neq v_2} \widehat{\mathbf{E}}^{(X_{v_1}(s_1),s_1)}[S_t]\widehat{\mathbf{E}}^{(X_{v_2}(s_2),s_2)}[S_t]\Big]. \quad (6.52)\end{aligned}$$

Proof. For $(v_1,s_1),(v_2,s_2) \in \mathscr{N}_t$ we write $(v_1,s_1) \wedge (v_2,s_2)$ for their most recent common ancestor. Then define for $l \geqslant 1$,

$$\begin{aligned}\mathscr{A}_1^{(l)} = \{((v_1,s_1),(v_2,s_2)) \in \mathscr{N}_t^2 : v_1 \neq v_2 \text{ and if } (v_1,s_1)\wedge(v_2,s_2) = (v_0,s_0), \\ \text{then } \tau_{l-1}(v_0) \leqslant s_0 < \sigma_l(v_0)\},\end{aligned}$$

and for $l \geqslant 0$,

$$\begin{aligned}\mathscr{A}_2^{(l)} = \{((v_1,s_1),(v_2,s_2)) \in \mathscr{N}_t^2 : v_1 \neq v_2 \text{ and if } (v_1,s_1)\wedge(v_2,s_2) = (v_0,s_0), \\ \text{then } \sigma_l(v_0) \leqslant s_0 < \tau_l(v_0)\},\end{aligned}$$

We then have

$$\begin{aligned}\widehat{\mathbf{E}}^x\Big[\sum_{((v_1,s_1),(v_2,s_2))\in\mathscr{A}_1^{(l)}} w(X_{v_1}(s_1))w(X_{v_2}(s_2)) \,\Big|\, \mathscr{G}_l\Big] \\ = \sum_{(u,s)\in\mathscr{R}_t^{(l)}} \sum_{(v_1,s_1),(v_2,s_2)\in\mathscr{S}^{(u,s)},\, v_1\neq v_2} \widehat{\mathbf{E}}^{(X_{v_1}(s_1),s_1)}[S_t]\widehat{\mathbf{E}}^{(X_{v_2}(s_2),s_2)}[S_t], \quad (6.53)\end{aligned}$$

and by Lemma 3.5 (see also Remark 3.4),

$$\begin{aligned}\widehat{\mathbf{E}}^x\Big[\sum_{((v_1,s_1),(v_2,s_2))\in\mathscr{A}_2^{(l)}} w(X_{v_1}(s_1))w(X_{v_2}(s_2))\Big] \\ = \beta_0 \int_0^t \int_0^a m_2(z,s)\widehat{\mathbf{p}}_s^{(l)}(x,z)\widehat{\mathbf{E}}^{(z,s)}[S_t]^2 \,\mathrm{d}z\,\mathrm{d}s. \quad (6.54)\end{aligned}$$

Equations (6.53) and (6.54) and the tower property of conditional expectation yield the lemma. $\square$

Remark 6.11. An analogous result holds for R_t.

Lemma 6.12. *We have for every* $t \leqslant \tau$,

$$\widehat{\mathrm{Var}}(Z_t) \leqslant C\varepsilon e^A\Big(\frac{t}{a^3}Z_0 + Y_0\Big) \quad \text{and} \quad \widehat{\mathrm{Var}}(R_t) \leqslant C\varepsilon e^A\Big(\frac{t}{a^3}Z_0 + Y_0\Big).$$

Proof. By Lemma 6.9 and the hypothesis $t \leqslant \tau \leqslant t_{6.21}$, we have for every $x \in (0,a)$ and $s \leqslant t$,

$$\widehat{\mathbf{E}}^{(x,s)}[Z_t] \leqslant CA\Big(w_Z(x)\frac{t}{a^3} + w_Y(x)\Big) \leqslant CAe^{-\frac{3}{4}(a-x)}. \tag{6.55}$$

Now, as in the proof of (5.16), we have by (6.55) and Corollary 6.7,

$$\int_0^t \int_0^a \widehat{\mathbf{p}}_s^{(0)}(x,z)\widehat{\mathbf{E}}^{(z,s)}[Z_t]^2 \,\mathrm{d}z\,\mathrm{d}s \leqslant CA^2\Big(\frac{t}{a^3}w_Z(x) + w_Y(x)\Big), \tag{6.56}$$

which yields

$$\begin{aligned} &\int_0^t \int_0^a \widehat{m_2}(z,s)\widehat{\mathbf{p}}_s(x,z)\widehat{\mathbf{E}}^{(z,s)}[Z_t]^2 \,\mathrm{d}z\,\mathrm{d}s \\ &\qquad \leqslant C\widehat{\mathbf{E}}^x\Big[\sum_{(u,s_0)\in\mathscr{S}_t}\int_{s_0}^t\int_0^a \widehat{\mathbf{p}}_{s-s_0}^{(0)}(X_u(s_0),z)\widehat{\mathbf{E}}^{(z,s)}[Z_t]^2\,\mathrm{d}z\,\mathrm{d}s\Big] \\ &\qquad \leqslant CA^2\Big(\frac{t}{a^3}\widehat{\mathbf{E}}^x[Z_{\varnothing,[0,t]}] + \widehat{\mathbf{E}}^x[Y_{\varnothing,[0,t]}]\Big) \\ &\qquad \leqslant CA^2\Big(\frac{t}{a^3}w_Z(x) + w_Y(x)\Big), \end{aligned} \tag{6.57}$$

by Lemma 6.8 and the hypothesis $t \leqslant \tau \leqslant t_{6.21}$. Furthermore, by (6.55), (6.3) and Lemma 5.8, we have

$$\begin{aligned} &\widehat{\mathbf{E}}^x\Big[\sum_{(u,s)\in\mathscr{R}_t}\ \sum_{(v_1,s_1),(v_2,s_2)\in\mathscr{S}^{(u,s)},\,v_1\neq v_2} \widehat{\mathbf{E}}^{(X_{v_1}(s_1),s_1)}[Z_t]\widehat{\mathbf{E}}^{(X_{v_2}(s_2),s_2)}[Z_t]\Big] \\ &\qquad\qquad \leqslant C\mathbf{Q}^a[Z^2]\widehat{\mathbf{E}}^x[R_t] \leqslant C\varepsilon e^A\Big(\frac{t}{a^3}w_Z(x) + w_Y(x)\Big), \end{aligned} \tag{6.58}$$

by (6.17) and Lemmas 5.6 and Corollary 6.7. Lemma 6.10, together with (6.57) and (6.58) as well as Lemma 6.9 and the inequality $w_Z^2 \leqslant Cw_Y$ gives

$$\widehat{\mathrm{Var}}^x(Z_t) \leqslant \widehat{\mathbf{E}}^x[(Z_t)^2] \leqslant C\varepsilon e^A\Big(\frac{t}{a^3}w_Z(x) + w_Y(x)\Big). \tag{6.59}$$

Summing over the initial particles yields the inequality for $\widehat{\mathrm{Var}}(Z_t)$. The proof of the second inequality is analogous, relying on Lemma 5.6 and Remark 6.11. □

We finish the section by a corollary which will be useful in the next section.

Corollary 6.13. *Suppose that* $t \leqslant \tau$ *and* $x \in (0,a)$. *Then for large* A *and* a, *we have* $\widehat{\mathbf{E}}^x[Z_t] \leqslant CAe^{-(a-x)/2}$, $\widehat{\mathbf{E}}^x[Z_t^2] \leqslant C\varepsilon e^A e^{-(a-x)/2}$ *and* $\widehat{\mathbf{E}}^x[Y_t] \leqslant Ce^{-(a-x)/2}$.

Proof. Immediate from Lemmas 6.9 and 6.12 and (5.2). □

6.4 The fugitive and its family

We now describe the BBM conditioned to break out at a given time. Recall that $\mathscr{U}$ denotes the fugitive. For simplicity, we write τ_l and σ_l for $\tau_l(\mathscr{U})$ and $\sigma_l(\mathscr{U})$, respectively, $l \in \mathbf{N}$. On the event $T = T^{(l)}$, we define $\mathscr{U}_j$ to be the ancestor of $\mathscr{U}$ alive at the time σ_j, $j = 0, \dots, l$. By the strong branching property, we have the following decomposition:

Lemma 6.14. *Let $k \in \mathbf{N}$, $l \in \mathbf{N} \cup \{\infty\}$ with $l \geqslant k$ and $t \geqslant 0$. Conditioned on $\mathscr{F}_0$, $T^{(0;l)} = T^{(k)} = t$, τ_0 and $\mathscr{U}_0$, the BBM admits the following recursive decomposition:*

1. *The initial particles $u \neq \mathscr{U}_0$ spawn independent BBM conditioned on $T^{(0;l)} > t$,*
2. *independently, the particle $\mathscr{U}_0$ spawns BBM conditioned on a particle (call it $\mathscr{U}_0'$) hitting a for the first time at the time τ_0, all the children of which born before τ_0 being conditioned on $T^{(0;l)} > t$.*
   - *(a) If $k = 0$, this particle $\mathscr{U}_0'$ spawns BBM conditioned on the event B of a breakout.*
   - *(b) If $k > 0$, it spawns BBM starting at the space-time point (a, τ_0) conditioned on $T^{(0;l)} = T^{(k)} = t$. In particular, if we write $\mathscr{S} = \mathscr{S}^{(\mathscr{U}_0',\tau_0)}$, then conditioned on $\mathscr{F}_{\mathscr{S}}$, the particles in $\mathscr{S}$ spawn BBM starting from the collection of space-time points $\mathscr{S}$, conditioned on $T^{(0;l-1)} = T^{(k-1)} = t$.*

Note that in the case 2b above, the subtree spawned by $(\mathscr{U}_0', \tau_0)$ follows the law $\mathbf{Q}$ *conditioned* on $T^{(0;l)} = T^{(k)} = t$, hence the law of $\mathscr{S}^{(\mathscr{U}_0',\tau_0)}$ is not the same as under $\mathbf{Q}$. In particular, $\mathbf{E}[Z^{(\mathscr{U}_0',\tau_0)}] \not\leqslant CA$. Indeed, conditioning on one of its descendants breaking out at a later time corresponds to a kind of size-bias on the number of particles. However, it is still true that $Z^{(\mathscr{U}_0',\tau_0)} < \varepsilon e^A$, by the definition of a breakout.

Lemma 6.14 gives a decomposition of the BBM conditioned on $T^{(k)}$ into $k+1$ pieces. In order to describe what happens in a single piece, define $\mathscr{F}_{\mathscr{U}}$ to be the σ-field generated by the family of events $(\{T^{(0;l)} = T^{(k)}\})_{k \in \mathbf{N}}$ and by $\mathscr{U}$, $(X_{\mathscr{U}}(s))_{\sigma_j \leqslant s < \tau_j}$, $(d_u, k_u)_{u \preceq \mathscr{U}}$ and $\mathscr{S}^{(\mathscr{U},\tau_j)}$, for $j = 0, \dots, k$, on the event $\{T^{(0;l)} = T^{(k)}\}$. Note that in particular, on the event $\{T^{(0;l)} = T^{(k)}\}$, τ_j, $(\mathscr{U}_j, \sigma_j)$ are $\mathscr{F}_{\mathscr{U}}$-measurable, $j = 0, \dots, k$. It is plain that conditioned on $\mathscr{F}_{\mathscr{U}}$, the subtrees spawned by the children of the ancestors of $\mathscr{U}$ during the intervals $[\sigma_j, \tau_j)$ are independent BBM conditioned not to break out before t.

It remains to describe the trajectory of the fugitive and its reproduction. We could do by a decomposition of the process at the first branching, but it is faster to use the Many-to-one lemma instead, which is the method of proof of the following lemma, which we state for general penalisations $p(s)$. This result is essentially [24, Theorem 1].

Lemma 6.15. *Let $t \geqslant 0$, $x \in (0, a)$ and $p : \mathbf{R}_+ \to [0,1]$ be measurable with $p(s) = 0$ for s large enough. Denote by $\widetilde{\mathbf{P}}$ the law associated to $p(s)$ as in Section 5.5. Recall the definition of $e(x,t)$ from (5.40). Then, given a family of $\mathscr{F}_{\mathscr{R}_t^{(0)}}$-measurable non-negative random variables $(Y_u)_{u \in U}$, we have*

$$\widetilde{\mathbf{E}}^x\Big[Y_u \,\Big|\, \exists u \in U : (u,t) \in \mathscr{R}_t^{(0)}\Big] = \frac{\widetilde{\mathbf{E}}^{*,x}\Big[Y_{\xi_t} e^{-\int_0^t e(\xi_s,s)\,\mathrm{d}s} \,\Big|\, H_0(\xi) > H_a(\xi) = t\Big]}{\widetilde{\mathbf{E}}^{*,x}\Big[e^{-\int_0^t e(\xi_s,s)\,\mathrm{d}s} \,\Big|\, H_0(\xi) > H_a(\xi) = t\Big]},$$

where under $\widetilde{\mathbf{P}}^{*,x}$, the spine moves according to standard Brownian motion and spawns particles with rate $\widetilde{m_1}(x,t)\beta_0 Q(x,s)$ according to the reproduction law $((\widetilde{m_1}(x,t))^{-1} k \widetilde{q}(x,t,k))_{k \geqslant 0}$, which start independent $\widetilde{\mathbf{P}}$-BBM. In particular, conditioned on $(X_u(s))_{0 \leqslant s \leqslant t}$ and $(d_v, k_v)_{v \preceq u}$, the children of the ancestors of u follow independent $\widetilde{\mathbf{P}}$-BBM.

Proof. We have

$$\widetilde{\mathbf{E}}^x\Big[Y_u \,\Big|\, \exists u \in U : (u,t) \in \mathscr{R}_t^{(0)}\Big] = \frac{\widetilde{\mathbf{E}}^x\Big[\sum_{(u,s) \in \mathscr{R}_t^{(0)}} \mathbf{1}_{(H_a(X_u) \in \mathrm{d}t)} Y_u\Big]}{\widetilde{\mathbf{E}}^x\Big[\sum_{(u,s) \in \mathscr{R}_t^{(0)}} \mathbf{1}_{(H_a(X_u) \in \mathrm{d}t)}\Big]}, \tag{6.60}$$

and by Lemma 3.1,

$$\widetilde{\mathbf{E}}^x\Big[\sum_{(u,s)\in\mathscr{R}_t^{(0)}}\mathbf{1}_{(H_a(X_u)\in\mathrm{d}t)}Y_u\Big]=\widetilde{\mathbf{E}}^{*,x}_{-\mu}\Big[Y_{\xi_t}e^{\int_0^t\widetilde{m}(\xi_s,s)\beta_0Q(\xi_s,s)\,\mathrm{d}s}\mathbf{1}_{(H_0(\xi)>H_a(\xi)\in\mathrm{d}t)}\Big].\tag{6.61}$$

According to the description of the conditioned process in Section 5.5 and the description of the spine in Section 3.3, the particles on the spine follow the Doob transform of Brownian motion with drift $-\mu$ by the harmonic function $h(x,t)$ and spawn particles with rate $\widetilde{m_1}(x,t)\beta_0Q(x,s)$ according to the reproduction law $((\widetilde{m_1}(x,t))^{-1}k\widetilde{q}(x,t,k))_{k\geqslant 0}$. With Girsanov's transform, (6.61) yields,

$$\widetilde{\mathbf{E}}^x\Big[\sum_{(u,s)\in\mathscr{R}_t^{(0)}}\mathbf{1}_{(H_a(X_u)\in\mathrm{d}t)}Y_u\Big]=e^{-\mu(a-x)}\frac{h(a,t)}{h(x,0)}\widetilde{\mathbf{E}}^{*,x}\Big[Y_{\xi_t}e^{-\int_0^t e(\xi_s,s)\,\mathrm{d}s}\mathbf{1}_{(H_0(\xi)>H_a(\xi)\in\mathrm{d}t)}\Big].\tag{6.62}$$

Equations (6.60) and (6.62) yield the first statement. The second statement follows by taking appropriate test functionals $(Y_u)_{u\in U}$. ☐

Corollary 6.16. *In addition to the assumptions in Lemma 6.15, suppose $t\in[0,a^3]$ and $\|p\|_\infty\leqslant c_p$, where c_p is some universal constant implicitly defined below. Then,*

$$\widetilde{\mathbf{E}}^x\Big[Y_u\,\Big|\,\exists u\in U:(u,t)\in\mathscr{R}_t^{(0)}\Big]\leqslant C\widetilde{\mathbf{E}}^{*,x}\Big[Y_{\xi_t}\,\Big|\,H_0(\xi)>H_a(\xi)\in\mathrm{d}t\Big].\tag{6.63}$$

Proof. By Lemma 2.2, the inequality $e^{-x}\geqslant 1-x$ and the hypothesis on t, we have

$$\widetilde{\mathbf{E}}^{*,x}\Big[e^{-\int_0^t e(\xi_s,s)\,\mathrm{d}s}\,\Big|\,H_0(\xi)>H_a(\xi)=t\Big]=W^{x,t,a}_{\mathrm{taboo}}[e^{-\int_0^t e(X_s,s)\,\mathrm{d}s}]\geqslant(1-C\|p\|_\infty).\tag{6.64}$$

The statement now follows from (6.64) and Lemma 6.15. ☐

We come back to the BBM conditioned to break out at a given time and set up the important definitions. Recall that $\mathscr{U}$ denotes the fugitive. We will denote by a *bar* the quantities referring to the particles spawned between the times $\sigma_l(\mathscr{U})$ and $\tau_l(\mathscr{U})$ for some $l\geqslant 0$, and by a *check* those referring to the particles spawned between the times $\tau_l(\mathscr{U})$ and $\sigma_{l+1}(\mathscr{U})$. Formally, we set

$$\overline{\mathscr{N}_t}=\{u\in\mathscr{N}_t:(u,t)\wedge(\mathscr{U},T)\in U\times\bigcup_{l\geqslant 0}[\sigma_l(\mathscr{U}),\tau_l(\mathscr{U}))\},$$

$$\bar{\mathscr{R}}_t=\{(u,s)\in\bigcup_{l\geqslant 0}\mathscr{R}_t^{(l)}:(u,s)\wedge(\mathscr{U},T)\in U\times\bigcup_{l\geqslant 0}[\sigma_l(\mathscr{U}),\tau_l(\mathscr{U})))\}$$

and

$$\widecheck{\mathscr{N}_t}=\{u\in\mathscr{N}_t:(u,t)\wedge(\mathscr{U},T)\in U\times\bigcup_{l\geqslant 1}[\tau_{l-1}(\mathscr{U}),\sigma_l(\mathscr{U}))\}.$$

We then define

$$\bar{Z}_t=\sum_{u\in\overline{\mathscr{N}_t}}w_Z(X_u(t)),\quad \bar{Y}_t=\sum_{u\in\overline{\mathscr{N}_t}}w_Y(X_u(t)),\quad \bar{R}_t=\#\bar{\mathscr{R}}_t,$$

and

$$\check{Z}_t=\sum_{u\in\overline{\mathscr{N}_t}}w_Z(X_u(t)),\quad \check{Y}_t=\sum_{u\in\overline{\mathscr{N}_t}}w_Y(X_u(t)).$$

Note that on the event $T=T^{(0)}$, we have $\widecheck{\mathscr{N}}(T)=\varnothing$ by definition.

By Corollary 6.13, the following $\mathscr{F}_{\mathscr{U}}$-measurable functional will be of use in the study of the bar-quantities.

$$\mathscr{E}_{\mathscr{U}}=\sum_{i=0}^{l}\sum_{u\leq\mathscr{U}}\mathbf{1}_{(\sigma_i\leqslant d_u<\tau_i)}(k_u-1)e^{-(a-X_u(d_u))/2},\qquad\text{on }\{T=T^{(l)}\}.\tag{6.65}$$

where we recall that d_u and k_u are respectively the time of death and the number of children of the individual u.

Lemma 6.17. *For large A and a, we have for every $l \in \mathbf{N}$,*

$$\mathbf{E}[\mathscr{E}_{\mathscr{U}} \,|\, T = T^{(l)} \leqslant t_{6.21}] \leqslant C(l+1).$$

Proof. By Lemma 6.14 and Corollary 6.16 (which can be applied because of (6.44)), we have for every $l \in \mathbf{N}$,

$$\mathbf{E}[\mathscr{E}_{\mathscr{U}} \,|\, T = T^{(l)} \leqslant t_{6.21}] \leqslant C \sum_{i=0}^{l} \mathbf{E}\Big[W_i\Big[\int_0^{\tau_i - \sigma_i} e^{-(a-B_t)/2}\,\mathrm{d}t\Big] \Big| T = T^{(l)} \leqslant t_{6.21}\Big],$$

where W_i is the law of a Brownian bridge of length $\tau_i - \sigma_i$ from $X_{\mathscr{U}}(\sigma_i)$ to $X_{\mathscr{U}}(\tau_i)$. The statement now follows readily from Lemma 2.2. ☐

We can now study the probability that the fugitive stems from a given tier.

Lemma 6.18. *Suppose that $C_1 e^A \leqslant Z_0 \leqslant C_2 e^A$ and $Y_0 \leqslant \eta Z_0$. Then for large A and a, we have for $l \in \{1,2\}$,*

$$\mathbf{P}(T^{(l+)} < T^{(0;l-1)}) \leqslant C(\varepsilon A)^l,$$

Proof. By (6.2), (6.14) and (6.46), we have for large A and a,

$$\mathbf{P}(T^{(0)} > t_{6.21}) \leqslant \exp(-CA/\varepsilon) \leqslant \exp(-C/\sqrt{\varepsilon}). \tag{6.66}$$

Now, for the rest of the proof, let $t \leqslant t_{6.21}$ and let $\nu = \sum_{i=1}^n \delta_{x_i}$. We have by the decomposition (6.20) of the process conditioned on $T^{(0;l-1)} = t$,

$$\mathbf{P}^{\nu}(T^{(l+)} > t \,|\, T^{(0;l-1)} = t) = \sum_{i=1}^{n} p_i \widehat{\mathbf{P}}^{\nu - \delta_{x_i}}(T^{(l+)} > t \,|\, T^{(0;l-1)} > t) \times \mathbf{P}^{x_i}(T^{(l+)} > t \,|\, T^{(0;l-1)} = t). \tag{6.67}$$

Define $c_t = \pi^2 A(t/a^3 + C_{6.48}\eta)$, which is less than $1/2$ for large A and a by the hypothesis on t. By Lemma 6.8 and the hypothesis, we have for every $j \leqslant l$,

$$\widehat{\mathbf{E}}_{l-j-1}[Z_{\varnothing}^{(l-j)}] \leqslant C Z_0 c_t^{l-j}, \tag{6.68}$$

and moreover for every $(x,s) \in (0,a) \times [0,t]$,

$$\widehat{\mathbf{E}}_{l-j-1}^{(x,s)}[Z_{\varnothing}^{(l-j)}] \leqslant C A e^{-(a-x)/2} c_t^{l-j-1}, \tag{6.69}$$

For every $k \leqslant l-1$, (6.68) applied to the particles in $\mathscr{S}_t^{(j)}$, $j = 1, \dots, k$ yields

$$\mathbf{E}[\check{Z}_{\varnothing}^{(l)} \,|\, T^{(k)} = T^{(0;l-1)} = t] \leqslant C\varepsilon e^A \sum_{j=1}^{l} c_t^{l-j} \leqslant C\varepsilon e^A c_t^{l-k}. \tag{6.70}$$

Moreover, by (6.69), Corollary 6.13 and Lemma 6.17, we have

$$\mathbf{E}[\bar{Z}_{\varnothing}^{(l)} \,|\, T^{(k)} = T^{(0;l-1)} = t] \leqslant C A c_t^{l-k}. \tag{6.71}$$

In total, we get by (6.68), (6.69) and (6.70), the hypothesis on Z_0 and (6.2),

$$\mathbf{E}[\widehat{Z}_{\varnothing}^{(l)} + \check{Z}_{\varnothing}^{(l)} + \bar{Z}_{\varnothing}^{(l)} \,|\, T^{(k)} = T^{(0;l-1)} = t] \leqslant C e^A \Big(c_t^l + \varepsilon c_t^{l-k}\Big). \tag{6.72}$$

Jensen's inequality and Proposition 6.2 together with (6.72) and the inequality $1 - e^{-x} \leqslant x$ give

$$\mathbf{P}(T^{(l+)} < t \,|\, T^{(k)} = T^{(0;l-1)} = t) \leqslant C c_t(\varepsilon^{-1} c_t^l + c_t^{l-k}). \tag{6.73}$$

Summing (6.73) over k, the law of total expectation gives

$$\mathbf{P}(T^{(l+)} < t \,|\, T^{(0;l-1)} = t) \leqslant C c_t(\varepsilon^{-1} c_t^l + c_t). \tag{6.74}$$

The lemma now follows by integrating (6.74) over t from 0 to $t_{6.21}$ and using Lemma 6.5 and (6.66). ☐

Remark 6.19. One may wonder whether one can simply calculate $\mathbf{P}(T^{(l)} \in \mathrm{d}t \,|\, T^{(0;l-1)} > t)$ for every $l \geqslant 1$ and $t \leqslant t_{6.21}$, using only the tools from Section 6.3. This would require fine estimates on the density of the point process formed by the particles from tier $l-1$ hitting a just before t. These estimates can be most easily obtained if one stops descendants of the particles hitting a at a (large) fixed time ζ instead of the line from Lemma 5.8, with which the results in this paper would hold as well. However, in order not to lose generality, we stick to Lemma 6.18, which is enough for our purposes.

7 The B-BBM

We will now define properly the BBM with the moving barrier, also called the B-BBM (the "B" stands for "barrier"), which will be used in the subsequent sections to approximate the N-BBM. We will still use all the definitions from Section 6.1, with one notational change: Recall that by (6.20), we can decompose the process into two parts; the first part consisting of the particles spawned by the ancestors of the fugitive and the second part consisting of the remaining particles. As in Section 6.4, the quantities which refer to the particles of the first part will be denoted by a bar (e.g. $\bar{Z}$) or check (e.g. $\check{Z}$). The quantities of the second part will be denoted with a hat in this section (e.g. $\widehat{Z}$), in reference to the law $\widehat{\mathbf{P}}$ from Section 6.3. Moreover, the quantities related to the descendants of the fugitive will be denoted by the superscript "fug" (e.g. Z^{fug}).

7.1 Definition of the model

Suppose that we are given a family $(f_x)_{x\geqslant 0}$ of non-decreasing functions $f_x \in \mathscr{C}^2(\mathbf{R},\mathbf{R}_+)$, such that for each $x \geqslant 0$, $f_x(t) = 0$ for $t \leqslant 0$, $f_x(+\infty) = x$ and for each $\delta > 0$ small enough there exist $x_0 = x_0(\delta)$, $t_0 = t_0(\delta)$, such that

1. $x_0(\delta) \to \infty$ as $\delta \to 0$,
2. $\|f_x\| \leqslant \delta^{-1}$ for all $x \in [0, x_0]$,
3. $f_x(t) \geqslant (1-\delta)x$ for all $t \geqslant t_0$ and
4. $t_0 \leqslant \delta^{-1}$.

where $\|f\|$ is defined in (5.4). A family of functions with the above properties exists, for example the following, which we will choose in Sections 8 and 9.

$$f_x(t) = \log\left(1 + (e^x - 1)\overline{\theta}(t)\right). \tag{7.1}$$

For every A and a, let $\nu_0 = \nu_0^{A,a}$ be a (possibly random) finite counting measure. We now define the B-BBM with initial configuration of particles ν_0. Starting from BBM with constant drift $-\mu$ with initial configuration ν_0, we define for each $n \in \mathbf{N}$ a stopping time Θ_n and for each $n \in \mathbf{N}^*$ the barrier process $(X_t^{[n]})_{t\in[\Theta_{n-1},\Theta_n]}$ as follows:

1. We set $\Theta_0 = 0$.
2. Denote by $T = T_1$ the time of the first breakout of the BBM absorbed at 0 and by $\mathscr{U}$ the fugitive, as in Section 6.1. We set $X_t^{[1]} = 0$ for $t \in [0, T]$.
3. Define $T^- = (T - e^A a^2) \wedge 0$ and
$$\Delta = \log\left(e^{-A}(\widehat{Z}_{T^-} + \check{Z}_{\varnothing\vee T^-} + Z^{(\mathscr{U},T)})\right) \vee 0. \tag{7.2}$$
where $\check{Z}_{\varnothing\vee T^-} = \check{Z}^{(1)}_{T^-}$ if $T^- \geqslant \tau_0(\mathscr{U})$ and $= \check{Z}^{(1)}_{\varnothing}$ otherwise. The above quantities are defined in Section 6.4, after Corollary 6.16.
4. Define $T^+ = T_1^+ = T + \tau_{\max}^{(\mathscr{U},T)}$ and $\Theta_1 = (T + e^A a^2) \vee T^+$. Note that T^+ and therefore also Θ_1 is a stopping time for the BBM. Now define
$$X_t^{[1]} = f_\Delta\Big(\frac{t - T^+}{a^2}\Big), t \in [T^+, \Theta_1].$$

We then give to the particles an additional drift $-(\mathrm{d}/\mathrm{d}t)X_t^{[1]}$ for $t \in [T^+, \Theta_1]$, in the meaning of Section 5.1.

5. We have now defined T_1, T_1^+, Θ_1 and $X^{[1]}$. We further define ν_1 to be the measure formed by the particles at time Θ_1, which have never hit 0. To define T_2, T_2^+, Θ_2 and $X^{[2]}$, we repeat the above steps with the process formed by the BBM started from those particles, with the definitions changed such that the barrier process starts at $X^{[2]}_{\Theta_1} = X^{[1]}_{\Theta_1}$, time starts at Θ_1 etc.
6. We now construct the barrier process $X_t^{[\infty]}$ from the pieces by $X_t^{[\infty]} = X_t^{[n]}$, if $t \in [\Theta_{n-1}, \Theta_n]$.

Define the event $G_0 = \{\operatorname{supp}\nu_0 \subset (0,a),\ |Z_0 - e^A| \leqslant \varepsilon^{3/2}e^A,\ Y_0 \leqslant \eta\}$. Recall the definition of the phrase "As A and a go to infinity" from Section 6.1. Define the predicate

(HB) (The law of) ν_0 is such that $\mathbf{P}(G_0) \to 1$ as A and a go to infinity.

(HB$_0$) ν_0 is deterministic and such that G_0 is verified.

Furthermore, if for some $n > 0$, $(t_j^{A,a})_{j=1}^n \in \mathbf{R}_+^n$ for all A and a, then define the predicate

(Ht) There exists $0 \leqslant t_1 < \cdots < t_n$, such that for all $j \in \{1,\dots,n\}$, $t_j^{A,a} \to t_j$ as A and a go to infinity, with $t_1^{A,a} \equiv 0$ if $t_1 = 0$.

Theorem 7.1. *Suppose (HB). Let $n \geqslant 1$ and $(t_j^{A,a})_{j=1}^n \in \mathbf{R}_+^n$ such that (Ht) is verified. Define the process*

$$(X_t)_{t\geqslant 0} = (X^{[\infty]}_{a^3t} - \pi^2 At)_{t\geqslant 0}.$$

Then, as A and a go to infinity, the law of the vector $(X_{t_j^{A,a}})_{j=1}^n$ converges to the law of $(L_{t_j})_{j=1}^n$, where $(L_t)_{t\geqslant 0}$ is the Lévy process from Theorem 1.1.

A stronger convergence than convergence in the sense of finite-dimensional distributions is convergence in law with respect to Skorokhod's (J_1-)topology (see [28, Chapter 3]). Obviously, the convergence in Theorem 7.1 does not hold in this stronger sense, because the barrier is continuous but the Lévy process is not and the set of continuous functions is closed in Skorokhod's J_1-topology[5]. However, if we create artificial jumps, we can rectify this:

Theorem 7.2. *Suppose (HB). Define $J_t = X^{[\infty]}_{\Theta_n}$, if $t \in [\Theta_n, \Theta_{n+1})$, for $n \in \mathbf{N}$. Then as A and a go to infinity, the process $(X'_t)_{t\geqslant 0} = (J_{a^3t} - \pi^2 At)_{t\geqslant 0}$ converges in law with respect to Skorokhod's topology to the Lévy process defined in the statement of Theorem 7.1.*

For each $n \geqslant 1$, we define the event G_n to be the intersection of G_{n-1} with the following events (see Section 6.1 for the definition of $\mathscr{N}_t$):

- $\operatorname{supp}\nu_n \subset (0,a)$,
- $\mathscr{N}_{\Theta_n} \subset U \times \{\Theta_n\}$ and $\Theta_n > T_n^+$ (for $n > 0$),
- $|e^{-A}Z_{\Theta_n} - 1| \leqslant \varepsilon^{3/2}$ and $Y_{\Theta_n} \leqslant \eta$.

Note that $G_n \in \mathscr{F}_{\Theta_n}$ for every $n \geqslant 0$.

The core of the proof of Theorems 7.1 and 7.2 will be the following proposition:

Proposition 7.3. *Fix $\lambda \in \mathbf{R}$ and define $\gamma_0 = (\pi p_B e^A)^{-1}$. There exists a numerical constant $b > 0$, such that for large A and a, we have $\mathbf{P}(G_n^c \,|\, \mathscr{F}_0)\mathbf{1}_{G_0} \leqslant n\varepsilon^{1+b}$ for every $n \geqslant 0$ and*

$$\log \mathbf{E}\Big[e^{i\lambda X^{[\infty]}_{\Theta_n}}\,\Big|\,\mathscr{F}_0\Big]\mathbf{1}_{G_0} = \Big(n\gamma_0(\kappa(\lambda) + i\lambda\pi^2 A + o(1) + O(\varepsilon^b))\Big)\mathbf{1}_{G_0}. \tag{7.3}$$

where $\kappa(\lambda)$ is the cumulant from (1.1) and where $o(1)$ and $O(\varepsilon^b)$ may depend on λ.

---

[5] One could prove convergence in the M_1-topology (cf. [58, Section 3.3]), but the use of this less standard topology would lead us away from our real goal: the proof of Theorem 1.1.

7.2 Proof of Proposition 7.3

First note that conditioning on $\mathscr{F}_0$, we can and will assume without loss of generality that the hypothesis (HB$_0$) holds and will only state it again in the lemmas. In the same vein, *we will always assume that A and a are large and will omit stating this fact, except again in the statements of the lemmas.*

Furthermore, it is only necessary to treat the case $n = 1$, since the rest can be obtained by induction: Suppose that we have shown the result for $n = 1$. Since the process starts afresh at the stopping time Θ_n, we then have $\mathbf{P}(G_n \backslash G_{n+1} \,|\, \mathscr{F}_{\Theta_n}) \leqslant \varepsilon^{1+b}\mathbf{1}_{G_n}$ for every $n \geqslant 0$, which implies the statement on the probability of G_n. Similarly, if we set $\kappa'(\lambda) = \gamma_0\left(\kappa(\lambda) + i\lambda\pi^2 A + o(1)\right)$, where $o(1)$ is the term from (7.3), then

$$\begin{aligned}\mathbf{E}\Big[e^{i\lambda(X^{[\infty]}_{\Theta_{n+1}} - X^{[\infty]}_{\Theta_n})}\mathbf{1}_{G_{n+1}} \,\big|\, \mathscr{F}_{\Theta_n}\Big] &= \Big(e^{\kappa'(\lambda)+O(\varepsilon^b)} + O\left(\mathbf{P}(G_n \backslash G_{n+1} \,|\, \mathscr{F}_{\Theta_n})\right)\Big)\mathbf{1}_{G_n} \\ &= \Big(1 + O(\varepsilon^{1+b})\Big)\, e^{\kappa'(\lambda)+O(\varepsilon^b)}\mathbf{1}_{G_n},\end{aligned}$$

because $\gamma_0 = \varepsilon(1 + o(1))$ by (6.14). It follows that

$$\mathbf{E}\Big[e^{i\lambda X^{[\infty]}_{\Theta_{n+1}}}\mathbf{1}_{G_{n+1}}\Big] = e^{\kappa'(\lambda)+O(\varepsilon^b)}\mathbf{E}\Big[e^{i\lambda X^{[\infty]}_{\Theta_n}}\Big(1 + O(\varepsilon^{1+b})\Big)\mathbf{1}_{G_n}\Big] = e^{\kappa'(\lambda)+O(\varepsilon^b)}\mathbf{E}\Big[e^{i\lambda X^{[\infty]}_{\Theta_n}}\mathbf{1}_{G_n}\Big]$$

By induction, and the fact that $\mathbf{P}(G^c_{n+1}) \leqslant n\varepsilon^{1+b} \leqslant e^{O(\varepsilon^{1+b})}$, we then obtain (7.3).

Throughout the proof, we will work on several "good sets", which we all define here for easy reference, since they will be reused in the following sections. For this reason, some of them (for example $\widehat{G}$) are actually more restrictive than what would be necessary for the proof of Proposition 7.3.

- $G_{\mathscr{U}} = \{e^A a^2 \leqslant T \leqslant \sqrt{\varepsilon}a^3,\ T = T^{(0;1)},\ \mathscr{E}_{\mathscr{U}} \leqslant e^{A/3}\}$.
- $G_{\mathrm{fug}} = \{Z^{(\mathscr{U},T)} \leqslant e^A/\varepsilon,\ Y^{(\mathscr{U},T)} \leqslant Z^{(\mathscr{U},T)}/y\} \cap G^{(\mathscr{U},T)}$ (the event $G^{(\mathscr{U},T)}$ was defined in (6.10)).
- $\widehat{G} = \{|\widehat{Z}_{T-} - e^A| \leqslant \varepsilon^{1/4}e^A,\ \widehat{Y}_{T-} \leqslant a^{-1}e^{3A/2}\}$.
- $\check{G} = \{\check{Z}_{\varnothing\vee T-} \leqslant \varepsilon^{1/4}e^A,\ \check{Y}_{\varnothing\vee T-} \leqslant e^{-A/2}\}$ (we define $\check{Y}_{\varnothing\vee T-}$ analogously to $\check{Z}_{\varnothing\vee T-}$).
- $G_\Delta = G_{\mathscr{U}} \cap G_{\mathrm{fug}} \cap \widehat{G} \cap \check{G}$.
- G_{nbab} ("nbab" stands for "no breakout after breakout") is the event that no bar-particle breaks out between T and $\Theta_1 + e^A a^2$ and that no hat-, fug- or check-particle (fug-particles = descendants of $\mathscr{S}^{(\mathscr{U},T)}$) hits a between T^- and $\Theta_1 + e^A a^2$.

Recall the definition of $\mathscr{F}_{\mathscr{U}}$ from Section 6.4 and set $\mathscr{F}_\Delta := \mathscr{F}_{\mathscr{U}} \vee \widehat{\mathscr{F}}_{T-} \vee \check{\mathscr{F}}_{T-}$, such that $G_\Delta \in \mathscr{F}_\Delta$ and the random variable Δ is measurable with respect to $\mathscr{F}_\Delta$. Define further $Z_\Delta = \widehat{Z}_{T-} + Z^{(\mathscr{U},T)} + \check{Z}_{\varnothing\vee T-}$ and Y_Δ analogously. Finally, define $\mathbf{P}_{\mathrm{nbab}} = \mathbf{P}(\cdot \,|\, G_{\mathrm{nbab}})$.

The probability of G_Δ.

Lemma 7.4. *Suppose (HB$_0$). For large A and a, we have $\mathbf{P}(G_\Delta) \geqslant 1 - C\varepsilon^{5/4}$.*

Proof. By Proposition 6.2 and Lemma 6.18, we have

$$\mathbf{P}(e^A a^2 \leqslant T \leqslant \sqrt{\varepsilon}a^3,\ T = T^{(0;1)}) \geqslant 1 - CA^2\varepsilon^2. \tag{7.4}$$

Since $\sqrt{\varepsilon} \leqslant t_{6.21}$ for large A by (6.1), Equation (7.4) and Markov's inequality applied to Lemma 6.17 yield

$$\mathbf{P}(G_{\mathscr{U}}) \geqslant 1 - CA^2\varepsilon^2. \tag{7.5}$$

Furthermore, by (6.13) and the last statement of Lemma 5.8, we have,

$$\mathbf{P}(G^c_{\mathrm{fug}}) \leqslant p_B^{-1}\left(\mathbf{P}^a\left(\pi W > e^A/\varepsilon - 2\eta\right) + \eta\right) \leqslant C\varepsilon^2, \tag{7.6}$$

by (4.5), (6.2), (6.3) and (6.14).

In order to estimate the probability of $\widehat{G}$, we will calculate first- and second moments of the quantities in the definition of $\widehat{G}$. These estimates will be needed again later on, when we calculate the Fourier transform of the variable Δ.

By Lemma 6.9 and the hypotheses, together with (6.14), (6.2) and (6.3),

$$\mathbf{E}[e^{-A}\widehat{Z}_{T-}\,|\,\mathscr{F}_{\mathscr{U}}]\mathbf{1}_{G_{\mathscr{U}}} = 1+\pi\mathbf{Q}^a[Z]\tfrac{T}{a^3}+O\left(\varepsilon^{3/2}+\left(A\tfrac{T}{a^3}\right)^2\right), \tag{7.7}$$

which together with Lemma 6.12 gives

$$\begin{aligned}\mathbf{E}[(e^{-A}\widehat{Z}_{T-}-1)^2\,|\,\mathscr{F}_{\mathscr{U}}]\mathbf{1}_{G_{\mathscr{U}}} &\leqslant \mathrm{Var}(e^{-A}\widehat{Z}_{T-}\,|\,\mathscr{F}_{\mathscr{U}})\mathbf{1}_{G_{\mathscr{U}}}+C\left((A\tfrac{T}{a^3})^2+\varepsilon^3\right)\\ &\leqslant C\left((A\tfrac{T}{a^3})^2+\varepsilon\tfrac{T}{a^3}+\varepsilon^3\right),\end{aligned} \tag{7.8}$$

by the hypotheses on the initial configuration. Lemma 6.5 and (7.7), together with (6.1), (6.4) and (6.15), now give

$$\mathbf{E}[e^{-A}\widehat{Z}_{T-}\,|\,G_{\mathscr{U}}] = 1+\pi^2\gamma_0(A+\log\varepsilon+c_{6.15}+o(1)). \tag{7.9}$$

Similarly, (7.7) and (7.8) and Lemma 6.5 give

$$\mathbf{E}[(e^{-A}\widehat{Z}_{T-}-1)^2\,|\,G_{\mathscr{U}}] = O(A^2\varepsilon^2). \tag{7.10}$$

Furthermore, by Lemma 6.9 and Markov's inequality, together with (6.3) and the hypotheses, we have

$$\mathbf{P}[\widehat{Y}_{T-} > e^{3A/2}/a\,|\,G_{\mathscr{U}}] \leqslant Ce^{-A/2}. \tag{7.11}$$

By the same arguments, and since $\widehat{Z}^{(1)}_{\varnothing} \leqslant \varepsilon e^A\mathbf{1}_{(T=T^{(1)})}$ by the definition of a breakout, we have

$$\mathbf{E}[e^{-A}\check{Z}_{\varnothing\vee T-}\,|\,\mathscr{F}_{\mathscr{U}}]\mathbf{1}_{G_{\mathscr{U}}} \leqslant C\varepsilon\mathbf{1}_{(T=T^{(1)})} \tag{7.12}$$

$$\mathbf{E}[(e^{-A}\check{Z}_{\varnothing\vee T-})^2\,|\,\mathscr{F}_{\mathscr{U}}]\mathbf{1}_{G_{\mathscr{U}}} \leqslant C\varepsilon^2\mathbf{1}_{(T=T^{(1)})} \tag{7.13}$$

$$\mathbf{E}[\check{Y}_{\varnothing\vee T-}\,|\,\mathscr{F}_{\mathscr{U}}]\mathbf{1}_{G_{\mathscr{U}}} \leqslant e^A\eta \leqslant e^{-A}, \tag{7.14}$$

be (6.3). Equations (7.11) and (7.14) and Chebychev's inequality applied to (7.10) and (7.13) together with (6.1) now give

$$\mathbf{P}(\widehat{G}\cup\check{G}\,|\,G_{\mathscr{U}}) \leqslant 1-\varepsilon^{5/4}. \tag{7.15}$$

The lemma now follows from (7.5), (7.6) and (7.15). □

The probability of G_{nbab}.

Lemma 7.5. *We have $\mathbf{P}((G_{\mathrm{nbab}})^c\,|\,\mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant C\varepsilon^2$ for large A and a.*

Proof. Define $R_1 = \widehat{R}_{[T^-,\Theta_1+e^Aa^2]}+R^{\mathrm{fug}}_{[T,\Theta_1+e^Aa^2]}+\check{R}_{[T^-,\Theta_1+e^Aa^2]}$. By Markov's inequality, Lemmas 5.6, and 5.7 and Corollary 6.7 we have

$$\mathbf{P}(R_1>0\,|\,\mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant C(Z_\Delta e^A/a+Y_\Delta) \leqslant C\varepsilon^2,$$

by the definition of G_Δ, (6.2) and (6.3). Now, by Corollary 6.13,

$$\mathbf{E}[\bar{Z}_T\,|\,\mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} \leqslant CA\mathscr{E}_{\mathscr{U}} \leqslant CAe^{A/3} \quad\text{and}\quad \mathbf{E}[\bar{Y}_T\,|\,\mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} \leqslant C\mathscr{E}_{\mathscr{U}} \leqslant Ce^{A/3},$$

where we used the fact that $\check{Z}^{(1)}_{\varnothing} \leqslant \varepsilon e^A$ by the definition of a breakout. By Proposition 6.2 and the inequality $1-e^{-x}\leqslant x$, the probability conditioned on $\mathscr{F}_\Delta$ and on G_Δ that a check- or bar-particle breaks out between T and Θ_1 is now bounded by $p_Be^{A/3}\leqslant C\varepsilon^2$ by (6.14) and (6.2). This yields the lemma. □

The particles at the time Θ_1. The probability of G_1. Recall that from the time $T^+ = T + \tau_{\max}^{(\mathscr{U},T)}$ on, we move the barrier according to the function f_Δ, which is equivalent to having the variable drift $-\mu_t = -\mu - f_\Delta(t/a^2)/a^2$. On G_{fug}, we have $T^+ \leqslant T + \zeta$ and $\Theta_1 = T + e^A a^2$. Furthermore, $0 < \Delta \leqslant C\log(1/\varepsilon)$ on G_Δ and by the hypotheses on the functions (f_x) and (6.2) it follows,

$$\text{on } G_\Delta : \|f_\Delta\| \leqslant \sqrt{a} \quad \text{and} \quad \Delta - X^{[1]}_{\Theta_1} = O(e^{-A/2}), \tag{7.16}$$

where $\|\cdot\|$ is defined in (5.4).

Lemma 7.6. *Suppose (HB$_0$). Then $\mathbf{P}(G_1) \geqslant 1 - C\varepsilon^{5/4}$ for large A and a.*

Proof. By Proposition 5.2, Lemmas 5.1 and 7.5, Corollaries 6.7 and 6.13 and (7.16),

$$\begin{aligned}\mathbf{E}_{\mathrm{nbab}}[e^{-A}Z_{\Theta_1} \,|\, \mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} &= e^{-\Delta-A}(Z_\Delta + A\mathscr{E}_{\mathscr{U}})(1+O(e^{-A/2})) = 1 + O(e^{-A/2})\\ \mathrm{Var}_{\mathrm{nbab}}(e^{-A}Z_{\Theta_1} \,|\, \mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} &\leqslant C\varepsilon e^{-\Delta-A}\left(Z_\Delta e^A/a + Y_\Delta + \mathscr{E}_{\mathscr{U}}\right) \leqslant C\varepsilon e^{-2A/3},\end{aligned}$$

such that by Chebychev's inequality and (6.2),

$$\mathbf{P}_{\mathrm{nbab}}(|e^{-A}Z_{\Theta_1} - 1| \geqslant \varepsilon^{3/2}) \leqslant C\varepsilon^2. \tag{7.17}$$

Furthermore, by Lemma 6.9, Corollary 6.13 and Markov's inequality, we have

$$\mathbf{P}_{\mathrm{nbab}}(\operatorname{supp}\nu_{\Theta_1} \subset (0,a),\ Y_{\Theta_1} \leqslant \eta \,|\, \mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant e^A/(a\eta) + C\eta e^A \leqslant C\varepsilon^2, \tag{7.18}$$

by (6.3) and (6.2). The lemma now follows from (7.17) and (7.18) together with Lemmas 7.4 and 7.5. □

Remark 7.7. Lemma 7.6 obviously still holds if one replaces G_1 by $G_1 \cap \{|e^{-A}Z_{\Theta_1} - 1| < \frac{1}{2}\varepsilon^{3/2}\}$. This will be needed in Sections 8 and 9.

The Fourier transform of the barrier process. Define $\Delta_{\mathrm{drift}} = e^{-A}(\widehat{Z}_{T-} + \check{Z}_{\varnothing\vee T-}) - 1$ and $\Delta_{\mathrm{jump}} = e^{-A}Z^{(\mathscr{U},T)}$, which are independent random variables. Recall that $\Delta > 0$ on G_Δ. By (7.12) and Lemmas 6.18 and 7.4, we have

$$\mathbf{E}[e^{-A}\check{Z}_{\varnothing\vee T-} \,|\, G_{\mathscr{U}}] \leqslant C\varepsilon\mathbf{P}(T = T^{(1)} \,|\, G_{\mathscr{U}}) \leqslant CA\varepsilon^2,$$

such that with (7.9), we get,

$$\mathbf{E}\left[\Delta_{\mathrm{drift}} \,|\, G_{\mathscr{U}}\right] = \pi^2\gamma_0(A + \log\varepsilon + c + o(1)). \tag{7.19}$$

Furthermore, (7.10) and (7.13) and the inequality $(x+y)^2 \leqslant 2(x^2+y^2)$ yield

$$\mathbf{E}\left[(\Delta_{\mathrm{drift}})^2 \,|\, G_{\mathscr{U}}\right] = O(\varepsilon^2 A^2). \tag{7.20}$$

Equation (7.20) and (6.1) imply

$$\mathbf{P}(|\Delta_{\mathrm{drift}}| \geqslant \varepsilon^{1/3},\ G_{\mathscr{U}}) = O(\varepsilon^{4/3}A^2) = O(\varepsilon^{7/6}). \tag{7.21}$$

Fix $\lambda \in \mathbf{R}$. Since $\log(1+x+y) = \log(1+x) + \log((1+y)/(1+x))$, by (7.16), (7.21) and Lemma 7.4,

$$\begin{aligned}\mathbf{E}[e^{i\lambda X^{[\infty]}_{\Theta_1}}] &= \mathbf{E}[e^{i\lambda\Delta}\mathbf{1}_{(|\Delta_{\mathrm{drift}}|<\varepsilon^{1/3})}\mathbf{1}_{G_\Delta}] + O(\varepsilon^{9/8})\\ &= \mathbf{E}[e^{i\lambda\log(1+\Delta_{\mathrm{drift}})}\mathbf{1}_{(|\Delta_{\mathrm{drift}}|<\varepsilon^{1/3})}\mathbf{1}_{G_\Delta}e^{i\lambda\log(1+\frac{\Delta_{\mathrm{jump}}}{1+\Delta_{\mathrm{drift}}})}] + O(\varepsilon^{9/8}),\end{aligned} \tag{7.22}$$

for any $\lambda \in \mathbf{R}$. We will first study the term concerning Δ_{jump}. Write $Z = Z^{(\mathscr{U},T)}$ and let ρ be a real-valued constant with $|\rho| < \varepsilon^{1/3}$. Then,

$$\begin{aligned}\mathbf{E}[e^{i\lambda\log(1+\frac{\Delta_{\text{jump}}}{1+\rho})}] &= \mathbf{E}[e^{i\lambda\log(1+\frac{e^{-A}Z}{1+\rho})}\,|\,Z > \varepsilon e^{A}] + O(\mathbf{P}^{a}((G^{(\varnothing,0)})^{c}))\\ &= \int_{\varepsilon}^{\infty} g(x)\,\mathbf{P}(e^{-A}Z \in \mathrm{d}x\,|\,Z > \varepsilon e^{A}) + O(\varepsilon^{2}),\end{aligned}\tag{7.23}$$

where

$$g(x) = \exp\Big(i\lambda\log\Big(1 + \frac{x}{1+\rho}\Big)\Big).$$

By definition of p_B and η, we have

$$\int_{\varepsilon}^{1} x\mathbf{P}(e^{-A}Z \in \mathrm{d}x\,|\,Z > \varepsilon e^{A}) = (p_B + O(\eta))^{-1}\mathbf{E}[e^{-A}Z\mathbf{1}_{(\varepsilon e^{A} < Z \leqslant e^{A})}] = \pi^{2}\gamma_0(-\log\varepsilon + o(1)),$$

by (4.6), (6.2), (6.3), (6.14) and (6.13). It follows that

$$\begin{aligned}&\int_{\varepsilon}^{\infty} g(x)\,\mathbf{P}(e^{-A}Z \in \mathrm{d}x\,|\,Z > \varepsilon e^{A})\\ &\qquad = 1 + i\lambda\pi^{2}\gamma_0(1+\rho)^{-1}(-\log\varepsilon + o(1)) + \int_{\varepsilon}^{\infty} h(x)\mathbf{P}(e^{-A}Z \in \mathrm{d}x\,|\,Z > \varepsilon e^{A}),\end{aligned}\tag{7.24}$$

where we define $h(x) = g(x) - 1 - i\lambda(1+\rho)^{-1}x\mathbf{1}_{(x\leqslant 1)}$ for $x \geqslant 0$. Denote by $h^{-}(x)$ its left-hand derivative. Note that $|h(x)| \leqslant C(1 \wedge x^{2})$ and $|h^{-}(x)| \leqslant C(x^{-1} \wedge x^{2})$ for $x \geqslant 0$. By integration by parts, (4.5) and (6.13), we have

$$\begin{aligned}&\int_{\varepsilon}^{\infty} h(x)\mathbf{P}(e^{-A}Z \in \mathrm{d}x\,|\,Z > \varepsilon e^{A})\\ &= h(\varepsilon) + p_B^{-1}(1+o(1))\Big(\int_{\varepsilon}^{\infty} h^{-}(x)\mathbf{P}(Z > xe^{A})\,\mathrm{d}x + (h(1+) - h(1))\mathbf{P}(Z > e^{A})\Big)\\ &= \pi^{2}\gamma_0\Big(\int_{0}^{\infty} h^{-}(x)\frac{1}{x}\,\mathrm{d}x + (h(1+) - h(1)) + o(1)\Big)\\ &= \pi^{2}\gamma_0(1+o(1))\int_{0}^{\infty} h(x)\frac{1}{x^{2}}\,\mathrm{d}x.\end{aligned}\tag{7.25}$$

Now, one readily sees that

$$\int_{0}^{\infty} h(x)\frac{1}{x^{2}}\,\mathrm{d}x = i\lambda(c' + o(1) + O(\rho)) + \int_{0}^{\infty} e^{i\lambda x} - 1 - i\lambda x\mathbf{1}_{(x\leqslant 1)}\,\Lambda(\mathrm{d}x),\tag{7.26}$$

where $\Lambda(\mathrm{d}x)$ is as in the statement of Proposition 7.3 and c' is a numerical constant. Equations (7.23), (7.24), (7.25) and (7.26) and the Taylor expansion of e^{-x} at $x = 0$ now yield

$$\begin{aligned}&\mathbf{E}[e^{i\lambda\log(1+\frac{\Delta_{\text{jump}}}{1+\rho})}]\\ &\quad = \exp\Big[\gamma_0\Big(i\lambda\pi^{2}(-\log\varepsilon + c' + o(1) + \pi^{2}\int_{0}^{\infty} e^{i\lambda x} - 1 - i\lambda x\mathbf{1}_{(x\leqslant 1)}\,\Lambda(\mathrm{d}x)\Big) + O(\varepsilon|\log\varepsilon|\rho)\Big].\end{aligned}\tag{7.27}$$

Coming back to (7.22), we have by the Taylor expansion of $(1+x)^{i\lambda}$ at $x = 0$,

$$\begin{aligned}&\mathbf{E}[e^{i\lambda(\log(1+\Delta_{\text{drift}})+O(\varepsilon|\log\varepsilon|\Delta_{\text{drift}}))}\mathbf{1}_{(|\Delta_{\text{drift}}|<\varepsilon^{1/3})}\mathbf{1}_{G_\Delta}]\\ &\quad = \mathbf{E}[(1 + i\lambda\Delta_{\text{drift}} + O(\Delta_{\text{drift}}^{2}))\mathbf{1}_{(T\leqslant\sqrt{\varepsilon}a^{3})}] + O(\varepsilon^{9/8}) && \text{by Lemma 7.4 and (7.21)}\\ &\quad = 1 + i\lambda\pi^{2}\gamma_0(A + \log\varepsilon + c + o(1)) + \mathbf{P}(T > \sqrt{\varepsilon}a^{3}) + O(\varepsilon^{9/8}) && \text{by (7.19) and (7.20)}\\ &\quad = \exp i\lambda\pi^{2}\gamma_0(A + \log\varepsilon + c + o(1) + O(\varepsilon^{1/8})),\end{aligned}$$

where the last equation follows from Lemma 6.3 and the Taylor expansion of e^x at $x = 0$. This equation, together with (7.22) and (7.27) and the fact that Δ_{jump} is independent from Δ_{drift}, T and G_{bulk}, yields (7.3) in the case $n = 1$.

7.3 Proof of Theorems 7.1 and 7.2

We define the process $(X''_t)_{t\geqslant 0}$ by $X''_t = X^{[\infty]}_{\Theta_{\lfloor t\gamma_0^{-1} a^3\rfloor}} - \pi^2 At$.

Proposition 7.8. *Suppose (HB). Then, as A and a go to infinity, the process $(X''_t)_{t\geqslant 0}$ converges in law (with respect to Skorokhod's topology) to the Lévy process $(L_t)_{t\geqslant 0}$ defined in Theorem 1.1.*

Proof. Conditioning on $\mathscr{F}_0$, we can assume without loss of generality that the initial configuration ν_0 is deterministic and that G_0 is verified. Denote by $(\mathscr{F}''_t)_{t\geqslant 0}$ the natural filtration of the process X''_t and note that $\mathscr{F}''_t = \mathscr{F}''_{\gamma_0\lfloor t\gamma_0^{-1}\rfloor} \subset \mathscr{F}_{T_{\lfloor t\gamma_0^{-1}\rfloor}}$. In order to show convergence of the finite-dimensional distributions, it is enough to show (see Proposition 3.1 in [41] or Lemma 8.1 in [28], p. 225), that for every $\lambda \in \mathbf{R}$ and $t, s \geqslant 0$,

$$\mathbf{E}\Big[\Big|\mathbf{E}[e^{i\lambda X''_{t+s}} \mid \mathscr{F}''_t] - e^{i\lambda X''_t} e^{s\kappa(\lambda)}\Big|\Big] \to 0, \tag{7.28}$$

as A and a go to infinity. Now, define $n := \lfloor t\gamma_0^{-1}\rfloor$ and $m := \lfloor (t+s)\gamma_0^{-1}\rfloor$, such that $(m-n)\gamma_0 = s + A^{-1}o(1)$, by (6.1) and (6.14). Then we have by Proposition 7.3,

$$\begin{aligned}\mathbf{E}[e^{i\lambda(X''_{t+s}-X''_t)} \mid \mathscr{F}_{\Theta_m}]\mathbf{1}_{G_m} &= e^{-i\lambda As}\mathbf{E}[e^{i\lambda(X^{[\infty]}_{\Theta_m}-X^{[\infty]}_{\Theta_n})} \mid \mathscr{F}_{\Theta_m}]\mathbf{1}_{G_n}\\ &= \exp\Big(s\big(\kappa(\lambda) + o(1) + O(\varepsilon^b)\big)\Big)\mathbf{1}_{G_n}.\end{aligned} \tag{7.29}$$

In total, we get for A and a large enough,

$$\mathbf{E}\Big[\Big|\mathbf{E}[e^{i\lambda(X''_{t+s}-X''_t)} \mid \mathscr{F}''_t] - e^{s\kappa(\lambda)}\Big|\Big] \leqslant e^{s\kappa(\lambda)}\mathbf{E}[|e^{s(o(1)+O(\varepsilon^b))} - 1|] + \mathbf{P}(G_m^c).$$

By Proposition 7.3, this goes to 0 as A and a go to infinity, which proves (7.28).

In order to show tightness in Skorokhod's topology, we use Aldous' famous criterion [1] (see also [10], Theorem 16.10): If for every $M > 0$, every family of $(\mathscr{F}''_t)$-stopping times $\tau = \tau(A,a)$ taking only finitely many values, all of which in $[0, M]$ and every $h = h(A,a) \geqslant 0$ with $h(A,a) \to 0$ as A and a go to infinity, we have

$$X''_{\tau+h} - X''_\tau \to 0, \qquad \text{in probability as } A \text{ and } a \text{ go to infinity}, \tag{7.30}$$

then tightness follows for the processes X''_t (note that the second point in the criterion, namely tightness of X''_t for every fixed t, follows from the convergence in finite-dimensional distributions proved above). Now let τ be such a stopping time and let V_τ be the (finite) set of values it takes. We first note that since $G_n \supset G_{n+1}$ for every $n \in \mathbf{N}$, we have for every $t \in V_\tau$ and every A and a large enough,

$$\mathbf{P}(G^c_{\lfloor t\gamma_0^{-1}\rfloor}) \leqslant \mathbf{P}(G^c_{\lfloor M\gamma_0^{-1}\rfloor}) = O(M\varepsilon^b). \tag{7.31}$$

by Proposition 7.3. Moreover, since $\mathscr{F}''_t \subset \mathscr{F}_{T_{\lfloor t\gamma_0^{-1}\rfloor}}$ for every $t \geqslant 0$, we have for every $\lambda > 0$,

$$\begin{aligned}\mathbf{E}[e^{i\lambda(X''_{\tau+h}-X''_\tau)}] &= \sum_{t\in V_\tau}\mathbf{E}\Big[e^{i\lambda(X''_{t+h}-X''_t)}\mathbf{1}_{(\tau=t)}\Big]\\ &= \sum_{t\in V_\tau}\mathbf{E}\Big[\mathbf{E}[e^{i\lambda(X''_{t+h}-X''_t)} \mid \mathscr{F}_{T_{\lfloor t\gamma_0^{-1}\rfloor}}]\mathbf{1}_{(\tau=t)}\mathbf{1}_{G_{\lfloor t\gamma_0^{-1}\rfloor}}\Big] + O(M\varepsilon^b) && \text{by (7.31)}\\ &= e^{h(\kappa(\lambda)+o(1)+O(\varepsilon^b))}(1 - O(M\varepsilon^b)) + O(M\varepsilon^b), && \text{by (7.29),}\end{aligned}$$

which converges to 1 as A and a go to infinity. This implies (7.30) and therefore proves tightness in Skorokhod's topology, since M was arbitrary. Together with the convergence in finite-dimensional distributions proved above, the lemma follows. □

A coupling with a Poisson process. Let $(V_n)_{n\geqslant 0}$ be a sequence of independent exponentially distributed random variable with mean γ_0. In order to prove convergence of the processes X'_t and X_t, we are going to couple the BBM with the sequence (V_n) in the following way: Suppose we have constructed the BBM until time Θ_{n-1}. Now, on the event G_{n-1}, by Lemma 6.6, the strong Markov property of BBM and the transfer theorem ([38], Theorem 5.10), we can construct the BBM up to time Θ_n such that $\mathbf{P}(|(T_n-\Theta_{n-1})/a^3 - V_n| > \varepsilon^{3/2}) = O(\varepsilon^2)$ (recall that T_n denotes the time of the first breakout after Θ_{n-1}). On the event G^c_{n-1}, we simply let the BBM evolve independently of $(V_j)_{j\geqslant n}$. Setting $G'_n = G_n \cap \{\forall j \leqslant n : |(T_j - \Theta_{j-1})/a^3 - V_j| \leqslant \varepsilon^{3/2}\}$, there is by Lemma 6.6 and Proposition 7.3 a $b > 0$, such that for large A and a,

$$\mathbf{P}(G'_n) \geqslant \mathbf{P}(G_0) - nO(\varepsilon^{1+b}) \tag{7.32}$$

Furthermore, we have $\Theta_n = T_n + a^{5/2}$ on G'_n, whence for large A and a,

$$\text{on } G'_n : |(\Theta_n - \Theta_{n-1})/a^3 - V_n| \leqslant 2\varepsilon^{3/2}. \tag{7.33}$$

This construction now permits us to do the

Proof of Theorem 7.2. Let d denote the Skorokhod metric on the space of cadlag functions $D([0,\infty))$ (see [28], Section 3.5). Let Φ be the space of strictly increasing, continuous, maps of $[0,\infty)$ onto itself. Let $x, x_1, x_2, \ldots$ be elements of $D([0,\infty))$. Then ([28], Proposition 3.5.3), $d(x_n, x) \to 0$ as $n \to \infty$ if and only if for every $M > 0$ there exists a sequence (φ_n) in Φ, such that

$$\sup_{t\in[0,M]} |\varphi_n(t) - t| \to 0, \qquad \text{and} \qquad \sup_{t\in[0,M]} |x_n(\varphi_n(t)) - x(t)| \to 0. \tag{7.34}$$

If $(x'_n)_{n\in\mathbf{N}}$ is another sequence of functions in $D([0,\infty))$, with $d(x'_n, x) \to 0$, then by the triangle inequality and the fact that Φ is stable under the operations of inverse and convolution, we have $d(x_n, x) \to 0$ if and only if there exists a sequence (φ_n) in Φ, such that the first inequality in (7.34) holds and

$$\sup_{t\in[0,M]} |x_n(\varphi_n(t)) - x'_n(t)| \to 0. \tag{7.35}$$

For every A and a, we define the (random) map $\varphi_{A,a} \in \Phi$ by

$$\varphi_{A,a}(\gamma_0(n+r)) = ((1-r)\Theta_n + r\Theta_{n+1})a^{-3}, \text{ with } n \in \mathbf{N},\ r \in [0,1].$$

Let $M > 0$ and define $n_M = \lceil M\gamma_0 \rceil$. Then we have by the monotonicity of $(X^{[\infty]}_t)_{t\geqslant 0}$,

$$\sup_{t\in[0,M]} |\varphi_{A,a}(t) - t| \leqslant \max_{n\in\{0,\ldots,n_M\}} \left|a^{-3}\Theta_n - \gamma_0 n\right|, \tag{7.36}$$

$$\sup_{t\in[0,M]} |X''_t - X'_{\varphi_{A,a}(t)}| \leqslant \max_{n\in\{0,\ldots,n_M\}} A\left|a^{-3}\Theta_n - \gamma_0 n\right|. \tag{7.37}$$

By Doob's L^2 inequality and the fact that $\gamma_0 = E[V_1]$, we get

$$\mathbf{P}\Big(\max_{n\in\{0,\ldots,n_M\}} \left|\sum_{i=1}^n V_i - n\gamma_0\right| > \varepsilon^{1/3}\Big) \leqslant C\varepsilon^{-2/3} n_M \operatorname{Var}(V_i) = O(\varepsilon^{1/3}).$$

Furthermore, on the set G'_{n_M}, we have for every $n \leqslant n_M$, $|\Theta_n - \sum_{i=0}^n V_i| \leqslant O(n_M\varepsilon^{3/2}) = O(\varepsilon^{1/2})$. In total, we get with (7.36) and (7.37), as A and a go to infinity,

$$\forall M > 0 : \sup_{t\in[0,M]} |\varphi_{A,a}(t) - t| \vee |X''_t - X'_{\varphi_{A,a}(t)}| \to 0, \quad \text{in probability}, \tag{7.38}$$

which is equivalent to

$$\sum_{M=1}^{\infty} 2^{-M}\Big[1 \wedge \Big(\sup_{t\in[0,M]} |\varphi_{A,a}(t)-t| \vee |X''_t - X'_{\varphi_{A,a}(t)}|\Big)\Big] \to 0, \quad \text{in probability.} \tag{7.39}$$

Now, suppose that A and a go to infinity along a sequence $(A_n,a_n)_{n\in\mathbf{N}}$ and denote by X'_{A_n,a_n}, X''_{A_n,a_n} and φ_{A_n,a_n} the processes corresponding to these parameters. By Proposition 7.8 and Skorokhod's representation theorem ([10], Theorem 6.7), there exists a probability space, on which the sequence (X''_{A_n,a_n}) converges almost surely as $n\to\infty$ to the limiting Lévy process $L=(L_t)_{t\geqslant 0}$ stated in the theorem. Applying again the representation theorem as well as the transfer theorem, we can transfer the processes X'_{A_n,a_n} and φ_{A_n,a_n} to this probability space in such a way that the convergence in (7.39) holds almost surely, which implies that the convergence in (7.38) holds almost surely as well. By the remarks at the beginning of the proof, it follows that on this new probability space,

$$d(X'_{A_n,a_n},L) \leqslant d(X'_{A_n,a_n},X''_{A_n,a_n}) + d(X''_{A_n,a_n},L) \to 0,$$

almost surely, as $n\to\infty$. This proves the theorem. □

Proof of Theorem 7.1. Let $(t_i^{A,a})_{i=1}^n$ be as in the statement of the theorem. By the virtue of Theorem 7.2, it suffices to show that

$$\mathbf{P}\Big(\forall i : X^{[\infty]}_{t_i^{A,a}a^3} = J_{t_i^{A,a}a^3}\Big) \to 1. \tag{7.40}$$

Suppose for simplicity that $t_i^{A,a}\equiv t_i$ for all i; the proof works exactly the same in the general case. Let $n := \lceil 2(t_k+2)/\gamma_0\rceil$, such that $n = O(\varepsilon^{-1})$, by (6.14). By Chebychev's inequality, we then have

$$\mathbf{P}(\sum_{i=1}^n V_i \leqslant t_k + 2) \leqslant \mathbf{P}(\sum_{i=1}^n (V_i-\gamma_0) \leqslant -\frac{n}{2}\gamma_0) = O(n\operatorname{Var}(V_i)) = O(\varepsilon). \tag{7.41}$$

Furthermore, define the intervals $I_i = t_i + [-2n\varepsilon^{3/2} - e^A/a, 2n\varepsilon^{3/2}]$, $i=1,\dots,k$ and denote by $\mathscr{P}$ the point process on the real line with points at the positions $V_1, V_1+V_2, V_1+V_2+V_3,\dots$. Then $\mathscr{P}$ is a Poisson process with intensity $\gamma_0^{-1} = O(\varepsilon^{-1})$ and thus,

$$\mathbf{P}\Big(\mathscr{P}\cap\bigcup_{i=1}^k I_i \neq \varnothing\Big) = O(\varepsilon^{1/2}). \tag{7.42}$$

We now have

$$\begin{aligned}
\mathbf{P}\Big(\forall i : X^{[\infty]}_{t_ia^3} = J_{t_ia^3}\Big) &\geqslant \mathbf{P}\Big(\nexists (i,j) : t_ia^3 \in [\Theta_j - T_{j-1}, \Theta_j]\Big) && \text{by definition}\\
&\geqslant \mathbf{P}\Big(G'_n,\ \sum_{i=1}^n V_i > t_k+2,\ \mathscr{P}\cap\bigcup_{i=1}^k I_i = \varnothing\Big) && \text{by definition of } G'_n\\
&\geqslant \mathbf{P}(G_0) - O(\varepsilon^b) && \text{by (7.32), (7.41), (7.42).}
\end{aligned}$$

Letting A and a go to infinity and using the hypothesis (HB) yields (7.40) and thus proves the theorem. □

8 The B$^\flat$-BBM

In this section, we define and study the B$^\flat$-BBM, which is obtained from the B-BBM by killing some of its particles. It will be used in Section 10 to bound the N-BBM from below.

8.1 Definition of the model

We will use throughout the notation from Section 7. Furthermore, we fix $\delta \in (0,1/2)$ and $K \geqslant 1$ such that $E_K \leqslant \delta/10$, where E_K was defined in (2.6). Define $N = \lceil 2\pi e^{A+\delta} a^{-3} e^{\mu a} \rceil$, where $\mu = \sqrt{1-\pi^2/a^2}$. We will then use interchangeably the phrases "as A and a go to infinity" and "as N go to infinity". Furthermore, in the definition of these phrases (see Section 6.1), A_0 and the function $a_0(A)$ may now also depend on δ. The symbols C_δ and $C_{\delta,\alpha}$ have the same meaning as C (see Section 5.1), except that they may depend on δ or δ and α as well, α being defined later.

The B$^\flat$-BBM is then defined as follows: Given a possibly random initial configuration ν_0 of particles in $(0,a)$, we let particles evolve according to B-BBM with barrier function given by (7.1), where, in addition, we colour the particles white and red as follows: Initially, all particles are coloured white. As soon as a white particle has N or more white particles to its right, it is coloured red[6]. Children inherit the colour of their parent. At each time Θ_n, $n \geqslant 1$, all the red particles are killed immediately and the process goes on with the remaining particles. See Figure 3 for a graphical description.



Figure 3: The B$^\flat$-BBM with parameter $N = 6$ (no breakout is shown). White and red particles are drawn with solid and dotted lines, respectively. The blobs indicate when a white particle is colored red, the circles show the six white particles living at that time (thick circles correspond to two particles at the same position).

For an interval $I \subset \mathbf{R}_+$, we define the stopping line $\mathscr{L}^{\mathrm{red}}_{\varnothing,I}$ by $(u,t) \in \mathscr{L}^{\mathrm{red}}_{\varnothing,I}$ if and only if the particle u gets coloured red at the time t and has been white up to the time t, with $t \in I$. We then set $Z^{\mathrm{red}}_{\varnothing,I}$ and $Y^{\mathrm{red}}_{\varnothing,I}$ by summing respectively w_Z and w_Y over the particles of this stopping line. Furthermore, we define $\mathscr{N}^{\mathrm{red}}_t$ and $\mathscr{N}^{\mathrm{white}}_t$ to be the subsets of $\mathscr{N}_t$ formed by the red and white particles, respectively, and define Z^{red}_t, Y^{red}_t, Z^{white}_t and Y^{white}_t accordingly.

---

[6] This can be ambiguous if there are more than one of the particles at the same position, for example when the left-most particle branches. In order to eliminate this ambiguity, induce for every $t \geqslant 0$ a total order on the particles in $\mathscr{N}(t)$ by $u < v$ iff $X_u(t) < X_v(t)$ or $X_u(t) = X_v(t)$ and u precedes v in the *lexicographical order on U*. Whenever there are more than N white particles, we then colour the particles in this order, which is well defined.

Let $\nu_t^\flat$ be the configuration of white particles at the time t and abuse notation by setting $\nu_n^\flat = \nu_{\Theta_n}^\flat$. We set $G_{-1}^\flat = \Omega$ and for each $n \in \mathbf{N}$, we define the event $G_n^\flat$ to be the intersection of $G_{n-1}^\flat$ with the following events (we omit the braces).

- $\operatorname{supp} \nu_n^\flat \subset (0,a)$,
- $\mathscr{N}_{\Theta_n}^{\text{white}} \subset U \times \{\Theta_n\}$ and $\Theta_n > T_n^+$ (for $n > 0$),
- $|e^{-A} Z_{\Theta_n}^{\text{white}} - 1| \leqslant \varepsilon^{3/2}$ and $Y_{\Theta_n}^{\text{white}} \leqslant \eta$.
- $\mathbf{P}\Big(Z^{\text{red}}_{\varnothing,[\Theta_n,\Theta_n+Ka^2]} + Y^{\text{red}}_{\varnothing,[\Theta_n,\Theta_n+Ka^2]} \leqslant a^{-1/2} \,\Big|\, \mathscr{F}_{\Theta_n}\Big) \geqslant 1 - \varepsilon^2$.

The last event is of course uniquely defined up to a set of probability zero. Note that $G_n^\flat \in \mathscr{F}_{\Theta_n}$ for each $n \in \mathbf{N}$. Furthermore, we define the predicates

(HB$^\flat$) (The law of) ν_0 is such that $\mathbf{P}(G_0^\flat) \to 1$ as A and a go to infinity.
(HB$_0^\flat$) ν_0 is deterministic and such that $G_0^\flat$ holds.
(HB$_\perp^\flat$) ν_0 is obtained from $\lfloor e^{-\delta} N \rfloor$ particles distributed independently according to the density proportional to $\sin(\pi x/a) e^{-\mu x} \mathbf{1}_{(0,a)}(x)$.

We now state the important results on the B$^\flat$-BBM.

Lemma 8.1. *(HB$_\perp^\flat$) implies (HB$^\flat$) for large A and a.*

Proposition 8.2. *Proposition 7.3 still holds for the B$^\flat$-BBM, with G_n replaced by $G_n^\flat$. The same is true for Theorems 7.1 and 7.2, with (HB) replaced by (HB$^\flat$).*

Recall the definition of qu_α^N and x_α from the introduction and of (Ht) from Section 7.

Proposition 8.3. *Suppose (HB$_\perp^\flat$). Let $(t_j^{A,a})$ satisfy (Ht) and let $\alpha \in (0, e^{-2\delta})$. As A and a go to infinity,*

$$\mathbf{P}(\forall j : \mathrm{qu}_\alpha^N(\nu^\flat_{a^3 t_j^{A,a}}) \geqslant x_{\alpha e^{2\delta}}) \to 1.$$

8.2 Preparatory lemmas

In this section, we will establish some fine estimates for the number of particles of the process, which will be used later to bound the number of creations of red particles. If a particle $u \in \mathscr{N}_0(t)$ satisfies $\tau_l(u) \leqslant t < \sigma_{l+1}(u)$ for some $l \geqslant 1$, we call it an "in between" particle (because it is in some kind of intermediary stage) and otherwise a "regular" particle. We then define

$$N_t(r) = \sum_{(u,s) \in \mathscr{N}_t} \mathbf{1}_{(X_u(s) \geqslant r)}.$$

The main lemma in this section is the following:

Lemma 8.4. *Suppose (HB$_0$). Let $Ka^2 \leqslant t \leqslant t_{6.21}$. Then, for $0 \leqslant r \leqslant 9a/10$ and every $\alpha > 0$ there exists $C_{\delta,\alpha}$, such that for large A and a,*

$$\mathbf{P}(N_t(r) \geqslant e^{-(\delta/2+r/3)} N \,|\, T > t) \leqslant C_{\delta,\alpha} A^2 \varepsilon \Big(\frac{t}{a^3} + \eta\Big) e^{-(\frac{4}{3}-\alpha) r}$$

Furthermore, conditioned on $\mathscr{F}_\Delta$, for $t \leqslant \Theta_1 + e^A a^2$, for large A and a,

$$\mathbf{P}_{\text{nbab}}(N_t(r) \geqslant e^{-(\delta/2+r/3)} N \,|\, \mathscr{F}_\Delta) \mathbf{1}_{G_\Delta} \leqslant C_{\delta,\alpha} A^2 \varepsilon^2 e^{-(\frac{4}{3}-\alpha) r}.$$

The following lemma about BBM conditioned not to break out will be used many times in the proof:

Lemma 8.5. *Let f be a barrier function, $0 \leqslant t \leqslant t_0 \leqslant t_{6.21}$ and suppose that* $\mathrm{Err}(f,t) \leqslant C$. *For $x \in (0,a)$, define $\widehat{\mathbf{P}}^x_f = \mathbf{P}^x_f(\cdot \,|\, T > t_0)$. Then, for all $r \leqslant (9/10)a$ and for large A and a,*

$$\widehat{\mathbf{E}}^x_f[N^{(1+)}_t(r)] \leqslant CAe^{-A}N(1+\mu r)e^{-\mu r}\big(1+(\eta r^2/a)\big)\Big(\frac{t}{a^3}w_Z(x)+w_Y(x)\Big) \tag{8.1}$$

$$\widehat{\mathbf{E}}^x_f[(N_t(r))^2] \leqslant C(1+r^4)e^{-2r}A^2\varepsilon e^{-A}N^2\Big(\frac{t}{a^3}w_Z(x)+w_Y(x)\Big). \tag{8.2}$$

Proof. By Lemma 5.7, the upper bound of Lemma 6.8 is still valid with varying drift (see also Remark 6.1). This gives,

$$\widehat{\mathbf{E}}^x_f[Z^{(1+)}_{\varnothing,[0,t]}] \leqslant CA\Big(\frac{t}{a^3}w_Z(x)+w_Y(x)\Big), \quad \text{and} \quad \widehat{\mathbf{E}}^x_f[Y^{(1+)}_{\varnothing,[0,t]}] \leqslant \eta\widehat{\mathbf{E}}^x_f[Z^{(1+)}_{\varnothing,[0,t]}]. \tag{8.3}$$

where the second inequality follows from Lemma 5.8. Equation (8.1) then follows from Lemma 5.4 and (8.3), noting that $a - y - f(\zeta/a^2) \geqslant r + a/20$ for large a, by the hypotheses on r and f.

For the second equation, suppose for simplicity that $f \equiv 0$. By Lemma 6.10, we have for all $x \in (0,a)$,

$$\begin{aligned}\widehat{\mathbf{E}}^x[(N_t(r))^2] &\leqslant \widehat{\mathbf{E}}^x[N_t(r)] + \beta_0 m_2 \int_0^t\int_0^a \widehat{\mathbf{p}}^{f,(0+)}_s(x,z)\widehat{\mathbf{E}}^{(z,s)}[N_t(r)]^2\,\mathrm{d}z\,\mathrm{d}s\\ &\quad + \widehat{\mathbf{E}}^x\Big[\sum_{(u,s)\in\mathscr{R}_t}\ \sum_{(v_1,s_1),(v_2,s_2)\in\mathscr{S}^{(u,s)},\, v_1\neq v_2} \widehat{\mathbf{E}}^{(X_{v_1}(s_1),s_1)}[N_t(r)]\widehat{\mathbf{E}}^{(X_{v_2}(s_2),s_2)}[N_t(r)]\Big].\end{aligned} \tag{8.4}$$

Now, for large A we have by (5.43) and Lemma 5.1,

$$\int_0^t\int_0^a \widehat{\mathbf{p}}^{f,(0+)}_s(x,z)\widehat{\mathbf{E}}^{(z,s)}[N_t(r)]^2\,\mathrm{d}z\,\mathrm{d}s \leqslant C\widehat{\mathbf{E}}\Big[\sum_{(v,s_0)\in\mathscr{S}_t}\int_{s_0}^t\int_0^a \mathbf{p}_{s-s_0}(X_v(s_0),z)\widehat{\mathbf{E}}^{(z,s)}[N_t(r)]^2\,\mathrm{d}z\,\mathrm{d}s\Big] \tag{8.5}$$

With Lemmas 5.4 and 6.8 and (8.1), we have as in the proof of Lemma 5.5, for any $s_0 \leqslant t$,

$$\int_{s_0}^t\int_0^a \mathbf{p}_{s-s_0}(x,z)\widehat{\mathbf{E}}^{(z,s)}[N_t(r)]^2\,\mathrm{d}z\,\mathrm{d}s \leqslant CA^2e^{-2A}N^2(1+r^4)e^{-2\mu r}\Big(\frac{t}{a^3}w_Z(x)+w_Y(x)\Big). \tag{8.6}$$

Equations (8.5) and (8.6) together with Lemma 6.8 now give

$$\int_0^t\int_0^a \widehat{\mathbf{p}}^{f,(0+)}_s(x,z)\widehat{\mathbf{E}}^{(z,s)}[N_t(r)]^2\,\mathrm{d}z\,\mathrm{d}s \leqslant CA^2e^{-2A}N^2(1+r^4)e^{-2\mu r}\Big(\frac{t}{a^3}w_Z(x)+Aw_Y(x)\Big). \tag{8.7}$$

Moreover, we have

$$\begin{aligned}&\widehat{\mathbf{E}}^x\Big[\sum_{(u,s)\in\mathscr{R}_t}\ \sum_{(v_1,s_1),(v_2,s_2)\in\mathscr{S}^{(u,s)},\, s_1<s_2} \widehat{\mathbf{E}}^{X_{v_1}(s_1)}[N_t(r)]\widehat{\mathbf{E}}^{X_{v_2}(s_2)}[N_t(r)]\Big]\\ &\quad\leqslant \widehat{\mathbf{E}}^x\Big[\sum_{(u,s)\in\mathscr{R}_t}\Big(\sum_{(v,s_0)\in\mathscr{S}^{(u,s)}}\widehat{\mathbf{E}}^{X_v(s_0)}[N_t(r)]\Big)^2\Big]\\ &\quad\leqslant C(1+r^4)e^{-2\mu r}A^2e^{-2A}N^2\widehat{\mathbf{E}}^x\Big[\sum_{(u,s)\in\mathscr{R}_t}(Z^{(u,s)})^2\Big]\\ &\quad\leqslant C(1+r^4)e^{-2\mu r}\varepsilon e^{-A}A^2N^2\Big(\frac{t}{a^3}w_Z(x)+w_Y(x)\Big),\end{aligned} \tag{8.8}$$

where the last inequality follows from (6.17) and Lemma 5.6. Equation (8.2) now follows from (6.2), (8.1), (8.4), (8.7) and (8.8). □

Corollary 8.6. *With the hypotheses from Lemma 8.5, we have for large A and a,*

$$\widehat{\mathbf{E}}^x_f[N_t(r)] \leqslant CAe^{-A}N(1+r^2)e^{-\mu r}e^{-(a-x)/2}.$$

Proof. This follows immediately from Lemmas 5.1, 5.4 and 8.5 and Corollary 6.7. □

Lemma 8.7. *Suppose (HB$_0$). Let $\alpha > 0$. For large A and a, we have for every $0 \leqslant r \leqslant 9a/10$ and $t \leqslant \Theta_1 + e^A a^2$,*

$$\mathbf{P}_{\mathrm{nbab}}(\overline{N}_t(r) > e^{-A/6-r/3}N \,|\, \mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant C_\alpha A^2\varepsilon e^{-2A/3}e^{-(\frac{4}{3}-\alpha)r}. \tag{8.9}$$

Proof. Conditioned on $\mathscr{F}_\Delta$, define $\widehat{\mathbf{P}} = \mathbf{P}(\cdot \,|\, T > \Theta_1 + e^A a^2)$. Then,

$$\mathrm{Var}_{\mathrm{nbab}}(\overline{N}_t(r) \,|\, \mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant C\sum_{i=0}^{l}\sum_{u\preceq\mathscr{U}} \mathbf{1}_{(\sigma_i\leqslant d_u<\tau_i)}(k_u-1)\widehat{\mathbf{E}}^{(X_u(d_u),d_u)}_{X^{[1]}}[N_t(r)^2], \tag{8.10}$$

and Lemma 8.5 now implies

$$\mathrm{Var}_{\mathrm{nbab}}(\overline{N}_t(r) \,|\, \mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant C(1+r^4)e^{-2r}A^2\varepsilon e^{-A}N^2\mathscr{E}_{\mathscr{U}}. \tag{8.11}$$

Furthermore, by Corollary 8.6,

$$\mathbf{E}_{\mathrm{nbab}}[\overline{N}_t(r) \,|\, \mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} \leqslant CAe^{-A}N(1+r^2)e^{-\mu r}\mathscr{E}_{\mathscr{U}}. \tag{8.12}$$

Equation (8.9) now follows from (8.11) and (8.12) together with the conditional Chebychev inequality and the fact that $\mathscr{E}_{\mathscr{U}} \leqslant e^{A/3}$ on $G_{\mathscr{U}} \subset G_\Delta$. □

Proof of Lemma 8.4. Assume $Ka^2 \leqslant t \leqslant t_{6.21}$. By the hypothesis on Z_0, the definition of K, Proposition 5.3 and Corollary 6.7, we have for large A and a,

$$\mathbf{E}[N^{(0)}_t(r) \,|\, T > t] \leqslant (1-7\delta/8)(1+\mu r)e^{-\mu r}N, \tag{8.13}$$

Moreover, we have by Lemma 8.5,

$$\mathbf{E}[N^{(1+)}_t(r) \,|\, T > t] \leqslant C(1+\mu r)e^{-\mu r}N(1+\eta r)Ae^{-A}(\frac{t}{a^3}Z_0 + Y_0) \leqslant C\sqrt{\varepsilon}A(1+r^2)e^{-\mu r}N, \tag{8.14}$$

by the hypotheses on Z_0 and Y_0. Equations (8.13) and (8.14) now give for large A and a, for all $r \in [0,a]$,

$$\mathbf{E}[N_t(r) \,|\, T > t] \leqslant (e^{-\delta/2} - \delta/8)e^{-r/3}N, \tag{8.15}$$

Furthermore, by Lemma 8.5 and the hypotheses on Z_0 and Y_0, we have

$$\mathrm{Var}(N_t(r) \,|\, T > t) \leqslant CA^2\varepsilon(1+r^4)e^{-2r}(t/a^3 + \eta), \tag{8.16}$$

Chebychev's inequality, (8.15) and (8.16) yield the first equation of the lemma.

Conditioned on $\mathscr{F}_\Delta$, let $T \leqslant t \leqslant \Theta_1 + e^A a^2$. Define $\widetilde{N}^{\mathrm{bulk}}_t(r)$ to be the number of hat- and check-particles to the right of r at time t that have not hit a between T^- and t and likewise $\widetilde{N}^{\mathrm{fug}}_t(r)$ the number of fug-particles with the same properties. Set $M_t = e^{-X^{[1]}_t} = (1+\Delta_t)^{-1}$, where $\Delta_t = \overline{\theta}((t-T^+)/a^2)\Delta$. Note that we have on G_Δ: $|\Delta - e^{-A}Z^{(\mathscr{U},T)}| \leqslant \varepsilon^{1/8}$.

Define $\alpha_r = (1+\mu r)e^{-\mu r}$. By Lemma 5.1 and Proposition 5.3, we have for large A and a,

$$\mathbf{E}[\widetilde{N}^{\mathrm{bulk}}_t(r) \,|\, \mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} \leqslant \alpha_r e^{o(1)}NM_t(1+E_K)e^{-A}(\widehat{Z}_{T^-} + \check{Z}_{\varnothing,T^-})\mathbf{1}_{G_\Delta} \leqslant \alpha_r NM_t(1-3\delta/4), \tag{8.17}$$

by the definition of $\widehat{G}$ and $\breve{G}$. Furthermore, we have by Proposition 5.3 and Lemma 5.1, for large a,

$$\begin{aligned}\mathbf{E}[\widetilde{N}_t^{\mathrm{fug}}(r)\,|\,\mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} &\leqslant \alpha_r e^{o(1)} N M_t \left(\overline{\theta}\left((t-T)/a^2\right) + O\left((y+\Delta+r)^2\eta/a\right)\right) e^{-A} Z^{(\mathscr{U},T)}\mathbf{1}_{G_\Delta} \\ &\leqslant \alpha_r N M_t \left(\Delta_t + O(\varepsilon^{1/8} + a^{-1}r^2)\right)\mathbf{1}_{G_\Delta},\end{aligned} \tag{8.18}$$

Equations (8.17) and (8.18) now give,

$$\mathbf{E}[\widetilde{N}_t^{\mathrm{bulk}}(r) + \widetilde{N}_t^{\mathrm{fug}}(r)\,|\,\mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} \leqslant \alpha_r N\left(1 - \delta/2 + O(a^{-1}r^2)\right). \tag{8.19}$$

Similarly, one has by Lemma 5.5,

$$\mathrm{Var}(\widetilde{N}_t^{\mathrm{bulk}}(r) + \widetilde{N}_t^{\mathrm{fug}}(r)\,|\,\mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant Ce^{-A}(1+r^4)e^{-2\mu r}. \tag{8.20}$$

Equations (8.19) and (8.20) and the conditional Chebychev inequality now yield for large A and a,

$$\mathbf{P}(\widetilde{N}_t^{\mathrm{bulk}}(r) + \widetilde{N}_t^{\mathrm{fug}}(r) \geqslant e^{-2\delta/5 - r/3}N\,|\,\mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant C_\delta(1+r^4)e^{-A}e^{-\frac{4}{3}\mu r}.$$

This, together with Lemma 8.7 and the fact that the hat-, fug- and check-particles do not hit a on G_{nbab} finishes the proof of the lemma. □

We finish this section with a result which fills the gap in Lemma 8.4 under the hypothesis $(\mathrm{HB}^\flat_\perp)$.

Lemma 8.8. *Suppose* $(HB^\flat_\perp)$. *Then* $\mathbf{P}(|e^{-A}Z_0 - 1| \leqslant \varepsilon^{3/2},\ Y_0 \leqslant \eta) \leqslant o(1)$. *Moreover, let* $\alpha > 0$. *Then for large* A *and* a, *we have for every* $t \leqslant Ka^2$ *and* $0 \leqslant r \leqslant 9a/10$,

$$\mathbf{P}(N_t(r) \geqslant e^{-(\delta/2 + r/3)}N,\ R_{Ka^2} = 0) \leqslant C_{\alpha,\delta}e^{-(\frac{4}{3}-\alpha)r}/a, \qquad \textit{and} \qquad \mathbf{P}(R_{Ka^2} = 0) \geqslant 1 - C_\delta e^A/a.$$

Proof. Recall that initially, there are $\lfloor e^{-\delta}N\rfloor$ particles distributed independently according to the density $\phi(x)$ proportional to $\sin(\pi x/a)e^{-\mu x}\mathbf{1}_{(0,a)}(x)$. An elementary calculation yields that

$$\mathbf{E}[Z_0] = e^A(1+o(1)), \quad \mathrm{Var}(Z_0) \leqslant Ce^A/a^3, \quad \mathbf{E}[Y_0] \leqslant Ce^A/a. \tag{8.21}$$

This immediately yields the first statement, by Chebychev's and Markov's inequalities. Moreover, (8.21) with Lemmas 5.6 and 5.7 yields

$$\mathbf{P}(R_{Ka^2} \geqslant 1) \leqslant \mathbf{E}[R_{Ka^2}] \leqslant C\Big(\frac{K}{a}\mathbf{E}[Z_0] + \mathbf{E}[Y_0]\Big) \leqslant C\frac{Ke^A}{a}, \tag{8.22}$$

which gives the last statement. Note that on the event $\{R_{Ka^2} = 0\}$, the $\mathrm{B}^\flat$-BBM equals BBM with absorption at 0 and a. Since the density $\phi(x)$ is stationary w.r.t. this process, a quick calculation shows that

$$\mathbf{E}[N_t^{(0)}(r)] = \mathbf{E}[N_0^{(0)}(r)] = \lfloor e^{-\delta}N\rfloor\int_r^a \phi(x)\,\mathrm{d}x \leqslant e^{-\delta}N(1+\mu r)e^{-\mu r}. \tag{8.23}$$

Furthermore, by the independence of the initial particles and the law of total variance,

$$\mathrm{Var}(N_t^{(0)}(r)) = \lfloor e^{-\delta}N\rfloor\Big(\mathrm{Var}(\mathbf{E}^X[N_t^{(0)}(r)]) + \mathbf{E}[\mathrm{Var}^X(N_t^{(0)}(r))]\Big), \tag{8.24}$$

where X is a random variable distributed according to the density ϕ and the outer variance and expectation are with respect to X. By Lemma 5.5, we have for every $x \in [0,a]$,

$$\mathrm{Var}^x(N_t^{(0)}(r)) \leqslant \mathbf{E}^x[N_t^{(0)}(r)^2] \leqslant Ce^{-2A}N^2(1+r^4)e^{-2\mu r}((t/a^3)w_Z(x) + w_Y(x)),$$

and a simple calculation then yields for $t \leqslant Ka^2$,

$$\mathbf{E}[\mathrm{Var}^X(N_t^{(0)}(r))] = \int_0^a \mathrm{Var}^x(N_t^{(0)}(r))\phi(x)\,\mathrm{d}x \leqslant C(K/a)e^{-A}N(1+r^4)e^{-2\mu r}. \tag{8.25}$$

Moreover, by Lemma 5.4 and the hypothesis on r, we have for every $x \in [0, a]$,

$$\mathbf{E}^x[N_t^{(0)}(r)] \leqslant C(1+r^2)e^{-\mu r}\Big(e^{\mu x}\mathbf{1}_{(x \leqslant \frac{19a}{20})} + e^{-A}N(w_Z(x)+w_Y(x))\mathbf{1}_{(x > \frac{19a}{20})}\Big),$$

which yields

$$\begin{aligned}\mathbf{E}\Big[(\mathbf{E}^X[N_t^{(0)}(r)])^2\Big] &\leqslant C(1+r^4)e^{-2\mu r}\Big(\int_0^{\frac{19a}{20}} a e^{\mu x}\,\mathrm{d}x + e^{-2A}N^2\int_{\frac{19a}{20}}^a (a-x+1)^3 e^{\mu(x-2a)}\,\mathrm{d}x\Big)\\ &\leqslant Ca^{-3}e^{-A}N(1+r^4)e^{-2\mu r}.\end{aligned} \tag{8.26}$$

Equations (8.24), (8.25) and (8.26) now yield for large a,

$$\mathrm{Var}(N_t^{(0)}(r)) \leqslant C(K/a)e^{-A}N(1+r^4)e^{-2\mu r}. \tag{8.27}$$

The lemma now follows from (8.23) and (8.27), together with Chebychev's inequality. □

8.3 The probability of $G_1^\flat$

Most of the work in this section will be devoted to bounding the number of creations of red particles. For this, we will discretize time: We set $t_n = Ka^2 + n\delta/4$ and $I_n = [t_n, t_{n+1})$ for all $n \in \mathbf{N}$. Furthermore, define the $\mathscr{F}_{\mathscr{Y}}$-measurable random variables $n_1 := \max\{n : t_n < T\}$ and $n_2 := \max\{n : t_n < \Theta_1 + e^A/a^2\}$.

For an interval $I \subset \mathbf{R}_+$ and $r \in [0, a]$, we now define

$$U_I(r) = \#\{(u,t) \in \mathscr{L}_{\varnothing,I}^{\mathrm{red}} : X_u(t) \geqslant r\},$$

the number of red particles created to the right of r during the time interval I and set $U_n(r) = U_{I_n}(r)$. Furthermore, we denote by $N_t^{\mathrm{white}}(r)$ the number of white particles with positions $\geqslant r$ at time t, *including* the "in between particles".

Lemma 8.9. *Suppose (HB$_0$). Let $I = [t_l, t_r]$, with $d := t_r - t_l \leqslant \delta/4$. Then for every $\alpha > 0$, for sufficiently small δ, there exists $C_{\delta,\alpha}$, such that for large A and a, for every $r \in [0, 9a/10]$, if $Ka^2 \leqslant t_l < t_{6.21}$,*

$$\mathbf{E}[U_I(r)\,|\,T > t_l] \leqslant C_{\delta,\alpha}A^2\varepsilon\Big(\frac{t_l}{a^3}+\eta\Big)e^{-(\frac{4}{3}-\alpha)r}N, \tag{8.28}$$

$$\mathbf{P}(\exists t \in I : N_t^{\mathrm{white}}(r) \geqslant N\,|\,T > t_l) \leqslant C_{\delta,\alpha}A^2\varepsilon\Big(\frac{t_l}{a^3}+\eta\Big)e^{-(\frac{4}{3}-\alpha)r}, \tag{8.29}$$

$$\mathbf{E}[U_I(r)\mathbf{1}_{(R_{[t_l-\zeta,t_r]}=0)}\,|\,T > t_l] \leqslant C_{\delta,\alpha}dA^2\varepsilon\Big(\frac{t_l}{a^3}+\eta\Big)e^{-(\frac{4}{3}-\alpha)r}N. \tag{8.30}$$

Likewise, conditioned on $\mathscr{F}_\Delta$, we have for $T \leqslant t_l < \Theta_1 + e^A a^2$,

$$\mathbf{E}_{\mathrm{nbab}}[U_I(r)\,|\,\mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} \leqslant C_{\delta,\alpha}A^2\varepsilon^2 e^{-(\frac{4}{3}-\alpha)r}N, \tag{8.31}$$

$$\mathbf{P}_{\mathrm{nbab}}(\exists t \in I : N_t^{\mathrm{white}}(r) \geqslant N\,|\,\mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant C_{\delta,\alpha}A^2\varepsilon^2 e^{-(\frac{4}{3}-\alpha)r}. \tag{8.32}$$

Finally, suppose (HB$^\flat_\perp$). Then

$$\mathbf{E}[U_{[0,Ka^2]}(r)\mathbf{1}_{(R_{Ka^2}=0)}] \leqslant C_{\delta,\alpha}ae^{-(\frac{4}{3}-\alpha)r}N. \tag{8.33}$$

Proof. Morally, (8.28) follows from the fact that particles get created with rate $N/2$ and the probability that a particle at the point r turns red is given by the probability that there are more than N particles to the right of r, an upper bound of which is given by Lemma 8.4. If there were no in-between particles, we could make this argument precise and give a short proof for the expected amount of red particles created in an infinitesimal time interval. However, the presence of in-between particles, which do not follow Markovian dynamics due to the conditioning, makes the proof more complicated and forced us to discretise time.

Define the stopping time $\tau = \inf\{t \in I : N_t^{\mathrm{white}}(r) \geqslant N\}$, with $\inf \varnothing = +\infty$. Define further the law $\widehat{\mathbf{P}} = \mathbf{P}(\cdot \,|\, T > t_l)$ and the event $E = \{N_{t_l}(r') < e^{-\delta/2 - r'/3}N\}$, where $r' = r - \sqrt{r} \vee 0$. Then $e^{-4r'/3} \leqslant C_\alpha e^{-r(4/3-\alpha)}$ for every $\alpha > 0$, such that by Lemma 8.4, we have

$$\widehat{\mathbf{P}}(\tau < \infty) \leqslant \widehat{\mathbf{P}}(\tau < \infty,\, E) + \widehat{\mathbf{P}}(E^c) \leqslant \widehat{\mathbf{P}}(\tau < \infty \,|\, E) + C_{\delta,\alpha} A^2 \varepsilon \Big(\frac{t_l}{a^3} + \eta\Big) e^{-(\frac{4}{3}-\alpha)r}. \tag{8.34}$$

For $t \in I$, we can bound $N_t^{\mathrm{white}}(r)$ by the sum of

1. $N_t^{[1]}$: the number of descendants at time t of {the particles in $\mathscr{N}_{t_l}$ which are to the *right* of r'},
2. $N^{[2]}$: the number of descendants of {the particles in $\mathscr{N}_{t_l}$ which are to the *left* of r'} and which reach the point r before the time t_r and
3. $N_t^{[3]}$: the number of "in between" particles at time t.

Conditioned on $\mathscr{F}_{t_l}$ and the event E, $N_t^{[1]}$ is stochastically bounded by the number of individuals at time $t - t_l$ in a Galton–Watson process with reproduction law q and branching rate β_0, starting from $\lfloor e^{-\delta/2 - r'/3}N \rfloor$ individuals. Let $(\mathrm{GW}_t)_{t \geqslant 0}$ be such a process, then $e^{-\beta_0 m t}\mathrm{GW}_t = e^{-t/2}\mathrm{GW}_t$ is a martingale and by Doob's L^2-inequality, we get

$$E[\sup_{t\in[0,d]} (e^{-t/2}\mathrm{GW}_t - \mathrm{GW}_0)^2] \leqslant 4 \operatorname{Var}(e^{-d/2}\mathrm{GW}_d) \leqslant CdNe^{-r'/3}. \tag{8.35}$$

By (8.35), Chebychev's inequality and the hypothesis on d, we get,

$$\begin{aligned}\widehat{\mathbf{P}}(\sup_{t\in I} N_t^{[1]} \geqslant Ne^{-\delta/4}) &\leqslant P(\sup_{t\in[0,d]} \mathrm{GW}_t \geqslant Ne^{-\delta/4}) \\ &\leqslant P(\sup_{t\in[0,d]} e^{-t/2}\mathrm{GW}_t - \mathrm{GW}_0 \geqslant Ne^{-3\delta/8}(1 - e^{-\delta/8})) \leqslant C_\delta d e^{-r'/3}/N. \end{aligned}\tag{8.36}$$

Now, if $r < 1$, we have $N^{[2]} = 0$, so suppose that $r \geqslant 1$. Let the random variable Γ_r denote the number of particles in BBM which reach r before the time d. Since there are at most N white particles to the left of r' at time t_l, we have by Lemma 3.1,

$$\widehat{\mathbf{E}}[N^{[2]} \,|\, \mathscr{F}_{t_l}] \leqslant N \sup_{x\in[0,r']} \mathbf{E}^x[\Gamma_r] \leqslant CNW(\sup_{t\in[0,d]} B_t \geqslant \sqrt{r}) \leqslant CNd^{-1/2}e^{-r/(2d)}, \tag{8.37}$$

where (B_t) follows standard Brownian motion under W. For each $(u,t) \in \mathscr{N}_{t_l}$, let $N_u^{[2]}$ be the contribution of the descendants of u to $N^{[2]}$. By Lemma 3.5,

$$\begin{aligned}\operatorname{Var}[N^{[2]} \,|\, \mathscr{F}_{t_l}] &\leqslant \sum_{(u,t)\in\mathscr{N}_{t_l}} \mathbf{E}[(N_u^{[2]})^2] \leqslant N \sup_{x\in[0,r']} \mathbf{E}^x[\Gamma_r^2] \\ &\leqslant N\Big(\sup_{x\in[0,r']} \mathbf{E}^x[\Gamma_r] + C\int_0^d \int_0^a \mathbf{p}_t(x,z)\mathbf{E}^{(z,t)}[\Gamma_r]^2 \,\mathrm{d}z\,\mathrm{d}t\Big).\end{aligned}\tag{8.38}$$

Trivially, $\mathbf{E}^x[\Gamma_r] \leqslant e^{d/2}$ and $\int_0^a \mathbf{p}_t(x,z)\,\mathrm{d}z \leqslant e^{d/2}$ for any $x \in \mathbf{R}$, $t \leqslant d$. Together with (8.37) and (8.38), this yields,

$$\operatorname{Var}[N^{[2]} \,|\, \mathscr{F}_{t_l}] \leqslant CN \sup_{x\in[0,r']} \mathbf{E}^x[\Gamma_r] \leqslant CNd^{-1/2}e^{-r/(2d)}. \tag{8.39}$$

Chebychev's inequality and the inequality $t^{-1/2}e^{-1/(4t)} \leqslant Ct$ now yields

$$\mathbf{P}(N^{[2]} \geqslant N(1-e^{-\delta/8}) \,|\, \mathscr{F}_{t_l}) \leqslant C_\delta d e^{-r/(4d)}/N. \tag{8.40}$$

As for $N^{[3]}$, we have $\widehat{\mathbf{E}}[R_{t_l}^2] = \widehat{\mathrm{Var}}(R_{t_2}) + \widehat{\mathbf{E}}[R_{t_l}]^2 \leqslant Ce^{2A}$ by Lemma 6.12 as well as Lemmas 5.6, 6.8 and Corollary 6.7, such that by the definition of a breakout event, $\widehat{\mathbf{E}}[\sup_{t\in I}(N_t^{[3]})^2] \leqslant Ce^{2A}\zeta^2$. Markov's inequality then gives

$$\widehat{\mathbf{P}}\big(\sup_{t\in I} N_t^{[3]} \geqslant (e^{-\delta/4} - e^{-\delta/8})N\big) \leqslant C_\delta e^{2A}\zeta^2/N^2. \tag{8.41}$$

Equations (8.34), (8.36), (8.40) and (8.41) now yield (8.29). As for (8.28) and (8.30), we first note that the expected number of individuals created until time t in a Galton–Watson process with reproduction law $(q(k))_{k\geqslant 0}$ is bounded by the number of individuals at time t of a Galton–Watson process with reproduction law $(1-q(0))^{-1}q(k)_{k\geqslant 1}$, whose expectation is bounded by Ct. This gives,

$$\widehat{\mathbf{E}}[U_{(\tau,t_r]} \,|\, \tau < \infty] \leqslant C(dN + \zeta e^A). \tag{8.42}$$

because there are at most N white particles at the time t_l and the total number of in-between particles is bounded by $\zeta\mathbf{E}[R_{t_l}] \leqslant C\zeta e^A$ by Lemmas 5.6 and 6.8 and Corollary 6.7. As for the overshoot, i.e. the particles that have been coloured red at the time τ, we first note that

$$\widehat{\mathbf{E}}[U_{\{\tau\}}\mathbf{1}_{(\tau<\infty)}] \leqslant \widehat{\mathbf{E}}[U_{\{\tau\}}\mathbf{1}_{(\tau<\infty,\, U_{\{\tau\}}\geqslant N)}] + N\widehat{\mathbf{P}}(\tau < \infty). \tag{8.43}$$

Now,

$$U_{\{\tau\}}\mathbf{1}_{(\tau<\infty,\, U_{\{\tau\}}\geqslant N)} \leqslant \max\{k_u : u \in U, d_u \in I,\, k_u \geqslant N,\, X_u(d_u-) \geqslant r\} \tag{8.44}$$

By the definition of a breakout event, the set on the right-hand side of (8.44) contains no "in between" particles for $N > \zeta$. Furthermore, the expected number of branching events of "regular" particles to the right of r is bounded by

$$C\int_{t_l}^{t_r} \widehat{\mathbf{E}}[N_t(r)]\,\mathrm{d}t \leqslant CdN(1+r^2)e^{-r}, \tag{8.45}$$

where the inequality follows from (8.13). This yields,

$$\widehat{\mathbf{E}}[U_{\{\tau\}}\mathbf{1}_{(\tau<\infty,\, U_{\{\tau\}}\geqslant N)}] \leqslant CdN(1+r^2)e^{-r}E[L\mathbf{1}_{(L\geqslant N)}] \leqslant Cd(1+r^2)e^{-r}, \tag{8.46}$$

where the last inequality follows from the Cauchy–Schwarz and Chebychev inqualities. By (8.29), (8.42), (8.43) and (8.46) we finally get for large A and a,

$$\widehat{\mathbf{E}}[U_I(r)] = \widehat{\mathbf{E}}[(U_{\{\tau\}}(r) + U_{(\tau,t_r]}(r))\mathbf{1}_{(\tau<\infty)}] \leqslant C_{\delta,\alpha}NA^2\varepsilon\Big(\frac{t_l}{a^3} + \eta\Big)e^{-(\frac{4}{3}-\alpha)r}. \tag{8.47}$$

For the proof of (8.30), we note that on the set $R_{[t_l-\zeta,t_r]} = 0$ there are no in-between particles during I. Taking away the corresponding terms in the above proof yields (8.30).

The proof of the other three equations is very similar. The proof of (8.31) and (8.32) uses the second half of Lemma 8.4 instead of the first and (8.12) in addition to (8.13) for the proof of (8.45). The proof of the last equation draws on Lemma 8.8 instead of Lemma 8.4 and requires covering the interval $[0, Ka^2]$ by pieces of length d. □

Corollary 8.10. *Suppose (HB$_0$). For an interval $I \subset \mathbf{R}_+$, let $G_{a/2,I}$ be the event that no particle is coloured red during the time interval I and to the right of $a/2$. Then,*

$$\mathbf{P}(G_{\mathscr{U}}\backslash G_{a/2,[Ka^2,T]}) \leqslant C_\delta o(1) \quad \textit{and} \quad \mathbf{P}_{\mathrm{nbab}}\Big((G_{a/2,[T,\Theta_1+e^Aa^2]})^c \,|\, \mathscr{F}_\Delta\Big)\,\mathbf{1}_{G_\Delta} \leqslant C_\delta o(1).$$

Proof. Follows immediately from (8.29) and (8.32) by summing over the intervals I_n (note that $T \leqslant \sqrt{\varepsilon} a^3$ on $G_{\mathscr{Y}}$). $\square$

Lemma 8.11. *Suppose (HB$_0$). Set $T' = T \wedge \sqrt{\varepsilon} a^3$ and define the intervals $I = [Ka^2, T']$ and $J = [T, \Theta_1 + e^A a^2]$ For large A and a,*

$$\mathbf{E}[(Z^{\mathrm{red}}_{\varnothing,I} + Y^{\mathrm{red}}_{\varnothing,I})\mathbf{1}_{G_{a/2,I}}] \leqslant C_\delta A^2 \varepsilon^3 e^A, \qquad \textit{and} \qquad \mathbf{E}_{\mathrm{nbab}}[(Z^{\mathrm{red}}_{\varnothing,J} + Y^{\mathrm{red}}_{\varnothing,J})\mathbf{1}_{G_{a/2,J}} \,|\, \mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} \leqslant C_\delta e^{2A}/a.$$

Now suppose (HB$^\flat_\perp$). Then $\mathbf{E}[(Z^{\mathrm{red}}_{\varnothing,[0,Ka^2]} + Y^{\mathrm{red}}_{\varnothing,[0,Ka^2]})\mathbf{1}_{(R_{Ka^2}=0)}] \leqslant C_\delta e^A/a^2$.

Proof. By Lemma 8.9, we have for every $r \in [0, a/2]$ and $n \leqslant n_0 := \lfloor (\sqrt{\varepsilon} a^3 - Ka^2)/(t_1 - t_0) \rfloor$,

$$\begin{aligned}\mathbf{E}[U_I(r)] &\leqslant \sum_{n=0}^{n_0} \mathbf{E}[U_n(r) \,|\, T > t_n]\mathbf{P}(T > t_n) \leqslant C_\delta A^2 \varepsilon e^{-\frac{8}{7}r} N \sum_{n=0}^{n_0} \Big(\frac{t_n}{a^3} + \eta\Big)\mathbf{P}(T > t_n) \\ &\leqslant C_\delta A^2 \varepsilon e^{-\frac{8}{7}r} e^A \mathbf{E}\Big[\Big(\frac{T \wedge \sqrt{\varepsilon} a^3}{a^3} + \eta\Big)^2\Big] \leqslant C_\delta A^2 \varepsilon^3 e^{-\frac{8}{7}r},\end{aligned} \tag{8.48}$$

by Lemma 6.5 and (6.3). Now, by integration by parts, we have for every (possibly random) interval I,

$$Z^{\mathrm{red}}_{\varnothing,I} = \int_0^a w_Z'(r) U_I(r)\,\mathrm{d}r \qquad \text{and} \qquad Y^{\mathrm{red}}_{\varnothing,I} = U_I(0) + \int_0^a w_Y'(r) U_I(r)\,\mathrm{d}r. \tag{8.49}$$

The first equation in the statement of the lemma now follows from (8.48) and (8.49), since $w_Z'(x) \leqslant C(1+x)e^{x-a}$ and $w_Y'(x) \leqslant Ce^{x-a}$ for $x \in [0, a]$. The remaining equations follow similarly. $\square$

The following lemma will only be needed in the proof of Proposition 8.3.

Lemma 8.12. *Suppose (HB$^\flat_0$). Let $r \in [0, 9a/10]$ and $(K+1)a^2 \leqslant t \leqslant t_{6.21}$. Then, for all $\alpha > 0$, for large A and a,*

$$\mathbf{P}[N_t^{\mathrm{red},(0)}(r) > \alpha N,\, T > t] \leqslant C_\delta\Big(\alpha^{-1}(1+\mu r)e^{-\mu r} A^2 \varepsilon \Big(\frac{t}{a^3} + \eta\Big) + \varepsilon^2\Big).$$

Proof. For an interval I, denote by $N_t^{\mathrm{red},(0)}(r, I)$ the number of red particles to the right of r at time t which have turned red during I. Define the event

$$G = G_{a/2,[0,t]} \cap \{R_{[t-a^2-\zeta,t]} = 0\} \cap \{Z^{\mathrm{red}}_{\varnothing,[0,Ka^2]} + Y^{\mathrm{red}}_{\varnothing,[0,Ka^2]} \leqslant a^{-1/2}\},$$

and note that $\mathbf{P}(G^c, T > t) \leqslant C\varepsilon^2$ for large A and a, by Corollary 8.10, Lemma 6.9 and the hypothesis.

Write $\widehat{\mathbf{E}} = \mathbf{E}[\cdot \,|\, T > t]$. If $\widehat{\mathbf{E}}^{(x,s)}[N_t^{(0)}(r)] \leqslant f(x,s)$ for some function f with $f(0,s) \equiv f(a,s) \equiv 0$, then by integration by parts,

$$\widehat{\mathbf{E}}[N_t^{\mathrm{red},(0)}(r, I)\mathbf{1}_G] \leqslant \int_I \int_0^{a/2} \widehat{\mathbf{E}}[U_{\mathrm{d}s}(x)\mathbf{1}_{(R_{[t-\zeta,t]}=0)}]\frac{\mathrm{d}}{\mathrm{d}x} f(x,s)\,\mathrm{d}x\,\mathrm{d}s.$$

Define $I_1 = [0, t-a^2]$ and $I_2 = [t-a^2, t]$. By (5.20), (5.21) and Corollary 6.7, we have for $x \in [0, a/2]$,

$$\widehat{\mathbf{E}}^{(x,s)}[N_t^{(0)}(r)] \leqslant \begin{cases} C(1+\mu r)e^{-\mu r} N e^{-A} w_Z(x), & \text{if } s \in I_1, \\ Ce^{\mu x} \int_r^\infty e^{-\mu z} p^a_{t-s}(x,z), & \text{if } s \in I_2, \end{cases}$$

Similarly to the proof of Lemma 8.11, we now have by the inequality $w_Z'(x) \leqslant C(1+x)e^{x-a}$, for large A and a,

$$\begin{aligned}\widehat{\mathbf{E}}[N_t^{\mathrm{red},(0)}(r, I_1)\mathbf{1}_G] &\leqslant Ce^{-A}N(1+\mu r)e^{-\mu r}\int_0^{a/2}\widehat{\mathbf{E}}[U_{I_1}(x)\mathbf{1}_G](1+x)e^{x-a}\,\mathrm{d}x\\ &\leqslant C_\delta A^2\varepsilon((t/a^3)+\eta)(t/a^3)(1+\mu r)e^{-\mu r}N,\end{aligned}$$

by Lemma 8.9 and the definition of G. As for the particles created during I_2, we have by Lemma 8.9, the definition of G and the hypothesis on t, for all $s \in I_2$,

$$\widehat{\mathbf{E}}[U_{\mathrm{d}s}(x)\mathbf{1}_G] \leqslant C_\delta A^2\varepsilon\Big(\frac{t}{a^3}+\eta\Big)Ne^{-\frac{5}{4}x}\mathrm{d}s.$$

Now, with (2.5), one easily sees that for large a, $(\mathrm{d}/\mathrm{d}x)(e^{\mu x}p_s^a(x,z)) \geqslant 0$ for all $x \leqslant a/2$ and $s \geqslant a^2$, whence, by (2.8) and Fubini's theorem, we get

$$\begin{aligned}\widehat{\mathbf{E}}[N_t^{\mathrm{red},(0)}(r, I_2)\mathbf{1}_G] &\leqslant C\int_0^{a^2}\int_0^{a/2}\widehat{\mathbf{E}}[U_{\mathrm{d}s}(x)\mathbf{1}_G]\int_r^\infty e^{-\mu z}\frac{\mathrm{d}}{\mathrm{d}x}\left(e^{\mu x}p_s^a(x,z)\right)\,\mathrm{d}z\,\mathrm{d}x\,\mathrm{d}s\\ &\leqslant C\int_0^{a/2}\widehat{\mathbf{E}}[U_{\mathrm{d}s}(x)\mathbf{1}_G]\int_r^\infty e^{-\mu z}\frac{\mathrm{d}}{\mathrm{d}x}\big(e^{\mu x}a^{-1}(x\wedge z)(a-x\vee z)\big)\,\mathrm{d}z\,\mathrm{d}x\\ &\leqslant C_\delta A^2\varepsilon\Big(\frac{t}{a^3}+\eta\Big)N\int_r^\infty\int_0^{a/2}e^{-\mu(x/4+z)}(1+x)\,\mathrm{d}x\,\mathrm{d}z\\ &\leqslant C_\delta A^2\varepsilon((t/a^3)+\eta)e^{-\mu r}N.\end{aligned}$$

In total, we have

$$\widehat{\mathbf{E}}[N_t^{\mathrm{red},(0)}(r)\mathbf{1}_G] \leqslant C_\delta A^2\varepsilon\Big(\frac{t}{a^3}+\eta\Big)(1+\mu r)e^{-\mu r}N.$$

An application of Markov's inequality finishes the proof. □

Lemma 8.13. *Suppose ($HB_0^\flat$). Let G_{fred} be the event that the fugitive does not get coloured red. Then, for large A and a,*

$$\mathbf{P}(G_{\mathscr{U}} \cap G_{a/2,[Ka^2,T]}\backslash G_{\mathrm{fred}}) \leqslant C_\delta\varepsilon^2.$$

Proof. Set $G_{a/2} = G_{a/2,[Ka^2,T]}$. Let $\mathbf{P}_0$ be the law of BBM with absorption at zero from Section 6 (i.e. we do not move the barrier when a breakout occurs). Let T_{red} be the first breakout of a red particle. Then

$$\mathbf{P}(G_{\mathscr{U}} \cap G_{a/2}\backslash G_{\mathrm{fred}}) = \mathbf{P}_0(T = T_{\mathrm{red}}, \mathscr{G}_{\mathscr{U}} \cap G_{a/2}) \leqslant \mathbf{P}_0(T_{\mathrm{red}} \leqslant \sqrt{\varepsilon}a^3, G_{a/2}). \tag{8.50}$$

Define $T' = T \wedge \sqrt{\varepsilon}a^3$. Then, since $\mathscr{L}_{\varnothing,T'}^{\mathrm{red}}$ is a stopping line, we have by Proposition 6.2 and Corollary 8.10,

$$\mathbf{P}_0(T_{\mathrm{red}} \leqslant \sqrt{\varepsilon}a^3 \,|\, \mathscr{F}_{\mathscr{L}_{\varnothing,T'}^{\mathrm{red}}})\mathbf{1}_{G_{a/2}} \leqslant C\varepsilon^{-1/2}e^{-A}(Z_{\varnothing,T'}^{\mathrm{red}} + Y_{\varnothing,T'}^{\mathrm{red}}). \tag{8.51}$$

Recall that $\mathbf{P}(Z_{\varnothing,[0,Ka^2]}^{\mathrm{red}} + Y_{\varnothing,[0,Ka^2]}^{\mathrm{red}} \leqslant a^{-1/2}) \geqslant 1-\varepsilon^2$ by hypothesis. By the tower property of conditional expectation and (8.51), this gives

$$\begin{aligned}\mathbf{P}_0(T_{\mathrm{red}} \leqslant \sqrt{\varepsilon}a^3, G_{a/2}) &\leqslant \mathbf{E}[\mathbf{P}_0(T_{\mathrm{red}} \leqslant \sqrt{\varepsilon}a^3 \,|\, \mathscr{F}_{\mathscr{L}_{\varnothing,T'}^{\mathrm{red}}})\mathbf{1}_{G_{a/2}}] + \varepsilon^2 + o(1)\\ &\leqslant C\mathbf{E}[C\varepsilon^{-1/2}e^{-A}(Z_{\varnothing,[Ka^2,T']}^{\mathrm{red}} + Y_{\varnothing,[Ka^2,T']}^{\mathrm{red}})\mathbf{1}_{G_{a/2}}] + C\varepsilon^2.\end{aligned}$$

The lemma now follows from (8.50) together with Lemma 8.11, (6.1) and the hypothesis. □

Lemma 8.14. *Suppose (HB$^\flat_0$). There exists a numerical constant $b > 0$, such that for large A and a,*

$$\mathbf{P}(G_1^\flat) \geqslant 1 - \varepsilon^{1+b}.$$

Proof. Recall the event G_1 from Section 7 and change its definition slightly by requiring that $|e^{-A}Z_{\Theta_1} - 1| \leqslant \varepsilon^{3/2}/2$, call the new event G_1'. Define the random variable

$$X = \mathbf{P}(Z^{\mathrm{red}}_{\varnothing,[\Theta_1,\Theta_1+Ka^2]} + Y^{\mathrm{red}}_{\varnothing,[\Theta_1,\Theta_1+Ka^2]} \leqslant a^{-1/2} \,|\, \mathscr{F}_{\Theta_1}).$$

By Proposition 7.3 and Remark 7.7, it suffices to show that

$$\mathbf{P}\left(G_1' \backslash \left\{Z^{\mathrm{red}}_{\Theta_1} \leqslant \varepsilon^{3/2}/2,\ X > 1-\varepsilon^2\right\}\right) \leqslant \varepsilon^{1+b},$$

for some numerical constant b.

Let T', I and J as in Lemma 8.11. We then have by Markov's inequality, Lemma 8.11 and Corollary 8.10,

$$\mathbf{E}_{\mathrm{nbab}}[1 - X \,|\, \mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} \leqslant C_\delta o(1). \tag{8.52}$$

Markov's inequality applied to (8.52) then yields

$$\mathbf{P}(X > 1-\varepsilon^2,\ G_\Delta \cap G_{\mathrm{nbab}}) \leqslant C_\delta o(1). \tag{8.53}$$

Now define $G_{\mathrm{red}} = \{T \leqslant \sqrt{\varepsilon}a^3,\ Z^{\mathrm{red}}_{\varnothing,[0,Ka^2]} + Y^{\mathrm{red}}_{\varnothing,[0,Ka^2]} \leqslant a^{-1/2}\} \cap G_{a/2,I} \cap G_{\mathrm{fred}}$. Conditioned on T and $\mathscr{F}_{\mathscr{L}^{\mathrm{red}}_{\varnothing,T}}$, on the set G_{red}, the particles from the stopping line $\mathscr{L}^{\mathrm{red}}_{\varnothing,T}$ then all spawn BBM conditioned not to break out before T (because neither the fugitive nor any in-between particles are on the stopping line). By Lemma 8.11, Lemma 6.9 and the tower property of conditional expectation, we then have

$$\mathbf{E}[Z^{\mathrm{red}}_T\mathbf{1}_{G_{\mathrm{red}}}] \leqslant C\Big(Aa^{-1/2} + \mathbf{E}[(Z^{\mathrm{red}}_{\varnothing,I} + AY^{\mathrm{red}}_{\varnothing,I})\mathbf{1}_{G_{a/2,I}}]\Big) \leqslant C_\delta(A\varepsilon)^3e^A$$

$$\mathbf{E}[Y^{\mathrm{red}}_T\mathbf{1}_{G_{\mathrm{red}}}] \leqslant C\Big(Aa^{-1/2} + \mathbf{E}[(Z^{\mathrm{red}}_{\varnothing,I} + Y^{\mathrm{red}}_{\varnothing,I})\mathbf{1}_{G_{a/2,I}}]\Big) \leqslant C_\delta A^2\varepsilon^3e^A.$$

By Lemma 6.9, we now have

$$\mathbf{E}_{\mathrm{nbab}}[Z^{\mathrm{red}}_{\Theta_1} \,|\, \mathscr{F}_\Delta \vee \mathscr{F}_T]\mathbf{1}_{G_\Delta} \leqslant C_\delta\Big(Z^{\mathrm{red}}_T + Z^{\mathrm{red}}_{\varnothing,J} + A(Y^{\mathrm{red}}_T + Y^{\mathrm{red}}_{\varnothing,J})\Big),$$

and Lemma 8.11 and the tower property of conditional expectation give

$$\mathbf{E}[Z^{\mathrm{red}}_{\Theta_1}\mathbf{1}_{G_\Delta\cap G_{\mathrm{nbab}}\cap G_{\mathrm{red}}}] \leqslant C_\delta(A\varepsilon)^3e^A. \tag{8.54}$$

Furthermore, by the hypothesis, Corollary 8.10 and Lemmas 7.4, 7.5 and 8.13, we have

$$\mathbf{P}(G_\Delta \cap G_{\mathrm{nbab}} \cap G_{\mathrm{red}}) \geqslant 1 - \varepsilon^{1+b}. \tag{8.55}$$

The lemma now follows from (8.53), (8.55) and Markov's inequality applied to (8.54), together with (6.1). □

8.4 Proofs of the main results

Proof of Lemma 8.1. By Lemma 8.8, it remains to estimate the probability of the last event in the definition of $G_0^\flat$. Define the random variable $X = \mathbf{P}(Z^{\mathrm{red}}_{\varnothing,[0,Ka^2]} + Y^{\mathrm{red}}_{\varnothing,[0,Ka^2]} \leqslant a^{-1/2} \,|\, \mathscr{F}_0)$. By Lemma 8.8 and Markov's inequality applied to the last equation of Lemma 8.11, we have $\mathbf{E}[(1-X)\mathbf{1}_{(R_{Ka^2}=0)}] \leqslant C_\delta e^A/a$. By Markov's inequality, we then have

$$\mathbf{P}(X \leqslant 1-\varepsilon^2,\ R_{Ka^2} = 0) \leqslant \varepsilon^{-2}\mathbf{E}[(1-X)\mathbf{1}_{(R_{Ka^2}=0)}] \leqslant C_\delta o(1).$$

Together with (8.22), this finishes the proof. □

Proof of Proposition 8.2. Let $n \geqslant 0$. Conditioned on $\mathscr{F}_{\Theta_n}$, the barrier process until Θ_{n+1} is by definition the same in B$^\flat$-BBM and B-BBM. Furthermore, $G_n \subset G_n^\flat$ and $G_n^\flat \in \mathscr{F}_{\Theta_n}$. The first statement then follows by induction from Lemma 8.14, as in the beginning of the proof of Proposition 7.3.

As for the second statement, inspection of the proofs of Theorems 7.1 and 7.2 shows that they only rely on Proposition 7.3 and on the existence of the coupling with a Poisson process constructed in Section 7.3. But this construction only relied on the law of T_{n+1} conditioned on $\mathscr{F}_{\Theta_n}$ and G_n and thus readily transfers to the B$^\flat$-BBM. $\square$

Proof of Proposition 8.3. Define $\rho = x_{\alpha e^{2\delta}}$, such that $(1+\mu\rho)e^{-\mu\rho} \geqslant \alpha e^{2\delta}(1+o(1))$. Recall from Section 7.3 the coupling of the process $(\Theta_n/a^3)_{n\geqslant 0}$ with a Poisson process $(V_n)_{n\geqslant 0}$ of intensity $\gamma_0^{-1} = p_B e^A \pi$. As indicated above, this coupling can be constructed for the B$^\flat$-BBM as well. In particular, setting $\widetilde{G}_n^\flat = G_n^\flat \cap \{\forall j \leqslant n : |(T_j - \Theta_{j-1})/a^3 - V_j| \leqslant \varepsilon^{3/2}\}$, we have as in (7.32), for every $n \geqslant 0$,

$$\mathbf{P}(\widetilde{G}_n^\flat) \geqslant \mathbf{P}(G_n^\flat) - nO(\varepsilon^2) \geqslant \mathbf{P}(G_0^\flat) - nO(\varepsilon^{1+b}), \tag{8.56}$$

by Proposition 8.2 (where $b > 0$ is a numerical constant). Now let $(t_i^{A,a})$ satisfy (Ht). For simplicity, suppose that $t_i^{A,a} \equiv t_i$ for all i, the proof of the general case is exactly the same. We want to show that $\mathbf{P}(\forall 1 \leqslant i \leqslant k : N_{t_i a^3}^{\text{white}}(\rho) \geqslant \alpha N) \to 1$ as A and a go to infinity. By (7.42), the probability that there exists (i,j), such that t_i and t_{i+1} are in the same interval $[\Theta_j, \Theta_{j+1}]$ is bounded by $O(\sqrt{\varepsilon})$ for large A and a. We can therefore suppose that $k = 1$, the general case is a straightforward extension.

It thus remains to show that for every $t > 0$, $\mathbf{P}(N_{ta^3}^{\text{white}}(\rho) \geqslant \alpha N) \to 1$ as A and a go to infinity (the case $t = 0$ is a straightforward calculation). Let $t > 0$ and $n_0 := \lceil \gamma_0^{-1}(t+1) \rceil$. Then $\mathbf{E}[V_{n_0}] \geqslant t+1$ and $\mathrm{Var}(V_{n_0}) \leqslant 2\gamma_0(t+1)$, such that $\mathbf{P}(V_{n_0} < t + 1/2) \leqslant C\gamma_0 = O(\varepsilon)$ (for this proof, we allow the constants C and the expression $O(\cdot)$ depend on t). Defining the event $E_n(s)$ for $n \geqslant 0$ by

$$E_n(s) = \{\Theta_n > s\} \cap \{\nexists j \in \{0,\dots,n\} : s \in [T_{j-1}, \Theta_j + e^A a^2]\} \cap \widetilde{G}_n^\flat,$$

(define $T_{-1} = 0$) we then have by (7.42) and (8.56), $\mathbf{P}(E_{n_0}) \geqslant 1 - O(\varepsilon^b)$.

We now prove by induction that there exists a numerical constant $b > 0$, such that for every $1 \leqslant n \leqslant n_0$,

$$S_n = \sup_{\nu \in G_0^\flat} \sup_{s \in [0, ta^3]} \mathbf{P}^\nu(N_s^{\text{white}}(\rho) < \alpha N,\, E_n(s)) \leqslant C_{\delta,\alpha} n \varepsilon^{1+b},$$

as A and a go to infinity (we accept the abuse of notation $\nu \in G_0^\flat$). By definition, $S_0 = 0$. Now, suppose the statement is true for some $n \geqslant 0$. We then have for every $\nu \in G_0^\flat$ and $s \in [0, ta^3]$,

$$\begin{aligned}\mathbf{P}^\nu(N_s^{\text{white}}(\rho) < \alpha N,\, E_{n+1}(s)) &\leqslant \mathbf{P}^\nu(N_s^{\text{white}}(\rho) < \alpha N,\, E_{n+1}(s),\, s > \Theta_1)\\ &\quad + \mathbf{P}^\nu(N_s^{\text{white}}(\rho) < \alpha N,\, E_{n+1}(s),\, s \leqslant \Theta_1) =: P_1 + P_2.\end{aligned}$$

We first have, by the definition of $E_{n+1}(s)$ and the fact that the process starts afresh at the stopping time Θ_1,

$$P_1 \leqslant \mathbf{E}^\nu[\mathbf{P}(N_s^{\text{white}}(\rho) < \alpha N,\, E_{n+1}(s) \,|\, \mathscr{F}_{\Theta_1}) \mathbf{1}_{G_1^\flat \cap \{\Theta_1 \leqslant s\}}] \leqslant S_n.$$

Second, we have again by the definition of $E_{n+1}(s)$,

$$P_2 \leqslant \mathbf{P}^\nu(N_s^{\text{white}}(\rho) < \alpha N,\, T > s)\mathbf{1}_{(s \geqslant e^A a^2)} \leqslant \mathbf{P}^\nu(N_s^{\text{white},(0)}(\rho) < \alpha N,\, T > s)\mathbf{1}_{(s \geqslant e^A a^2)}. \tag{8.57}$$

Write $\widehat{\mathbf{P}}^\nu = \mathbf{P}^\nu(\cdot \,|\, T > s)$. For $e^A a^2 \leqslant s \leqslant t_{6.21}$, we now have by Proposition 5.3 and Corollary 6.7 for the first inequality and Lemma 5.5 and Corollary 6.7 for the second,

$$\begin{aligned}&\widehat{\mathbf{E}}^\nu[N_s^{(0)}(\rho)] \geqslant (1 - O(\varepsilon^{3/2}))e^{2\delta}\alpha e^{-\delta} N \geqslant (1 - O(\varepsilon^{3/2}))(1+\delta)\alpha,\\ &\widehat{\mathrm{Var}}^\nu(N_s^{(0)}(\rho)) \leqslant C_\alpha \alpha^2 e^{-2A} N^2,\end{aligned}$$

such that

$$\begin{aligned}\mathbf{P}^\nu(N_s^{\mathrm{white},(0)}(\rho) < \alpha N,\, T > s) &\leqslant \widehat{\mathbf{P}}^\nu(N_s^{(0)}(\rho) < (1+\delta/2)\alpha N) + \mathbf{P}^\nu(N_s^{\mathrm{red},(0)}(\rho) \geqslant (\delta/2)\alpha N,\, T > s)\\ &\leqslant C_{\delta,\alpha}\left(e^{-2A} + A^2\varepsilon(s/a^3+\eta)\right).\end{aligned}$$

by Chebychev's inequality applied to the first term and Lemma 8.12 to the second. Altogether, this gives,

$$P_2 \leqslant C_{\delta,\alpha}\Big(\mathbf{P}(T > \sqrt{\varepsilon}a^3) + e^{-2A} + A^2\varepsilon^{3/2}\Big) \leqslant C_{\delta,\alpha}A^2\varepsilon^{3/2},$$

for large A and a. Finally, we get $S_{n+1} \leqslant S_n + C_{\delta,\alpha}A^2\varepsilon^{3/2}$ for all $0 \leqslant n \leqslant n_0$, which yields $S_n \leqslant C_{\delta,\alpha}n\varepsilon^{5/4}$ for large A and a, by (6.1). This gives

$$\mathbf{P}(N_{ta^3}^{\mathrm{white}}(\rho) \leqslant \alpha N) \leqslant S_{n_0} + \mathbf{P}(V_{n_0} < t + 1/2) + (1 - \mathbf{P}(E_{n_0}(s))) \to 0,$$

as A and a go to infinity. The statement follows. □

9 The B$^\sharp$-BBM

In this section, we define and study the B$^\sharp$-BBM, a model which will be used in Section 10 to bound the N-BBM from *above.*

9.1 Definition of the model

As in the previous section, we will use throughout the notation from Section 7, fix $\delta \in (0,1/2)$ and define K to be the smallest number, such that $K \leqslant 1$ and $E_K \leqslant \delta/10$. We now define however $N = \lfloor 2\pi e^{A-\delta}a^{-3}e^{\mu a}\rfloor$. Again, we will use interchangeably the phrases "as A and a go to infinity" and "as N go to infinity" and A_0 and the function $a_0(A)$ in the definition of these phrases may now also depend on δ. The symbols C_δ and $C_{\delta,\alpha}$ have the same meaning as in the last section.

The B$^\sharp$-BBM is then defined as follows: Given a possibly random initial configuration ν_0 of particles in $(0,a)$, we let particles evolve according to B-BBM with barrier function given by (7.1) and with the following changes: Define $t_n = n(K+3)a^2$ and $I_n = [t_n, t_{n+1})$ (note that these definitions differ from those of Section 8). Colour all initial particles white. When a white particle hits 0 during the time interval I_n and has at least N particles to its right, it is killed immediately. If less than N particles are to its right, it is coloured blue and survives until the time $t_{n+2} \wedge \Theta_1$, where all of its descendants *to the left of* 0 are killed and the remaining survive and are coloured white again. At the time Θ_1, the process starts afresh. See Figure 4 for a graphical description.

For bookkeeping, we add a shade of grey to the white particles which have hit 0 at least once (and call them hence the grey particles). We then add the superscripts "nw", "gr", "blue" or "tot" to the quantities referring respectively to the non-white, grey, blue or all the particles. Quantities without this superscript refer to the white particles.

In particular, we define B_I and B_I^{tot} to be the number of white, respectively, white and grey particles touching the left barrier during the time interval I with less than N particles to their right (i.e. those which are coloured blue). We set $B_n = B_{I_n}$ and $B_n^{\mathrm{tot}} = B_{I_n}^{\mathrm{tot}}$.

Let $\nu_t^\sharp$ be the configuration of (all) the particles at the time t and abuse notation by setting $\nu_n^\sharp = \nu_{\Theta_n}^\sharp$. We set $G_{-1}^\sharp = \Omega$ and for each $n \in \mathbf{N}$, we define the event $G_n^\sharp$ to be the intersection of $G_{n-1}^\sharp$ with the following events (we omit the braces).

- $\operatorname{supp}\nu_n^\sharp \subset (0,a)$,
- $\mathscr{N}_{\Theta_n}^{\mathrm{tot}} \subset U \times \{\Theta_n\}$ and $\Theta_n > T_n^+$ (for $n > 0$),
- $|e^{-A}Z_{\Theta_n}^{\mathrm{tot}} - 1| \leqslant \varepsilon^{3/2}$ and $Y_{\Theta_n}^{\mathrm{tot}} \leqslant \eta$,



Figure 4: The B$^\sharp$-BBM with parameter $N = 3$ (no breakout is shown). White, blue and grey particles are drawn with solid, dotted and dashed lines, respectively. A cross indicates that a blue particle is killed.

- $N(\Theta_n) \geqslant N$ and for all $j \geqslant 0$ with $\Theta_{n-1} + t_j < \Theta_n$: $N(\Theta_{n-1} + t_j) \geqslant N$,
- $\mathbf{P}\Big(B^{\mathrm{tot}}_{[\Theta_n,\Theta_n+t_1]} \leqslant e^{-A}e^{\mu a}/a \,\Big|\, \mathscr{F}_{\Theta_n}\Big) \geqslant 1 - \varepsilon^2$.

The last event is of course uniquely defined up to a set of probability zero. Note that $G^\sharp_n \in \mathscr{F}_{\Theta_n}$ for each $n \in \mathbf{N}$. Furthermore, we define the predicates

(HB$^\sharp$) (The law of) ν_0 is such that $\mathbf{P}(G^\sharp_0) \to 1$ as A and a go to infinity.
(HB$^\sharp_0$) ν_0 is deterministic and such that $G^\sharp_0$ holds.
(HB$^\sharp_\perp$) ν_0 is obtained from $\lceil e^\delta N\rceil$ particles distributed independently according to the density proportional to $\sin(\pi x/a)e^{-\mu x}\mathbf{1}_{(0,a)}(x)$.

We now state the important results on the B$^\sharp$-BBM.

Lemma 9.1. *(HB$^\sharp_\perp$) implies (HB$^\sharp$) for large A and a.*

Proposition 9.2. *Proposition 7.3 still holds for the B$^\sharp$-BBM, with G_n replaced by $G^\sharp_n$. The same is true for Theorems 7.1 and 7.2, with (HB) replaced by (HB$^\sharp$).*

Recall the definition of qu^N_α and x_α from the introduction and of (Ht) from Section 7.

Proposition 9.3. *Suppose (HB$^\sharp_\perp$). Let $(t^{A,a}_j)$ satisfy (Ht). Let $\alpha \in (0,1)$. As A and a go to infinity,*

$$\mathbf{P}(\forall j : \mathrm{qu}^N_\alpha(\nu^\sharp_{a^3 t^{A,a}_j}) \leqslant x_{\alpha e^{-2\delta}}) \to 1.$$

Lemma 9.4. *Define a variant called C$^\sharp$-BBM of the B$^\sharp$-BBM by killing blue particles only if there are at least N particles to their right. Then Propositions 9.2 and 9.3 hold for the C$^\sharp$-BBM as well.*

9.2 Preparatory lemmas

We first derive upper bounds on the probability that the number of white particles is less than N at a given time t. We do not impose any particular condition on the initial configuration.

Lemma 9.5. *Let $\widehat{\mathbf{P}} = \mathbf{P}(\cdot\,|\,T > s)$ with $s \leqslant t_{6.21}$ and let $Ka^2 \leqslant t \leqslant s$. We have for small δ and for A and a large enough,*

$$\widehat{\mathbf{P}}(N_t^{(0)} < N \,|\, \mathscr{F}_0)\mathbf{1}_{(Z_0 \geqslant (1-\delta/2)e^A)} \leqslant C_\delta e^{-A}(a^{-1} + e^{-A}Y_0).$$

Proof. By (5.20), Corollary 6.7 and the definition of K, we have for A and a large enough,

$$\widehat{\mathbf{E}}[N_t^{(0)} \,|\, \mathscr{F}_0] \geqslant (1 - Cp_B)\mathbf{E}[N_t^{(0)} \,|\, \mathscr{F}_0] \geqslant (1 + \delta/4)Ne^{-A}Z_0.$$

By the conditional Chebychev inequality, we then have

$$\widehat{\mathbf{P}}(N_t^{(0)} < N \,|\, \mathscr{F}_0)\mathbf{1}_{(Z_0 \geqslant (1-\delta/2)e^A)} \leqslant C\frac{\widehat{\mathrm{Var}}(N_t^{(0)} \,|\, \mathscr{F}_0)}{(e^{-A}N\delta Z_0)^2}.$$

The lemma now follows from Lemma 5.5 and Corollary 6.7. ☐

Corollary 9.6. *Under the conditions of Lemma 9.5, suppose furthermore that $t \geqslant (K+1)a^2$. Then, for small δ and for A and a large enough,*

$$\widehat{\mathbf{P}}(N_t^{(0)} < N,\, Z_{t-Ka^2}^{(0)} \geqslant (1-\delta/2)e^A \,|\, \mathscr{F}_0) \leqslant C_\delta e^{-A}(1 + e^{-A}Z_0)/a.$$

Proof. First condition on $\mathscr{F}_{t-Ka^2}$ and apply Lemma 9.5. Then condition on $\mathscr{F}_0$ and apply (5.10) and Corollary 6.7. ☐

For $s \geqslant 0$, define the event

$$G_{Z,s} = \left\{ \sup_{0 \leqslant t \leqslant s} \left| e^{-A}Z_t^{(0)} - 1 \right| < \delta/4 \right\},$$

and set $G_{Z,n} = G_{Z,t_n}$. In using the last two results, the following lemma will be crucial:

Lemma 9.7. *Suppose $|e^{-A}Z_0 - 1| \leqslant \delta/8$. Let $s \leqslant t_{6.21}$ and set $\widehat{\mathbf{P}} = \mathbf{P}(\cdot\,|\,T > s)$. Then $\widehat{\mathbf{P}}(G_{Z,s}^c) \leqslant C_\delta e^{-A}$ for large A and a.*

Proof. Let $h(x,t) = \mathbf{P}^{(x,t)}(T > s)$ and note that $h(x,t) \geqslant 1 - Cp_B$ by (6.45). Define

$$Z_t^h = \sum_{u \in \widehat{\mathscr{N}}_t^{(0)}} (h(X_u(t), t))^{-1} w_Z(X_u(t)),$$

which is a supermartingale under $\widehat{\mathbf{P}}$ by Lemma 5.9, with $\widehat{\mathbf{E}}[Z_s^h] = (1 + O(p_B))Z_0^h$ by Corollary 6.7. For large A and a, we then have by Doob's L^2 inequality,

$$\widehat{\mathbf{P}}(G_{Z,s}^c) \leqslant \widehat{\mathbf{P}}(\sup_{0 \leqslant t \leqslant s} \left|Z_t^h - \mathbf{E}[Z_s^h]\right| \geqslant e^A\delta/16) \leqslant C_\delta e^{-2A}\widehat{\mathrm{Var}}(Z_s^h).$$

The lemma now follows from Proposition 5.2 and Corollary 6.7. ☐

The following lemma is the analogue of Lemma 9.5 for the system after the breakout. For an individual u, we define $\widehat{N}_t^{\neg u}$ and $N_t^{\mathrm{fug},\neg u}$ to be the number of hat-, respectively, fug-particles which are not descendants of u.

Lemma 9.8. *Let $t \in [T, \Theta_1]$ and let $(u, t_u) \in \widehat{\mathscr{N}_{T-}} \cup \mathscr{S}^{(\mathscr{U},T)}$. Then, for large A and a,*

$$\mathbf{P}_{\mathrm{nbab}}(\widehat{N}_t^{\neg u} + N_t^{\mathrm{fug},\neg u} < N \,|\, \mathscr{F}_\Delta)\mathbf{1}_{G_\Delta} \leqslant C_\delta \eta e^{-A}/\varepsilon.$$

Proof. As in the proof of Lemma 8.4, set $M_t = e^{-X_t^{[1]}} = (1+\Delta_t)^{-1}$, where $\Delta_t = \overline{\theta}((t-T^+)/a^2)\Delta$. Recall that on G_Δ: $|\Delta - e^{-A}Z^{(\mathscr{U},T)}| \leqslant \varepsilon^{1/8}$. For a particle $(u,t_u) \in \widehat{\mathscr{N}_{T^-}} \cup \mathscr{S}^{(\mathscr{U},T)}$, we define $\widehat{N}_t^{\neg u,0}$ and $N_t^{\mathrm{fug},\neg u,0}$ to be the number of hat-, respectively, fug-particles which are not descendants of u and which have not hit a after the time T^-, respectively, T.

Now, by Lemma 5.1 and Proposition 5.3, we have for large A and a,

$$\begin{aligned}\mathbf{E}[\widehat{N}_t^{\neg u,0} \,|\, \mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} &\geqslant Ne^{-A}M_t(\widehat{Z}_{T^-} - \|w_Z\|_\infty)(1-\delta/8)\mathbf{1}_{G_\Delta} \geqslant NM_t(1+\delta/2)\mathbf{1}_{G_\Delta}, \text{ and}\\ \mathbf{E}[N_t^{\mathrm{fug},0,\neg u} \,|\, \mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} &= Ne^{-A}M_tZ^{(\mathscr{U},T)}(\overline{\theta}((t-T)/a^2) + O(y/a))\mathbf{1}_{G_\Delta}\\ &\geqslant NM_t(\Delta_t(1+\delta) - \delta/8)\mathbf{1}_{G_\Delta}.\end{aligned}$$

In total, this gives for large A and a,

$$\mathbf{E}[\widehat{N}_t^{\neg u,0} + N_t^{\mathrm{fug},\neg u,0} \,|\, \mathscr{F}_\Delta]\mathbf{1}_{G_\Delta} \geqslant N(1+\delta/4)\mathbf{1}_{G_\Delta}. \tag{9.1}$$

Moreover, by Lemma 5.5, we have for large A and a,

$$\begin{aligned}&\mathrm{Var}(\widehat{N}_t^{\neg u,0} + N_t^{\mathrm{fug},\neg u,0} \,|\, \mathscr{F}_\Delta)\\ &\qquad \leqslant C(Ne^{-A})^2((\widehat{Z}_{T^-} + Z^{(\mathscr{U},T)})(t-T^-)/a^3 + (\widehat{Y}_{T^-}^{(0)} + Y^{(\mathscr{U},T)})) \leqslant CN^2\eta e^{-A}/\varepsilon,\end{aligned} \tag{9.2}$$

by the definition of G_Δ. Now note that under $\mathbf{P}_{\mathrm{nbab}}$, $\widehat{N}_t^{\neg u} = \widehat{N}_t^{\neg u,0}$ and $N_t^{\mathrm{fug},\neg u} = N_t^{\mathrm{fug},\neg u,0}$ almost surely. Equations (9.1) and (9.2), together with Lemma 7.5 and the conditional Chebychev inequality now yield the lemma. □

For a barrier function f, let $\mathbf{P}_f^0$ denote the law of BBM with drift $-\mu_s$ defined in (5.5) starting from a single particle at 0 (without absorption). Let R_t be the number of particles hitting a before the time t and define Z_t, Y_t by summing w_Z, respectively w_Y over the particles at time t.

Lemma 9.9. *Let $t \geqslant 0$ and f be a barrier function. Then,*

$$\mathbf{E}_f^0[Z_t] \leqslant ae^{\frac{\pi^2}{2a^2}t - \mu a}, \quad \mathbf{E}_f^0[Y_t] \leqslant e^{\frac{\pi^2}{2a^2}t-\mu a} \quad \text{and} \quad \mathbf{E}_f^0[R_t] \leqslant e^{\frac{\pi^2}{2a^2}t-\mu a}.$$

Proof. By Lemma 3.2 and Girsanov's theorem, we have

$$\mathbf{E}_f^0[Y_t] = e^{\beta_0 mt}e^{-\mu a}W_{-\mu_s}^0\Big[e^{\mu X_t}\mathbf{1}_{(X_t \geqslant 0)}\Big] \leqslant e^{\beta_0 mt}e^{-\mu a}W_{-\mu}^0\Big[e^{\mu(X_t - f(t/a^2))}\Big] \leqslant e^{\frac{\pi^2}{2a^2}t-\mu a},$$

which implies the first two inequalities, since $Z_t \leqslant aY_t$. As for the third one, by Lemma 3.1 and again Girsanov's theorem,

$$\mathbf{E}^0[R_t] = W_{-\mu}^0\Big[e^{\beta_0 mH_a}\mathbf{1}_{(H_a \leqslant t)}\Big] \leqslant e^{-\mu a}W^0\Big[e^{\frac{\pi^2}{2a^2}H_a}\mathbf{1}_{(H_a \leqslant t)}\Big] \leqslant e^{\frac{\pi^2}{2a^2}t-\mu a}.$$

Furthermore, we have $\mathbf{E}_f^0[R_t] \leqslant \mathbf{E}^0[R_t]$ as in Lemma 5.7. This finishes the proof of the lemma. □

9.3 The probability of $G_1^\sharp$

For an interval I, we define the random variable $L_I = \sum_{(u,s)\in\mathscr{L}_{H_0}} \mathbf{1}_{(s\in I)}$ counting the number of particles hitting the origin during the time interval I and set $L_n = L_{I_n}$.

Lemma 9.10. *Let $n \geqslant 0$, $s \leqslant t_{6.21}$, $I = [t_l, t_r] \subset [0,s]$ and $x \in [0,a]$. Furthermore, let f be a barrier function with $\mathrm{Err}(f, t_r) \leqslant C$. Then, for large A and a, we have*

$$\mathbf{E}_f^x[L_I \,|\, T > s] \leqslant \begin{cases} Ce^{\mu a}a^{-3}(t_r - t_l)(Aw_Y(x) + w_Z(x)), & \text{if } t_l \geqslant a^2 \text{ or } x \geqslant a/2\\ Ce^{\mu a}(w_Y(x) + a^{-3}(t_r-t_l)w_Z(x)), & \text{for any } t_l \text{ and } x.\end{cases}$$

Proof. Write $\widehat{\mathbf{E}}[\cdot] = \mathbf{E}[\cdot\,|\,T > s]$. Define $L_I^{(0)} = \sum_{(u,s)\in\mathscr{L}_{H_0}} \mathbf{1}_{(H_a(X_u)>s\in I)}$. As in the proof of Lemma 5.1, we have for every interval I, $x \in [0,a]$ and $t \leqslant s$,

$$\widehat{\mathbf{E}}_f^x[L_I^{(0)}] \leqslant e^{O(\mathrm{Err}(f,t))}\widehat{\mathbf{E}}^x[L_I^{(0)}] \leqslant Ce^{\mu x} I^a(a-x,I), \tag{9.3}$$

by (2.10) and Corollary 6.7. Lemma 2.1 and (9.3) now give,

$$\begin{aligned}\widehat{\mathbf{E}}_f^x[L_I] = \widehat{\mathbf{E}}_f^x[L_I^{(0)}] + \widehat{\mathbf{E}}_f^x\Big[\sum_{(u,s)\in\mathscr{S}_{t_r}^{(1+)}} \widehat{\mathbf{E}}_f^{(X_u(s),s)}[L_I^{(0)}]\Big]&\\ \leqslant Ce^{\mu a}\left(w_Y(x)I^a(a-x,I) + \frac{t_r - t_l}{a}\widehat{\mathbf{E}}_f^x[Z_{\varnothing,[0,t_r]}^{(1+)}]\right).&\end{aligned} \tag{9.4}$$

Lemma 2.1 now gives for large a, $I^a(a-x,I) \leqslant C\left(a^{-3}(t_r-t_l)\sin(\pi x/a) + \mathbf{1}_{(t_l<a^2 \text{ and } x\leqslant a/2)}\right)$, and by (5.38), Lemmas 5.7 and 6.8, Corollary 6.7 and (6.15), we have

$$\widehat{\mathbf{E}}_f^x[Z_{\varnothing,[0,t_r]}^{(1+)}] \leqslant CAw_Y(x)\left(1 + \frac{t_r}{a^2}\sin(\pi x/a)\right). \tag{9.5}$$

The lemma now follows from (9.4) and (9.5), together with the hypothesis $t_r \leqslant s \leqslant t_{6.21} \leqslant Ca^3/A$. □

The following lemma is crucial. It will permit to estimate the number of particles turning blue upon hitting the origin.

Lemma 9.11. *Suppose that ν_0 is deterministic with $Z_0 \geqslant (1-\delta/4)e^A$. Let $s \leqslant t_{6.21}$ and $I = [t_l,t_r] \subset [Ka^2, s]$. Write $\widehat{\mathbf{P}} = \mathbf{P}(\cdot\,|\,T > s)$. Then, for large A and a,*

$$\widehat{\mathbf{E}}\Big[\sum_{(u,t)\in\mathscr{L}_{H_0}} \mathbf{1}_{(t\in I,\, N_t^{(0)}<N)}\Big] \leqslant C_\delta e^{-A}(a^{-1} + e^{-A}Y_0)\widehat{\mathbf{E}}[L_I].$$

Proof. For an individual $u \in U$ and $t \geqslant 0$, define $N_t^{(0),\neg u} = \sum_{(v,t)\in\mathscr{N}_t^{(0)}} \mathbf{1}_{(v\not\geq u_0)}$, where u_0 is the ancestor of u at the time 0. By the trivial inequality $N_t^{(0)} \geqslant N_t^{(0),\neg u}$ for every u and by the independence of the initial particles,

$$\widehat{\mathbf{E}}\Big[\sum_{(u,t)\in\mathscr{L}_{H_0}} \mathbf{1}_{(t\in I,\, N_t^{(0)}<N)}\Big] \leqslant \widehat{\mathbf{E}}\Big[\sum_{(u,t)\in\mathscr{L}_{H_0}} \mathbf{1}_{(t\in I)}\mathbf{P}(N_t^{(0),\neg u} < N)\Big].$$

The lemma now follows from Lemma 9.5, since for every $u \in \mathscr{N}(0)$, we have $Z_0 - w_Z(X_u(0)) \geqslant Z_0 - C \geqslant (1-\delta/2)e^A$ for large A, by hypothesis. □

Before the breakout. For every $n \geqslant 0$, define the (sub-probability) measure $\mathbf{P}_n = \mathbf{P}(\cdot,\ T > t_{n+1})$. The following lemma gives an estimate on the number of *white* particles (not counting the grey ones) turning blue during the interval I_n.

Lemma 9.12. *For every $n \geqslant 1$ with $t_{n+1} \leqslant t_{6.21}$, we have for large A and a,*

$$\mathbf{E}_n[B_n\mathbf{1}_{G_{Z,n}}] \leqslant C_\delta\frac{e^{\mu a}}{a^2}.$$



Figure 5: The key step in the proof of Lemma 9.12: Conditioned on $\mathscr{F}_{t'}$, the descendants of u are independent of the descendants of the other particles by the strong branching property. Furthermore, conditioned on $\mathscr{F}_{t'}$, the probability that at time t there are less than N particles not descending from u is bounded by $C_\delta e^{-A}(a^{-1} + e^{-A} Y_{t'}^{(0),\neg u})$. These facts imply that the expected proportion of white particles turning blue upon hitting a is bounded by $C_\delta e^{-A}/a$.

Proof. Let $B_n^{(0)}$ be the number of white particles turning blue during I_n and which have not hit a before the time $t' = t_n - (K+1)a^2 = t_{n-1} + 2a^2$ and let $B_n^{(1+)} = B_n - B_n^{(0)}$. Lemmas 9.10 and 9.11 now yield,

$$\begin{aligned}\mathbf{E}_n[B_n^{(0)} \mathbf{1}_{G_{Z,n}} \,|\, \mathscr{F}_{t'}] &\leqslant C_\delta e^{-A}\Big(\frac{1}{a} + e^{-A} Y_{t'}^{(0)}\Big) \mathbf{1}_{G_{Z,t'}} \sum_{(u,s)\in \mathscr{N}_{t'}^{(0)}} \mathbf{E}_n^{(X_u(s),s)}[L_n] \\ &\leqslant C_\delta e^{\mu a - A}\Big(\frac{1}{a} Y_{t'}^{(0)} + e^{-A} (Y_{t'}^{(0)})^2\Big) \mathbf{1}_{G_{Z,t'}},\end{aligned}$$

Proposition 5.2 now gives

$$\mathbf{E}_n[B_n^{(0)} \mathbf{1}_{G_{Z,n}}] \leqslant C_\delta e^{\mu a}\Big(a^{-2} + e^{-2A} a^{-1} \mathbf{E}[Y_{t_{n-1}+a^2}^{(0)} \mathbf{1}_{G_{Z,t_{n-1}}}]\Big) \leqslant C_\delta e^{\mu a}/a^2. \tag{9.6}$$

As for the remaining particles, by the strong branching property,

$$\begin{aligned}\mathbf{E}_n[B_n^{(1+)} \mathbf{1}_{G_{Z,n}} \,|\, \mathscr{L}_{H_a \wedge t'}] &= \sum_{(u,s)\in \mathscr{R}_{t'}^{(0)}} \int_{I_n} \mathbf{E}_n^{(a,s)}[L_{[r,r+\mathrm{d}r)}] \mathbf{P}_n(N_r^{(0)} < N,\, G_{Z,n} \,|\, \mathscr{N}_{t'}) \\ &\leqslant C_\delta (e^{-A}/a) R_{t'}^{(0)} (e^{\mu a}/a) \widehat{\mathbf{Q}}^a[Z],\end{aligned}$$

by Corollary 9.6 and Lemma 9.10. Lemma 5.6 and (6.46) then give

$$\mathbf{E}_n[B_n^{(1+)} \mathbf{1}_{G_{Z,n}}] \leqslant C_\delta (e^{\mu a}/a^2) A t_{n+1}/a^3 \leqslant C_\delta e^{\mu a}/a^2, \tag{9.7}$$

by the hypothesis on t_{n+1}. The lemma now follows from (9.6) and (9.7). □

Up to now, we have only considered the white particles turning blue and will now turn to B_n^{tot}, $n \geqslant 0$. For $n \geqslant 0$ and $0 \leqslant k \leqslant n-2$, we define $B_{k,n}$ to be the number of particles that turn blue at a time $t \in I_n$, have an ancestor that turned blue at a time $t' \in I_k$, have none that has hit 0 between t_{k+2} and t_n, have not hit a between t' and t_{k+2} and did not break out between t_{k+2} and t.

Let $G_{\mathrm{b},n}$ the event that no blue particle hits a before t_n and $G_{\mathrm{g},n}$ the event that no particle that turned from white to blue before t_{n-1} has a grey descendant that breaks out before $t_{6.21}$). Then set

$$G_n^{\mathrm{tot}} = G_{Z,n} \cap G_{\mathrm{b},n} \cap G_{\mathrm{g},n} \cap \{B_0 \leqslant e^A e^{\mu a}/a^2,\, Y_0 \leqslant \eta\}.$$

Lemma 9.13. *For every $n \geqslant 0$ with $t_{n+1} \leqslant t_{6.21}$ and $k \leqslant n-2$, we have for large A and a,*

$$\mathbf{E}_n[B_{k,n}\mathbf{1}_{G_n^{\mathrm{tot}}}] \leqslant C_\delta \mathbf{E}_k[B_k^{\mathrm{tot}}\mathbf{1}_{G_k^{\mathrm{tot}}}]\frac{e^{-A}}{a}.$$

Proof. Let $\mathscr{B}_k$ be the stopping line consisting of the particles that turn blue during I_k, at the moment at which they turn blue (hence, $B_k^{\mathrm{tot}} = \#\mathscr{B}_k$). Note that $\mathscr{B}_k \wedge \mathscr{N}^{(0)}_{t_{n-1}} = \mathscr{B}_k \cup \mathscr{N}^{(0)}_{t_{n-1}}$, such that the descendants of $\mathscr{B}_k$ and of $\mathscr{N}^{(0)}_{t_{n-1}}$ are independent, given their past, by the strong branching property. Note also that $G_k^{\mathrm{tot}} \in \mathscr{F}_{\mathscr{B}_k \wedge \mathscr{N}^{(0)}_{t_{n-1}}}$, since by definition this σ-field contains all the information about the descendants of the particles in $\mathscr{B}_j$, $j < k$. By Corollary 9.6, we then have

$$\begin{aligned}\mathbf{E}_n&[B_{k,n}\mathbf{1}_{G_n^{\mathrm{tot}}} \mid \mathscr{F}_{\mathscr{B}_k \wedge \mathscr{N}^{(0)}_{t_{n-1}}}]\\ &\leqslant C_\delta \frac{e^{-A}}{a}\mathbf{1}_{G_k^{\mathrm{tot}}}\mathbf{1}_{(T>t_{k+1})}\sum_{(u,s)\in\mathscr{B}_k}\mathbf{E}_n^{(0,s)}\Big[\sum_{(v,t)\in\mathscr{L}_{H_0^{k+2}}}\mathbf{1}_{(H_a(X_v)>t_{k+2},\,X_v(t_{k+2})\in(0,a),\,t\in I_n)}\Big],\end{aligned} \tag{9.8}$$

where here $H_0^{k+2}(X) = \inf\{t \geqslant t_{k+2} : X_t = 0\}$. By Lemmas 9.9 and 9.10, each summand on the right-hand side of the above inequality is bounded by $C_\delta e^{\mu a}\mathbf{E}_n^0[Y_{t_{k+2}-s}] \leqslant C_\delta$. The lemma follows. □

Lemma 9.14. *For all $n \geqslant 1$ with $t_{n+1} \leqslant t_{6.21}$, we have $\mathbf{E}_n[B_n^{\mathrm{tot}}\mathbf{1}_{G_n^{\mathrm{tot}}}] \leqslant C_\delta e^{\mu a}/a^2$ for large A and a.*

Proof. By Lemma 9.12, the statement is true for $n = 1$, because $B_1^{\mathrm{tot}} = B_1$ by definition. Now we have for every n, by Lemmas 9.12 and 9.13 and the fact that $B_0^{\mathrm{tot}} = B_0 \leqslant e^A e^{\mu a}/a^2$ on G_n^{tot},

$$\mathbf{E}_n[B_n^{\mathrm{tot}}\mathbf{1}_{G_n^{\mathrm{tot}}}] = \sum_{k=0}^{n-2}\mathbf{E}_n[B_{k,n}\mathbf{1}_{G_n^{\mathrm{tot}}}] + \mathbf{E}_n[B_n\mathbf{1}_{G_n^{\mathrm{tot}}}] \leqslant C_\delta\Big(\frac{e^{-A}}{a}\sum_{k=1}^{n-2}\mathbf{E}_k[B_k^{\mathrm{tot}}\mathbf{1}_{G_k^{\mathrm{tot}}}] + \frac{e^{\mu a}}{a^2}\Big). \tag{9.9}$$

The lemma now follows easily by induction over n, since $n \leqslant a$ by hypothesis. □

Lemma 9.15. *Suppose ($HB_0^\sharp$). For large A and a, we have $\mathbf{P}_n((G_n^{\mathrm{tot}})^c) \leqslant C_\delta e^{-2A/3}n/a + \varepsilon^2$ for every $n \geqslant 0$.*

Proof. Let $n \geqslant 1$. By definition, the event $G_{n-1}^{\mathrm{tot}}\backslash G_{\mathrm{b},n}$ implies that a descendant of a particle in $\mathscr{B}_{n-2}$ hits a before t_n. Markov's inequality and Lemma 9.9 then imply,

$$\mathbf{P}_n(G_{n-1}^{\mathrm{tot}}\backslash G_{\mathrm{b},n}) \leqslant \mathbf{E}_n[B_{n-2}^{\mathrm{tot}}\mathbf{1}_{G_{n-1}^{\mathrm{tot}}}] \times \sup_{t\leqslant t_2}\mathbf{E}^0[R_t] \leqslant C_\delta a^{-2}, \tag{9.10}$$

by Lemma 9.14. Furthermore, by Proposition 6.2 and Lemmas 9.9 and 9.14, we have

$$\mathbf{P}_n((G_{n-1}^{\mathrm{tot}} \cap G_{\mathrm{b},n})\backslash G_{\mathrm{g},n}) \leqslant \mathbf{E}_n[B_{n-2}^{\mathrm{tot}}\mathbf{1}_{G_{n-1}^{\mathrm{tot}}}] \times \sup_{t\leqslant t_2} p_B\mathbf{E}^0[t_{6.21}Z_t + Y_t] \leqslant C_\delta e^{-2A/3}a^{-1}, \tag{9.11}$$

by (6.14) and (6.2). By (9.10) and (9.11), we have for large A and a,

$$\mathbf{P}_n(G_{Z,n}\backslash G_n^{\mathrm{tot}}) \leqslant \mathbf{P}_n(G_{Z,n}\backslash G_{n-1}^{\mathrm{tot}}) + C_\delta e^{-2A/3}a^{-1} \leqslant \mathbf{P}_{n-1}(G_{Z,n-1}\backslash G_{n-1}^{\mathrm{tot}}) + C_\delta e^{-2A/3}a^{-1}.$$

By induction over n and the hypothesis, this gives for every n,

$$\mathbf{P}_n(G_{Z,n}\backslash G_n^{\mathrm{tot}}) \leqslant C_\delta e^{-2A/3}(n/a) + \varepsilon^2.$$

Together with Lemma 9.7, the statement follows. □

Define the random variable $n_0 := \lfloor T/t_1 \rfloor$ and note that $t_{n_0-1} > T^-$ for large A. Define the events $G_{\mathrm{nw}} = \{Z^{\mathrm{gr}}_{t_{n_0}} \leqslant e^{A/2},\ Y^{\mathrm{gr}}_{t_{n_0}} \leqslant e^{A/2}/a,\ B^{\mathrm{gr}}_{n_0-1} \leqslant \eta e^{\mu a}/a\}$ and $G^{\mathrm{tot}}_{n_0} = \bigcup_n (G^{\mathrm{tot}}_n \cap \{n_0 = n\})$.

Lemma 9.16. *For large A and a, $\mathbf{P}((G^{\mathrm{tot}}_{n_0} \cap G_{\mathscr{U}}) \backslash G_{\mathrm{nw}}) \leqslant C_\delta e^{-A/2}$.*

Proof. By definition, $\mathscr{F}_{\mathscr{U}} \subset \mathscr{F}_{\mathscr{L}_{H_0}}$. Lemmas 6.9, 9.9 and 9.14 then give

$$\begin{aligned}\mathbf{E}[Z^{\mathrm{gr}}_{t_{n_0}} \mathbf{1}_{G^{\mathrm{tot}}_{n_0} \cap G_{\mathscr{U}}}] &\leqslant C \sup_{t \leqslant t_2} \mathbf{E}^0[Z_t + AY_t] \Big(\sum_{n=0}^{\infty} \mathbf{E}\Big[\mathbf{E}_n[B^{\mathrm{tot}}_n \mathbf{1}_{G^{\mathrm{tot}}_n} \,|\, \mathscr{F}_{\mathscr{L}_{H_0}}] \mathbf{1}_{G_{\mathscr{U}}}\Big]\Big) \\ &\leqslant C_\delta a e^{\mu a} \sum_{n=0}^{\lfloor \sqrt{\varepsilon} a^3/t_1 \rfloor} \mathbf{E}_n[B^{\mathrm{tot}}_n \mathbf{1}_{G^{\mathrm{tot}}_n}] \leqslant C_\delta.\end{aligned} \tag{9.12}$$

Similarly, we have

$$\mathbf{P}(R^{\mathrm{gr}}_{I_{n_0-1}} > 0,\ G^{\mathrm{tot}}_{n_0},\ G_{\mathscr{U}}) \leqslant C_\delta \eta, \quad \text{and} \quad \mathbf{E}[Y^{\mathrm{gr}}_{t_{n_0}} \mathbf{1}_{(R^{\mathrm{gr}}_{I_{n_0-1}}=0)} \mathbf{1}_{G^{\mathrm{tot}}_{n_0} \cap G_{\mathscr{U}}}] \leqslant C_\delta/a. \tag{9.13}$$

Finally, we have as in (9.12), by Lemmas 9.13 and 9.14,

$$\begin{aligned}\mathbf{E}[B^{\mathrm{gr}}_{n_0-1} \mathbf{1}_{G^{\mathrm{tot}}_{n_0-1} \cap G_{\mathscr{U}}}] &= \mathbf{E}\Big[\sum_{k=0}^{n_0-2} \mathbf{E}[B_{k,n_0-1} \mathbf{1}_{G^{\mathrm{tot}}_{n_0-1}} \,|\, \mathscr{F}_{\mathscr{L}_{H_0}}] \mathbf{1}_{G_{\mathscr{U}}}\Big] \\ &\leqslant C_\delta (e^{-A}/a) \sum_{k=0}^{\lfloor \sqrt{\varepsilon} a^3/t_1 \rfloor} \mathbf{E}_k[B^{\mathrm{tot}}_k \mathbf{1}_{G^{\mathrm{tot}}_k}] \leqslant C_\delta e^{-A+\mu a}/a^2.\end{aligned} \tag{9.14}$$

The lemma now follows from (9.12), (9.13) and (9.14), together with Markov's inequality. □

After the breakout. We study now the system after the breakout. Recall that $n_0 = \lfloor T/t_1 \rfloor$ by definition and define $n_1 := \lfloor \Theta_1/t_1 + 1 \rfloor$. We define $\mathscr{F}^\sharp_\Delta = \mathscr{F}_\Delta \cap \mathscr{F}_{\mathscr{N}^{\mathrm{gr}}_{t_{n_0}} \wedge \mathscr{B}_{n_0-1}}$ and the event $G^\sharp_\Delta = G_\Delta \cap G_{\mathrm{nw}} \in \mathscr{F}^\sharp_\Delta$. We denote by the superscript "gr≤" the quantities relative to particles the descending from those that were grey before or at time t_{n_0}. Then let $G^\sharp_{\mathrm{nbab}}$ be the intersection of G_{nbab} with the event that none of these particles hits a between t_{n_0} and $\Theta_1 + e^A a^2$ before hitting 0. Define then the (sub-probability) measure $\mathbf{P}^\sharp_{\mathrm{nbab}} = \mathbf{P}(\cdot,\, G^\sharp_{\mathrm{nbab}})$.

Lemma 9.17. *For large A and a, $\mathbf{P}(G^\sharp_\Delta) \geqslant 1 - C\varepsilon^{5/4}$ and $\mathbf{P}((G^\sharp_{\mathrm{nbab}})^c \,|\, \mathscr{F}^\sharp_\Delta) \mathbf{1}_{G^\sharp_\Delta} \leqslant C\varepsilon^2$.*

Proof. On G_Δ, we have $t_{n_0} \leqslant t_{6.21}$ for large A and a, in particular, $n_0 \leqslant a$. By Lemmas 9.15 and 9.16 and (6.2), we then have for large A and a,

$$\mathbf{P}(G_\Delta \backslash G^\sharp_\Delta) \leqslant \mathbf{P}(G_\Delta \backslash G^{\mathrm{tot}}_{n_0}) + \mathbf{P}((G_\Delta \cap G^{\mathrm{tot}}_{n_0}) \backslash G_{\mathrm{nw}}) \leqslant C\varepsilon^2.$$

With Lemma 7.4, this yields the first statement. For the second statement, we have by Lemmas 5.6 and 5.7,

$$\mathbf{P}(R^{\mathrm{gr}\preceq}_{[t_{n_0}, \Theta_1 + e^A a^2]} > 0 \,|\, \mathscr{F}^\sharp_\Delta) \mathbf{1}_{G^\sharp_\Delta} \leqslant C((e^A/a) Z^{\mathrm{gr}}_{t_{n_0}} + Y^{\mathrm{gr}}_{t_{n_0}}) \mathbf{1}_{G^\sharp_\Delta} \leqslant C e^{A/2} \eta.$$

Together with Lemma 7.5, this implies the lemma. □

We change the notation of B_n a bit: it is defined to be the number of particles hitting 0 during I_n for the first time *after* t_{n_0-1}. In particular, we also count the descendants of the grey particles at that time.

Lemma 9.18. *For large A and a and $n \geqslant n_0 - 1$, we have $\mathbf{E}^\sharp_{\mathrm{nbab}}[B_n \,|\, \mathscr{F}^\sharp_\Delta] \mathbf{1}_{G^\sharp_\Delta} \leqslant C_\delta e^{-5A/3} e^{\mu a}/a$.*

Proof. Define $\mathscr{L} := \widehat{\mathscr{N}_{T-}} \cup \mathscr{S}^{(\mathscr{U},T)}$. We have for $t \in I_n$, as in Lemma 9.11,

$$\begin{aligned}\mathbf{E}^{\sharp}_{\mathrm{nbab}}[\widehat{B}_n + B_n^{\mathrm{fug}} \,|\, \mathscr{F}^{\sharp}_{\Delta}]\mathbf{1}_{G^{\sharp}_{\Delta}} &\leqslant C \sum_{(u,t)\in\mathscr{L}} \widehat{\mathbf{E}}^{(X_u(t),t)}_{X^{[1]}}[L_n]\mathbf{P}(\widehat{N}_t'^{\neg u} + N_t'^{\mathrm{fug},\neg u} < N \,|\, \mathscr{F}_{\Delta})\mathbf{1}_{G^{\sharp}_{\Delta}}\\ &\leqslant C_\delta \eta (\varepsilon e^A)^{-1} \sum_{(u,t)\in\mathscr{L}} \widehat{\mathbf{E}}^{(X_u(t),t)}_{X^{[1]}}[L_n]\mathbf{1}_{G^{\sharp}_{\Delta}}\end{aligned}$$

by Lemma 9.8. By Lemma 9.10, (6.1) and the definition of $\widehat{G}$ and G_{fug}, this gives for large A and a,

$$\begin{aligned}\mathbf{E}^{\sharp}_{\mathrm{nbab}}[\widehat{B}_n + B_n^{\mathrm{fug}} \,|\, \mathscr{F}^{\sharp}_{\Delta}]\mathbf{1}_{G^{\sharp}_{\Delta}} &\leqslant C_\delta e^{\mu a}\eta(\varepsilon e^A a)^{-1}(\widehat{Z}_{T-} + A\widehat{Y}_{T-} + Z^{(\mathscr{U},T)})\mathbf{1}_{G^{\sharp}_{\Delta}}\\ &\leqslant C_\delta e^{\mu a}\eta\varepsilon^{-2}/a.\end{aligned} \tag{9.15}$$

Furthermore, by the independence of the check- and bar particles from the others, we have by Lemmas 9.8 and 9.10 and by the inequality $w_Y(x) \leqslant a^{-1}e^{-(a-x)/2}$, valid for every $x \leqslant a/2$ and a large enough,

$$\begin{aligned}&\mathbf{E}^{\sharp}_{\mathrm{nbab}}[\check{B}_n + \bar{B}_n \,|\, \mathscr{F}^{\sharp}_{\Delta}]\mathbf{1}_{G^{\sharp}_{\Delta}}\\ &\qquad\leqslant C_\delta e^{\mu a}\eta(\varepsilon e^A)^{-1}a^{-1}(\check{Z}_{\varnothing\vee T-} + A\check{Y}_{\varnothing\vee T-} + A\mathscr{E}_{\mathscr{U}} + Z^{\mathrm{gr}}_{t_{n_0}} + AY^{\mathrm{gr}}_{t_{n_0}})\mathbf{1}_{G^{\sharp}_{\Delta}} \leqslant C_\delta e^{\mu a}\eta/a,\end{aligned} \tag{9.16}$$

by (6.1) and the definition of $G^{\sharp}_{\Delta}$. Similarly, we have for $n \geqslant n_0$,

$$\mathbf{E}^{\sharp}_{\mathrm{nbab}}[B_n^{\mathrm{gr}\precsim} \,|\, \mathscr{F}^{\sharp}_{\Delta}]\mathbf{1}_{G^{\sharp}_{\Delta}} \leqslant C_\delta e^{\mu a}\eta(\varepsilon e^A)^{-1}Y^{\mathrm{gr}}_{t_{n_0}}\mathbf{1}_{G^{\sharp}_{\Delta}} \leqslant C_\delta e^{\mu a}\eta/a, \tag{9.17}$$

and $B^{\mathrm{gr}\precsim}_{n_0-1} \leqslant \eta e^{\mu a}/a$ on the event $G^{\sharp}_{\Delta}$. The lemma now follows from (9.15), (9.16) and (9.17), together with (6.3) and (6.2). ☐

For $n \geqslant n_0 - 1$, define $\widetilde{G}^{\mathrm{tot}}_n$ to be the event that no descendant of a particle which has been coloured blue between t_{n_0-1} and t_n hits a before $\Theta_1 + e^A a^2$.

Lemma 9.19. *We have for every $n \geqslant n_0$ and for large A and a,*

$$\mathbf{E}^{\sharp}_{\mathrm{nbab}}[B^{\mathrm{tot}}_n \mathbf{1}_{\widetilde{G}^{\mathrm{tot}}_n} \,|\, \mathscr{F}^{\sharp}_{\Delta}]\mathbf{1}_{G^{\sharp}_{\Delta}} \leqslant C_\delta e^{-5A/3}\frac{e^{\mu a}}{a}.$$

Proof. The proof is similar to the proof of Lemma 9.14: Substituting Lemma 9.5 by Lemma 9.8, one first shows that for every $n_0 - 1 \leqslant k \leqslant n-2$, one has

$$\mathbf{E}^{\sharp}_{\mathrm{nbab}}[B_{k,n}\mathbf{1}_{\widetilde{G}^{\mathrm{tot}}_n} \,|\, \mathscr{F}_{\mathscr{B}_k}]\mathbf{1}_{G^{\sharp}_{\Delta}} \leqslant C_\delta \eta e^{-A}\mathbf{E}^{\sharp}_{\mathrm{nbab}}[B^{\mathrm{tot}}_k \mathbf{1}_{\widetilde{G}^{\mathrm{tot}}_k} \,|\, \mathscr{F}^{\sharp}_{\Delta}]\mathbf{1}_{G^{\sharp}_{\Delta}}, \tag{9.18}$$

which, by a recurrence similar to (9.9), yields the lemma. ☐

Lemma 9.20. *For large A and a, $\mathbf{P}^{\sharp}_{\mathrm{nbab}}((\widetilde{G}^{\mathrm{tot}}_{n_1})^c \,|\, \mathscr{F}^{\sharp}_{\Delta})\mathbf{1}_{G^{\sharp}_{\Delta}} \leqslant 1/a$.*

Proof. Similarly to the proof of Lemma 9.15, but using Lemmas 5.6 and 5.7 instead of Proposition 6.2, we have for large A and a,

$$\mathbf{P}^{\sharp}_{\mathrm{nbab}}(\widetilde{G}^{\mathrm{tot}}_{n-1}\backslash\widetilde{G}^{\mathrm{tot}}_n \,|\, \mathscr{F}^{\sharp}_{\Delta})\mathbf{1}_{G^{\sharp}_{\Delta}} \leqslant C_\delta \mathbf{E}^{\sharp}_{\mathrm{nbab}}[B^{\mathrm{tot}}_{n-2}\mathbf{1}_{\widetilde{G}^{\mathrm{tot}}_{n-1}} \,|\, \mathscr{F}^{\sharp}_{\Delta}]\mathbf{1}_{G^{\sharp}_{\Delta}} e^{A-\mu a}/a \leqslant C_\delta a^{-2},$$

where the last inequality follows from Lemma 9.19 and (6.3). The lemma now follows by induction over n. ☐

Lemma 9.21. *Suppose* (*HB*$_0^\sharp$). *Then* $\mathbf{P}(G_1^\sharp) \geqslant 1 - \varepsilon^{4/3}$ *for large* A *and* a.

Proof. By Lemma 9.5, Corollary 9.6 and the union bound we have

$$\mathbf{P}(\exists n \leqslant n_0 : N(t_n) < N,\ G_{Z,n_0},\ G_{\mathscr{U}}) \leqslant C_\delta e^{-A}. \tag{9.19}$$

Furthermore, by Lemma 9.8, we have

$$\mathbf{P}(\exists n_0 < n < n_1 : N(t_n) < N \text{ or } N(\Theta_1) < N,\ G_\Delta) \leqslant C_\delta \eta/\varepsilon \leqslant C_\delta e^{-A}, \tag{9.20}$$

by (6.3) and (6.2). Now, by Proposition 5.2 and Lemmas 9.9 and 9.17,

$$\begin{aligned} \mathbf{E}^\sharp_{\mathrm{nbab}}[Z^{\mathrm{tot}}_{\Theta_1}\mathbf{1}_{\widetilde{G}^{\mathrm{tot}}_{n_1}} \mid \mathscr{F}^\sharp_\Delta]\mathbf{1}_{G^\sharp_\Delta} &\leqslant C\Big(Z^{\mathrm{nw}}_{t_{n_0}} + C_\delta a e^{\mu a}\sum_{n=n_0-1}^{n_1-1}\mathbf{E}^\sharp_{\mathrm{nbab}}[B^{\mathrm{tot}}_n \mathbf{1}_{\widetilde{G}^{\mathrm{tot}}_n} \mid \mathscr{F}^\sharp_\Delta]\Big)\mathbf{1}_{G^\sharp_\Delta} \\ &\leqslant Ce^{A/2} + C_\delta \leqslant Ce^{A/2}, \end{aligned} \tag{9.21}$$

by Lemma 9.19. Similarly, we get

$$\mathbf{E}^\sharp_{\mathrm{nbab}}[Y^{\mathrm{tot}}_{\Theta_1}\mathbf{1}_{\widetilde{G}^{\mathrm{tot}}_{n_1}} \mid \mathscr{F}^\sharp_\Delta]\mathbf{1}_{G^\sharp_\Delta} \leqslant Ce^{A/2}/a. \tag{9.22}$$

Moreover, we have by Lemma 9.19,

$$\mathbf{E}^\sharp_{\mathrm{nbab}}[(B^{\mathrm{tot}}_{n_1-1} + B^{\mathrm{tot}}_{n_1})\mathbf{1}_{\widetilde{G}^{\mathrm{tot}}_{n_1}} \mid \mathscr{F}^\sharp_\Delta]\mathbf{1}_{G^\sharp_\Delta} \leqslant C_\delta e^{\mu a - 5A/3}/a.$$

Setting $X = \mathbf{P}\Big(B^{\mathrm{tot}}_{[\Theta_1,\Theta_1+t_1]} \leqslant e^{-A}e^{\mu a}/a \,\Big|\, \mathscr{F}_{\Theta_1}\Big)$, we then have $\mathbf{E}[(1-X)\mathbf{1}_{G^\sharp_{\mathrm{nbab}}\cap\widetilde{G}^{\mathrm{tot}}_{n_1}\cap G^\sharp_\Delta}] \leqslant C_\delta e^{-2A/3}$ by Markov's inequality. Applying the Markov inequality once more to X as in the proof of Lemma 8.1 yields

$$\mathbf{P}(X \leqslant 1 - \varepsilon^2,\ G^\sharp_{\mathrm{nbab}} \cap \widetilde{G}^{\mathrm{tot}}_{n_1} \cap G^\sharp_\Delta) \leqslant C_\delta \varepsilon^{-2} e^{-2A/3} \leqslant C_\delta \varepsilon^2, \tag{9.23}$$

by (6.2). The lemma now follows from (9.19), (9.20) and (9.23), Lemmas 9.17 and 9.20 and Markov's inequality applied to (9.21) and (9.22). □

9.4 Proofs of the main results

Proof of Lemma 9.1. The property about Z_0 and Y_0 follows as in the proof of Lemma 8.8 from (8.21). Moreover, again as in the proof of Lemma 8.8, we have by (8.21), (8.23) and (8.25), for every $i \leqslant \lceil e^\delta N\rceil$,

$$\mathbf{P}(N^{-i}_{t_1} < N,\ R_{t_1} = 0) \leqslant C_\delta e^{-A}/a \quad \text{and} \quad \mathbf{P}(R_{t_1} > 0) \leqslant C_\delta e^{A}/a,$$

where we denote the initial particles by $1, \dots, \lceil e^\delta N\rceil$. By the independence of the particles, we now have,

$$\begin{aligned} \mathbf{E}[B_0 \mathbf{1}_{(R_{t_1}=0)}] &\leqslant \sum_{i=1}^{\lceil e^\delta N\rceil} \mathbf{P}(N^{-i}_{t_1} < N,\ R_{t_1} = 0)\mathbf{E}^{X_i(0)}[L_0\mathbf{1}_{(R_{t_1}=0)}] \\ &\leqslant C_\delta (e^{-A}/a)e^{\mu a}\mathbf{E}[Y_0] \leqslant C_\delta e^{\mu a}/a^2, \end{aligned}$$

by (8.21). Setting $X = \mathbf{P}\Big(B_0 \leqslant e^{-A}e^{\mu a}/a \,\Big|\, \mathscr{F}_0\Big)$, we then have $\mathbf{E}[1 - X] \leqslant C_\delta e^{A}/a$ by Markov's inequality. Applying Markov's inequality once more to X as in the proof of Lemma 8.1 yields the lemma. □

Proof of Proposition 9.2. Exactly the same reasoning as in the proof of Proposition 8.2, but using Lemma 9.21 instead of Lemma 8.14. ☐

Proof of Proposition 9.3. Define $\rho = x_{\alpha e^{-2\delta}}$, such that $(1+\mu\rho)e^{-\mu\rho} \leqslant \alpha e^{-2\delta}(1+o(1))$. As in the proof of Proposition 8.3 (see (8.57)), it is enough to show that

$$\sup_{\nu\in G_0^\sharp}\sup_{t\geqslant e^A a^2} \mathbf{P}^\nu(N_t^{\mathrm{tot}}(\rho) > \alpha N,\, T > t) \leqslant C_\delta \varepsilon^{5/4}, \tag{9.24}$$

for large A and a. Let $\nu \in G_0^\sharp$ and $t \geqslant e^A a^2$. If $t > \sqrt{\varepsilon}a^3$, then the above probability is bounded by $C\varepsilon^2$ by Proposition 6.2. Suppose therefore that $e^A a^2 \leqslant t \leqslant \sqrt{\varepsilon}a^3$ and write $\widehat{\mathbf{P}} = \mathbf{P}(\cdot \,|\, T > t)$. By definition, we have $(1+x_\alpha)e^{-x_\alpha} = \alpha$. Let n be the largest integer such that $t_{n+1} < t$; note that $n \geqslant 1$ for large A. By Lemmas 6.9 and 6.12 together with Chebychev's and Markov's inequalities, (6.2) and (6.1), we have for large A and a,

$$\widehat{\mathbf{P}}(|e^{-A}Z_{t_n} - 1| \leqslant \varepsilon^{1/8},\, Y_{t_n} \leqslant e^{4A/3}/a) \geqslant 1 - C_\delta\varepsilon^{5/4}. \tag{9.25}$$

And as in the proof of Lemma 9.16, the same holds for $Z_{t_n}^{\mathrm{gr}}$ and $Y_{t_n}^{\mathrm{gr}}$ as well, which yields

$$\widehat{\mathbf{P}}(|e^{-A}Z_{t_n}^{\mathrm{wg}} - 1| \leqslant \varepsilon^{1/8},\, Y_{t_n}^{\mathrm{wg}} \leqslant e^{4A/3}/a) \geqslant 1 - C_\delta\varepsilon^{5/4}, \tag{9.26}$$

Lemmas 5.6 and 9.9 and Corollary 6.7 then show that $\mathbf{P}(R_{[t_n,t]}^{\mathrm{tot}} > 0) \leqslant o(1)$.

Let $\widetilde{N}_t^{\mathrm{wg}}(\rho)$ be the white and grey particles to the right of ρ at time t which descend from the white and grey particles at time t_n and which have not hit a between t_n and t. By Proposition 5.3 and Corollary 6.7, we have for large A and a,

$$\widehat{\mathbf{E}}[\widetilde{N}_t^{\mathrm{wg}}(\rho) \,|\, \mathscr{F}_{t_n}] \leqslant e^{-\delta/2}\alpha N e^{-A} Z_{t_n}^{\mathrm{wg}}, \tag{9.27}$$

$$\widehat{\mathrm{Var}}(\widetilde{N}_t^{\mathrm{wg}}(\rho) \,|\, \mathscr{F}_{t_n}) \leqslant C_{\delta,\alpha}\alpha^2 N^2 e^{-2A}(a^{-1}Z_{t_n}^{\mathrm{wg}} + Y_{t_n}^{\mathrm{wg}}). \tag{9.28}$$

Chebychev's inequality and (9.26) then give for large A and a,

$$\mathbf{P}(\widetilde{N}_t^{\mathrm{wg}}(\rho) > e^{-\delta/4}\alpha N) \leqslant C\varepsilon^{5/4}. \tag{9.29}$$

Furthermore, denote by $\widetilde{N}_t^{\mathrm{blue}}(\rho)$ the number of blue particles to the right of ρ at time t which have turned blue after $t_n + a^2$ and which have not hit a between t_n and t. Then by Lemma 9.11 and (9.3), we have

$$\begin{aligned}\widehat{\mathbf{E}}[\widetilde{N}_t^{\mathrm{blue}}(\rho) \,|\, \mathscr{F}_{t_n}] &\leqslant C_\delta e^{-A}(a^{-1} + e^{-A}Y_{t_n}^{\mathrm{wg}}) \sum_{(u,s)\in\mathscr{N}_{t_n}^{\mathrm{wg}}} \int_{t_n+a^2}^{t} \widehat{\mathbf{E}}[L_{[\tau,\tau+\mathrm{d}\tau]}(u,s)]\mathbf{E}^0[N_{t-\tau}(\rho)]\\ &\leqslant C_\delta e^{-A}(a^{-1} + e^{-A}Y_{t_n}^{\mathrm{wg}}) \sum_{(u,s)\in\mathscr{N}_{t_n}^{\mathrm{wg}}} e^{\mu X_u(s)} \int_{a^2}^{d} I^a(a - X_u(s), \mathrm{d}\tau)\mathbf{E}^0[N_{t-t_n-\tau}(\rho)].\end{aligned} \tag{9.30}$$

Set $d = t - t_n$. Note that $(K+3)a^2 \leqslant d \leqslant 2(K+3)a^2$. By Lemma 3.2 and Girsanov's theorem, we have now for every $\tau \geqslant 0$,

$$\mathbf{E}^0[N_\tau(\rho)] = e^{\beta_0 m\tau}W_{-\mu}^0(X_\tau > \rho) = e^{\frac{\pi^2}{2a^2}\tau}W^0[e^{-\mu\rho}\mathbf{1}_{(X_\tau>\rho)}] = e^{\frac{\pi^2}{2a^2}\tau}\int_\rho^\infty e^{-\mu z}g_{d-\tau}(z)\,\mathrm{d}z, \tag{9.31}$$

where $g_\tau(x) = (2\pi\tau)^{-1/2}e^{-x^2/(2\tau)}$ is the Gaussian density with variance τ. If $\tau \geqslant a^2$, then $\sup_z g_\tau(z) \leqslant C/a$, such that for every $x \in [0,a]$ and $z \geqslant \rho$,

$$\int_{a^2}^{d-a^2} I^a(a-x, \mathrm{d}\tau)g_{d-\tau}(z) \leqslant Ca^{-1}I^a(a-x, [a^2, d-a^2]) \leqslant CKa^{-1}\sin(\pi x/a), \tag{9.32}$$

by Lemma 2.1. Moreover, by Lemma 2.1, we have $I^a(a-x,\mathrm{d}\tau) \leqslant Ca^{-2}\sin(\pi x/a)\,\mathrm{d}\tau$ for every $\tau \geqslant a^2$, such that

$$\int_{d-a^2}^{d} I^a(a-x,\mathrm{d}\tau) g_{d-\tau}(z) \leqslant Ca^{-2}\sin(\pi x/a)\int_0^{a^2} \tau^{-1/2}\,\mathrm{d}\tau \leqslant Ca^{-1}\sin(\pi x/a). \tag{9.33}$$

Equations (9.30), (9.31), (9.32) and (9.33) now yield

$$\widehat{\mathbf{E}}[\widetilde{N}_t^{\mathrm{blue}}(\rho)\,|\,\mathscr{F}_{t_n}] \leqslant Ce^{-\mu\rho}e^{\mu a - A}a^{-2}(Ka^{-1} + e^{-A}Y_{t_n}^{\mathrm{wg}})Z_{t_n}^{\mathrm{wg}} \leqslant C_\delta \alpha e^{-2A}N(1+ae^{-A}Y_{t_n}^{\mathrm{wg}})Z_{t_n}^{\mathrm{wg}}. \tag{9.34}$$

Furthermore, if $N_t^{\mathrm{rest}}(\rho)$ denotes the particles to the right of ρ at time t descending from those turning blue between t_{n-1} and $t_n + a^2$, then by (9.31) and the supremum bound on $g_\tau(z)$,

$$\widehat{\mathbf{E}}[N_t^{\mathrm{rest}}(\rho)G_n^{\mathrm{tot}}] \leqslant C_\delta a^{-1}e^{-\mu\rho}\mathbf{E}[(B_n^{\mathrm{tot}} + B_{n-1}^{\mathrm{tot}})\mathbf{1}_{G_n^{\mathrm{tot}}}] \leqslant C_\delta \alpha e^{-A}N, \tag{9.35}$$

by Lemma 9.14. Markov's inequality applied to (9.35) and (9.34) together with (9.26), Lemmas 9.15 and 9.7 give

$$\mathbf{P}(N_t^{\mathrm{rest}} + \widetilde{N}_t^{\mathrm{blue}} > \varepsilon\alpha N) \leqslant C_\delta\varepsilon^2, \tag{9.36}$$

by (6.2). The statement now follows from (9.29), (9.36) and the above-mentioned bound on $\mathbf{P}(R_{[t_n,t]}^{\mathrm{tot}} > 0)$. ☐

Proof of Lemma 9.4. By definition, the B$^\sharp$-BBM and C$^\sharp$-BBM coincide until Θ_n on the set $G_n^\sharp$. By Proposition 9.2, we have $\mathbf{P}(G_n^\sharp) \geqslant 1 - n\varepsilon^{1+b}$ for some numerical constant $b > 0$ and by the coupling of $(\Theta_n)_{n\geqslant 0}$ with a Poisson process of intensity $p_B e^A \pi \sim \varepsilon^{-1}$, we have $\mathbf{P}(\Theta_{\varepsilon^{-1-b/2}} > \varepsilon^{-b/4}) \to 1$, as A and a go to infinity. Propositions 9.2 and 9.3 then readily transfer to the C$^\sharp$-BBM. ☐

10 Coupling the N-BBM with other braching Brownian motions

We will first establish a monotone coupling between the N-BBM and a class of slightly more general BBM with selection which includes the B$^\flat$-BBM from Section 8 and the C$^\sharp$-BBM from Lemma 9.4. In a second part, Theorem 1.1 is proven.

10.1 A monotone coupling between N-BBM and more general particle systems

A *selection mechanism* for branching Brownian motion is by definition a stopping line $\mathscr{L}$, which has the interpretation that if $(u,t) \in \mathscr{L}$, we think of u being *killed* at the time t. The set of particles in the system at time t then consists of all the particles $u \in \mathscr{N}(t)$ which do not have an ancestor which has been killed at a time $s \leqslant t$, i.e. all the particles $u \in \mathscr{N}(t)$ with $\mathscr{L} \not\preceq (u,t)$.

Now suppose we have two systems of BBM with selection, the N^+-BBM and the N^--BBM, whose selection mechanisms satisfy the following rules.

1. Only left-most particles are killed, i.e. if $(u,t) \in \mathscr{L}$, then $X_u(t) \leqslant X_v(t)$ for all $v \in \mathscr{N}(t)$ with $\mathscr{L} \not\preceq (v,t)$.
2. N^+-BBM: Whenever a particle gets killed, there are at least N particles to its right (but not necessarily all the particles which have N particles to their right get killed). N^--BBM: Whenever at least N particles are to the right of a particle, it gets killed (but possibly more particles get killed).

Let ν_t^+, ν_t^- and ν_t^N be the empirical measures of the particles at time t in N^+-BBM, N^--BBM and N-BBM, respectively. On the space of finite counting measures on $\mathbf{R}$ we denote by $\preceq$ the usual stochastic ordering: For two counting measures ν_1 and ν_2, we write $\nu_1 \preceq \nu_2$ if and only if $\nu_1([x,\infty)) \leqslant \nu_2([x,\infty))$ for every $x \in \mathbf{R}$. If $x_1,\dots,x_n$ and $y_1,\dots,y_m$ denote the atoms of ν_1 and ν_2

respectively, then this is equivalent to the existence of an injective map $\phi : \{1, \ldots, n\} \to \{1, \ldots, m\}$ with $x_i \leqslant y_{\phi(i)}$ for all $i \in \{1, \ldots, n\}$. Furthermore, for two families of counting measures $(\nu_1(t))_{t\geqslant 0}$ and $(\nu_2(t))_{t\geqslant 0}$, we write $(\nu_1(t))_{t\geqslant 0} \preceq (\nu_2(t))_{t\geqslant 0}$ if $\nu_1(t) \preceq \nu_2(t)$ for every $t \geqslant 0$. If $(\nu_1(t))_{t\geqslant 0}$ and $(\nu_2(t))_{t\geqslant 0}$ are random, then we write $(\nu_1(t))_{t\geqslant 0} \overset{\text{st}}{\preceq} (\nu_2(t))_{t\geqslant 0}$ if there exists a coupling between the two (i.e. a realisation of both on the same probability space), such that $(\nu_1(t))_{t\geqslant 0} \preceq (\nu_2(t))_{t\geqslant 0}$.

Lemma 10.1. *Suppose that* $\nu_0^- \overset{\text{st}}{\preceq} \nu_0^N \overset{\text{st}}{\preceq} \nu_0^+$. *Then* $(\nu_t^-)_{t\geqslant 0} \overset{\text{st}}{\preceq} (\nu_t^N)_{t\geqslant 0} \overset{\text{st}}{\preceq} (\nu_t^+)_{t\geqslant 0}$.

Proof. We only prove the second inequality $\nu_t^N \overset{\text{st}}{\preceq} \nu_t^+$, the proof of the first one is similar. By a coupling argument and conditioning on $\mathscr{F}_0$ it is enough to show it for deterministic ν_0^N and ν_0^+. Let $n^+ = \nu_0^+(\mathbf{R})$ and let $\Pi = (\Pi^{(1)}, \Pi^{(2)}, \ldots, \Pi^{(n^+)})$ be a forest of independent BBM trees with the atoms of ν_0^+ as initial positions. We denote by $\mathscr{N}^\Pi(t)$ the set of individuals alive[7] at time t and by $X_u^\Pi(t)$ the position of an individual $u \in \mathscr{N}^\Pi(t)$. Denote by $\mathscr{N}^+(t) \subset \mathscr{N}^\Pi(t)$ the subset of individuals which form the N^+-BBM (i.e. those which have not been killed by the selection mechanism of the N^+-BBM). We set $\nu_t^+ = \sum_{u\in\mathscr{N}^+(t)} \delta_{X_u^\Pi(t)}$.

From the forest Π we will construct a family of forests $\left(\Xi_t = (\Xi_t^{(1)}, \ldots, \Xi_t^{(N)})\right)_{t\geqslant 0}$ (not necessarily comprised of independent BBM trees), such that

- if $t_1 \leqslant t_2$, then the forests Ξ_{t_1} and Ξ_{t_2} agree on the time interval $[0, t_1]$,
- the initial positions in the forest Ξ_0 are the atoms of ν_0^N,
- for every $t \geqslant 0$, the N-BBM is embedded in Ξ_t up to the time t, i.e. for $0 \leqslant s \leqslant t$, if $\mathscr{N}^\Xi(s)$ denotes the set of individuals[8] from Ξ_t alive at time s and $X_u^\Xi(s)$ the position of the individual $u \in \mathscr{N}^\Xi(s)$, then there is a subset $\mathscr{N}^N(s) \subset \mathscr{N}^\Xi(s)$ such that $(\nu_s^N)_{0\leqslant s\leqslant t} = \left(\sum_{u\in\mathscr{N}^N(s)} \delta_{X_u^\Xi(s)}\right)_{0\leqslant s\leqslant t}$ is equal in law to the empirical measure of N-BBM run until time t,
- for every $t \geqslant 0$, there exists a (random) injective map $\phi_t : \mathscr{N}^N(t) \to \mathscr{N}^+(t)$, such that $X_u^\Xi(t) \leqslant X_{\phi_t(u)}^\Pi(t)$ for every $u \in \mathscr{N}^N(t)$.

We will say that the individuals u and $\phi_t(u)$ are *connected*. If at a time t an individual $v \in \mathscr{N}^+(t)$ is not connected to another individual (i.e. $v \notin \phi_t(\mathscr{N}^N(t))$), we say that v is *free*.

The construction of the coupling goes as follows: Since $\nu_0^N \preceq \nu_0^+$, we can construct Ξ_0, $\mathscr{N}^N(0)$ and ϕ_0, such that for every $u \in \mathscr{N}^N(0)$ we have $X_u^\Xi(0) \leqslant X_{\phi_0(u)}^\Pi(0)$ and the subtrees $\Xi_0^{(u,0)}$ and $\Pi^{(\phi_0(u),0)}$ are the same up to translation. We now define a sequence of random times $(t_n)_{n\geqslant 0}$ recursively by $t_0 = 0$ and for each n, we define t_{n+1} to be the first time after t_n at which either

1. a particle of the N-BBM branches, or
2. the left-most particle of the N^+-BBM dies without a particle of the N-BBM branching.

Note that between these times, there is nothing to do, i.e. if Ξ_t and ϕ_t are defined for all $t \leqslant t_n$, then we simply set $\Xi_t = \Xi_{t_n}$ and $\phi_t = \phi_{t_n}$ for all $t \in [t_n, t_{n+1})$. This is clear if no branching occurs between the times t_n and t_{n+1}, but even if a particle of the N^+-BBM branches at a time $t \in (t_n, t_{n+1})$, since no branching occurs in the N-BBM between t_n and t_{n+1}, this particle must be free at the time t and therefore no change in Ξ_t and ϕ_t are necessary. At the times t_n however, we may need to "rewire" particles. We show that this is always possible in such a way that the above conditions on the family Ξ_t are satisfied. Figure 6 is a graphical description of the key step.

Let $n \in \mathbf{N}$ and suppose that

a) Ξ_t and ϕ_t have been defined for all $t < t_{n+1}$ and are equal to Ξ_{t_n} and ϕ_{t_n}, respectively,

b) for each $u \in \mathscr{N}^N(t_n)$, the subtrees $\Xi_{t_n}^{(u,t_n)}$ and $\Pi_{t_n}^{(\phi_{t_n}(u),t_n)}$ are the same, up to translation.

Note that this is the case for $n = 0$. We now distinguish between the two cases above, starting with the second:

---

[7] The term "alive" has the same meaning here as in Section 3.1.

[8] Note that this does not depend on t.



Figure 6: The connection between the particles w of the N-BBM and w' of the N^+-BBM breaks. By definition of the N^+-BBM, there exists a free particle v' to the right of w' and w is rewired to that particle.

Case 2: The left-most particle w' of the N^+-BBM gets killed without a particle of the N-BBM branching. If w' is free, nothing has to be done. Suppose therefore that w' is connected to a particle w of the N-BBM. Then, since there are at most $N-1$ remaining particles in the N-BBM and there are at least N particles to the right of w' in the N^+-BBM (otherwise it would not have been killed), at least one of those particles is free. Denote this particle by v'. We then "rewire" the particle w to v' by setting $\phi_{t_{n+1}}(w) := v'$ and define $\Xi_{t_{n+1}}$ by replacing the subtree $\Xi_{t_n}^{(w,t_{n+1})}$ in Ξ_{t_n} by $\Pi_{t_n}^{(v',t_{n+1})}$, properly translated. Note that we then have

$$X^{\Pi}_{\phi_{t_{n+1}}(w)}(t_{n+1}) = X^{\Pi}_{v'}(t_{n+1}-) \geqslant X^{\Pi}_{w'}(t_{n+1}-) \geqslant X^{\Xi}_{w}(t_{n+1}-),$$

where the first inequality follows from the fact that w' is the left-most individual in N^+-BBM at time t_{n+1} and the second inequality holds by hypothesis.

If more than one particle of the N^+-BBM gets killed at the time t_{n+1}, we repeat the above procedure for every particle, starting from the left-most.

Case 1: A particle u of the N-BBM branches at time t_{n+1}. By the hypothesis b), the particle $\phi_{t_n}(u)$ then branches as well into the same number of children. We then define $\phi_{t_n}(uk) = \phi_{t_n}(u)k$ for each $k \in \{1, \dots, C_u\}$ (recall that C_u denotes the number of children of u), i.e. we connect each child of u to the corresponding child of $\phi_{t_n}(u)$. Now first define $\phi'_{t_{n+1}}$ to be the restriction of ϕ_{t_n} to the surviving particles, i.e. to $\mathscr{N}^N(t_{n+1})$. Then continue as in Case 2, i.e. for each particle w' of the N^+-BBM which gets killed and which is connected through $\phi'_{t_{n+1}}$ to a particle w of the N-BBM, rewire w to a free particle v'. In the end, we get $\phi_{t_{n+1}}$.

In both cases, we then set $\Xi_t = \Xi_{t_{n+1}}$ and $\phi_t = \phi_{t_{n+1}}$ for each $t \in [t_{n+1}, t_{n+2})$, as explained above. Note that each time we are rewiring a particle, we rewire it to a particle whose subtree is independent of the others by the strong branching property, whence the particles from $\mathscr{N}^N(t)$ and $\mathscr{N}^+(t)$ still follow the law of N-BBM and N^+-BBM, respectively. Furthermore, we have for every $\omega \in \Omega$: $\nu_t^N(\omega) \preceq \nu_t^+(\omega)$ for every $t \geqslant 0$. This finishes the proof. □

10.2 Proof of Theorem 1.1

Let $(\nu_t^N)_{t\geqslant 0}$ be the empirical measure process of N-BBM starting from N particles independently distributed according to the density proportional to $\sin(\pi x/a_N)e^{-x}\mathbf{1}_{(x\in(0,a_N))}$, where $a_N = (\log N + 3\log\log N)$. We wish to show that for every $\alpha \in (0,1)$, the finite-dimensional distributions of the process $(M_\alpha^N(t\log^3 N))_{t\geqslant 0}$ (defined in the introduction) converge weakly as $N \to \infty$ to those of the Lévy process $(L_t)_{t\geqslant 0}$ stated in Theorem 1.1 starting from x_α. We will do this by proving seperately a lower and an upper bound and show that in the limit these bounds coincide and equal the Lévy process $(L_t)_{t\geqslant 0}$.

Lower bound. Fix $\alpha \in (0,1)$ and $\delta > 0$. We will let N and in parallel A and a go to infinity (in the meaning of Section 7) in such a way that $N = 2\pi e^{A+\delta} a^{-3} e^{\mu a}$ and such that A goes to infinity sufficiently slowly such that the results from Section 8 hold. We then have with $c = \log(2\pi)$,

$$a = a_N - (A + \delta + c + o(1)), \qquad \text{and} \qquad \mu = \mu_N - \frac{\pi^2}{a^3}(A + \delta + c + o(1)), \tag{10.1}$$

where μ is defined in (5.1). Let $(\nu_t^\flat)_{t\geqslant 0}$ be the empirical measure process of B$^\flat$-BBM starting from the initial configuration (HB$^\flat_\perp$), i.e. $\nu_0^\flat$ is obtained from $\lfloor e^{-\delta}N\rfloor$ particles distributed independently according to the density proportional to $\sin(\pi x/a)e^{-\mu x}\mathbf{1}_{(0,a)}(x)$. An easy calculation now shows that $\nu_0^\flat \overset{\mathrm{st}}{\preceq} \nu_0^N$ for large N, by (10.1). Now, if $X_t^\flat$ denotes the barrier process of the B$^\flat$-BBM, then $(\nu_t^-)_{t\geqslant 0} = (\nu_t^\flat + \mu t + X_t^\flat)_{t\geqslant 0}$ is by definition an instance of the N^--BBM, as defined earlier in this section. Here, for a measure ν and a number x we denote by $\nu + x$ the measure ν translated by x. Lemma 10.1 now gives for large N, $(\nu^-)_{t\geqslant 0} \overset{\mathrm{st}}{\preceq} (\nu_t^N)_{t\geqslant 0}$, which by definition implies

$$(\mathrm{qu}_\alpha^N(\nu_t^N))_{t\geqslant 0} \overset{\mathrm{st}}{\geqslant} (\mathrm{qu}_\alpha^N(\nu_t^-))_{t\geqslant 0}. \tag{10.2}$$

Given $0 \leqslant t_1 < \ldots < t_n$, we now define $t_i^N = a^{-3}t_i \log^3 N$, such that $t_i^N \to t_i$ for every i, as $N \to \infty$. We then have

$$\begin{aligned}\Big(M_\alpha^N\big(t_i \log^3 N\big)\Big)_{i=1,\ldots,n} &\overset{\mathrm{st}}{\geqslant} \big(\,\mathrm{qu}_\alpha^N(\nu^-_{t_i \log^3 N}) - \mu_N t_i \log^3 N\big)_{i=1,\ldots,n} && \text{by (10.2)}\\ &= \big(\,\mathrm{qu}_\alpha^N(\nu^\flat_{a^3 t_i^N}) + X_{a^3 t_i^N} - \pi^2(A+\delta+c+o(1))t_i^N\big)_{i=1,\ldots,n} && \text{by (10.1)}\\ &\overset{\mathrm{st}}{\geqslant} \big(x_{\alpha e^{2\delta}} + L_{t_i} - O(\delta) + c\big)_{i=1,\ldots,n} && \text{for large } N,\end{aligned}$$

where the last inequality follows from Propositions 8.2 and 8.3, with $(L_t)_{t\geqslant 0}$ being the Lévy process from the statement of Theorem 1.1 starting from 0. Letting first $N \to \infty$, then $\delta \to 0$ yields the proof of the lower bound.

Upper bound. The proof is analogous to the previous case, relying on Propositions 9.2 and 9.3 instead of Propositions 8.2 and 8.3. There are only two differences to notice: First, the B$^\sharp$-BBM is not a realisation of the N^+-BBM. However, the C$^\sharp$-BBM (defined in Lemma 9.4) is such a realisation and by that lemma, Propositions 9.2 and 9.3 hold for the C$^\sharp$-BBM as well. Second, if $\nu_0^\sharp$ is distributed according to (HB$^\sharp_\perp$), we do not have $\nu_0^N \overset{\mathrm{st}}{\preceq} \nu_0^\sharp$. However, if $\widetilde{\nu}_0^N$ is obtained from ν_0^N by killing the particles in the interval $[a_N - A^2, a_N]$, then by (10.1), a quick calculation shows that $\widetilde{\nu}_0^N = \nu_0^N$ with high probability and $\widetilde{\nu}_0^N \overset{\mathrm{st}}{\preceq} \nu_0^\sharp$ as N goes to infinity. This finishes the proof of the upper bound and of Theorem 1.1.

Acknowledgments

I wish to thank my PhD advisor Zhan Shi for his continuous support and encouragements and Julien Berestycki for numerous discussions and his interest in this work.

References

[1] D. Aldous (1978). Stopping times and tightness. *The Annals of Probability*, **6**, 2, 335–340

[2] A. Asselah, P. A. Ferrari, P. Groisman, M. Jonckheere (2012). Fleming-Viot selects the minimal quasi-stationary distribution: The Galton-Watson case. arXiv:1206.6114

[3] A. Asselah, M.-N. Thai (2012). A note on the rightmost particle in a Fleming-Viot process. arXiv:1212.4168

[4] K. B. Athreya, P. E. Ney (1972). *Branching processes*, volume 196 of *Die Grundlehren der mathematischen Wissenschaften.* Springer-Verlag, New York

[5] R. Bellman (1961). *A brief introduction to theta functions.* Athena Series: Selected Topics in Mathematics. Holt, Rinehart and Winston, New York

[6] J. Bérard, J.-B. Gouéré (2010). Brunet-Derrida behavior of branching-selection particle systems on the line. *Communications in Mathematical Physics*, **298**, 2, 323–342

[7] J. Berestycki, N. Berestycki, J. Schweinsberg (2010). The genealogy of branching Brownian motion with absorption. arXiv:1001.2337

[8] J. Bertoin, J.-F. Le Gall (2000). The Bolthausen-Sznitman coalescent and the genealogy of continuous-state branching processes. *Probability Theory and Related Fields*, **117**, 2, 249–266

[9] J. D. Biggins, A. E. Kyprianou (2004). Measure change in multitype branching. *Advances in Applied Probability*, **36**, 2, 544–581

[10] P. Billingsley (1999). *Convergence of probability measures.* Wiley Series in Probability and Statistics: Probability and Statistics, second edition. John Wiley & Sons Inc., New York

[11] N. H. Bingham, R. A. Doney (1974). Asymptotic properties of supercritical branching processes I: The Galton-Watson process. *Advances in Applied Probability*, **6**, 4, 711–731

[12] N. H. Bingham, C. M. Goldie, J. L. Teugels (1987). *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications.* Cambridge University Press, Cambridge

[13] A. N. Borodin, P. Salminen (2002). *Handbook of Brownian motion—facts and formulae.* Probability and its Applications, second edition. Birkhäuser Verlag, Basel

[14] M. D. Bramson (1983). Convergence of solutions of the Kolmogorov equation to travelling waves. *Memoirs of the American Mathematical Society*, **44**, 285

[15] E. Brunet, B. Derrida (1997). Shift in the velocity of a front due to a cutoff. *Physical Review E*, **56**, 3, 2597–2604

[16] — (2001). Effect of microscopic noise on front propagation. *Journal of Statistical Physics*, **103**, 1, 269–282

[17] E. Brunet, B. Derrida, A. Mueller, S. Munier (2006). Phenomenological theory giving the full statistics of the position of fluctuating pulled fronts. *Physical Review E*, **73**, 5, 056126

[18] E. Brunet, B. Derrida, A. H. Mueller, S. Munier (2006). Noisy traveling waves: Effect of selection on genealogies. *Europhysics Letters (EPL)*, **76**, 1, 1–7

[19] E. Brunet, I. M. Rouzine, C. O. Wilke (2008). The Stochastic Edge in Adaptive Evolution. *Genetics*, **179**, 1, 603–620

[20] D. Buraczewski (2009). On tails of fixed points of the smoothing transform in the boundary case. *Stochastic Processes and their Applications*, **119**, 11, 3955–3961

[21] K. Burdzy, R. Holyst, D. Ingerman, P. March (1996). Configurational transition in a Fleming–Viot-type model and probabilistic interpretation of Laplacian eigenfunctions. *Journal of Physics A: Mathematical and General*, **29**, 11, 2633–2642

[22] B. Chauvin (1991). Product martingales and stopping lines for branching Brownian motion. *The Annals of Probability*, **19**, 3, 1195–1205

[23] B. Chauvin, A. Rouault (1988). KPP equation and supercritical branching Brownian motion in the subcritical speed area. Application to spatial trees. *Probability Theory and Related Fields*, **80**, 2, 299–314

[24] B. Chauvin, A. Rouault, A. Wakolbinger (1991). Growing conditioned trees. *Stochastic Processes and their Applications*, **39**, 1, 117–130

[25] D. A. Dawson, B. Maisonneuve, J. Spencer (1993). *École d'Été de Probabilités de Saint-Flour XXI—1991*, volume 1541 of *Lecture Notes in Mathematics.* Springer-Verlag, Berlin

[26] B. Derrida, D. Simon (2007). The survival probability of a branching random walk in presence of an absorbing wall. *Europhysics Letters (EPL)*, **78**, 6, 60006

[27] R. Durrett, D. Remenik (2011). Brunet–Derrida particle systems, free boundary problems and Wiener–Hopf equations. *The Annals of Probability*, **39**, 6, 2043–2078

[28] S. N. Ethier, T. G. Kurtz (1986). *Markov processes.* Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons Inc., New York

[29] W. Feller (1971). *An introduction to probability theory and its applications. Vol II.* Second edition. John Wiley & Sons Inc., New York

[30] N. Gantert, Y. Hu, Z. Shi (2011). Asymptotics for the survival probability in a killed branching random walk. *Annales de l'Institut Henri Poincaré (B) Probabilités et Statistiques*, **47**, 1, 111–129

[31] R. Hardy, S. C. Harris (2006). A new formulation of the spine approach to branching diffusions. arXiv:math/0611054

[32] J. W. Harris, S. C. Harris, A. E. Kyprianou (2006). Further probabilistic analysis of the Fisher-Kolmogorov-Petrovskii-Piscounov equation: One sided travelling-waves. *Annales de l'Institut Henri Poincaré (B) Probabilités et Statistiques*, **42**, 1, 125–145

[33] S. C. Harris, M. I. Roberts (2011). The many-to-few lemma and multiple spines. arXiv:1106.4761

[34] T. E. Harris (1963). *The theory of branching processes*, volume 119 of *Die Grundlehren der Mathematischen Wissenschaften.* Springer-Verlag, Berlin

[35] N. Ikeda, M. Nagasawa, S. Watanabe (1968-69). Branching Markov processes. I–III. *Journal of Mathematics of Kyoto University.* I: 8, 233–278, 1968. II: 8, 365–410, 1968. III: 9, 95-160

[36] K. Itô, H. P. McKean Jr. (1974). *Diffusion processes and their sample paths*, volume 125 of *Die Grundlehren der mathematischen Wissenschaften.* Second edition. Springer-Verlag, Berlin

[37] P. Jagers (1989). General branching processes as Markov fields. *Stochastic Processes and their Applications*, **32**, 2, 183–212

[38] O. Kallenberg (1997). *Foundations of modern probability.* Probability and its Applications (New York). Springer-Verlag, New York

[39] H. Kesten (1978). Branching Brownian motion with absorption. *Stochastic Processes and their Applications*, **7**, 1, 9–47

[40] F. B. Knight (1969). Brownian local times and taboo processes. *Transactions of the American Mathematical Society*, **143**, 173–185

[41] T. G. Kurtz (1975). Semigroups of conditioned shifts and approximation of Markov processes. *The Annals of Probability*, **3**, 4, 618–642

[42] A. E. Kyprianou (2004). Travelling wave solutions to the K-P-P equation: Alternatives to Simon Harris' probabilistic analysis. *Annales de l'Institut Henri Poincaré (B) Probabilités et Statistiques*, **40**, 1, 53–72

[43] A. Lambert (2000). Completely asymmetric Lévy processes confined in a finite interval. *Annales de l'Institut Henri Poincaré (B) Probabilités et Statistiques*, **36**, 2, 251–274

[44] Q. Liu (2000). On generalized multiplicative cascades. *Stochastic Processes and their Applications*, **86**, 2, 263–286

[45] R. Lyons, R. Pemantle, Y. Peres (1995). Conceptual proofs of L Log L criteria for mean behavior of branching processes. *The Annals of Probability*, **23**, 3, 1125–1138

[46] P. Maillard (2010). The number of absorbed individuals in branching Brownian motion with a barrier. arXiv:1004.1426, to appear in *Annales de l'Institut Henri Poincaré (B) Probabilités et Statistiques.*

[47] — (2012). *Branching Browian motion with selection.* Ph.D. thesis, Université Pierre et Marie Curie, arXiv:1210.3500

[48] H. P. McKean (1975). Application of Brownian motion to the equation of Kolmogorov-Petrovskii-Piskunov. *Communications on Pure and Applied Mathematics*, **28**, 3, 323–331

[49] C. Mueller, L. Mytnik, J. Quastel (2010). Effect of noise on front propagation in reaction-diffusion equations of KPP type. *Inventiones mathematicae*, **184**, 2, 405–453

[50] J. Neveu (1986). Arbres et processus de Galton-Watson. *Annales de l'Institut Henri Poincaré (B) Probabilités et Statistiques*, **22**, 2, 199–207

[51] — (1988). Multiplicative martingales for spatial branching processes. In *Seminar on Stochastic Processes, 1987 (Princeton, NJ, 1987)*, volume 15 of *Progress in Probability and Statistics*, 223–242. Birkhäuser Boston, Boston, MA

[52] D. Panja (2004). Effects of fluctuations on propagating fronts. *Physics Reports*, **393**, 2, 87–174

[53] D. Revuz, M. Yor (1999). *Continuous martingales and Brownian motion*, volume 293 of *Die Grundlehren der Mathematischen Wissenschaften.* Third edition. Springer-Verlag, Berlin

[54] S. A. Sawyer (1976). Branching diffusion processes in population genetics. *Advances in Applied Probability*, **8**, 4, 659–689

[55] T. Shiga (1988). Stepping stone models in population genetics and population dynamics. In *Stochastic processes in physics and engineering (Bielefeld, 1986)*, volume 42 of *Mathematics and Applications*, 345–355. Reidel, Dordrecht

[56] W. Van Saarloos (2003). Front propagation into unstable states. *Physics Reports*, **386**, 2-6, 29–222

[57] S. Watanabe (1967). Limit theorem for a class of branching processes. In *Markov Processes and Potential Theory (Proc. Sympos. Math. Res. Center, Madison, Wis., 1967)*, 205–232. Wiley, New York

[58] W. Whitt (2002). *Stochastic-process limits.* Springer Series in Operations Research. Springer-Verlag, New York

[59] T. Yang, Y.-X. Ren (2011). Limit theorem for derivative martingale at criticality w.r.t branching Brownian motion. *Statistics & Probability Letters*, **81**, 2, 195–200